

E-20% w/v Oral Solution

E-20 w/v Oral Solution appears to be a yellow colour, clear solution.

Composition:

Each liter contains

Enrofloxacin	200 g
Benzyl Alcohol (as Preservative)	10 g

Pharmacodynamics:

Enrofloxacin inhibits bacterial DNA gyrase or topoisomerase IV by non-covalent binding to an intermediate state during which DNA is cleaved, thereby preventing DNA replication. The inhibition of DNA and mRNA synthesis triggers events resulting in a rapid, drug concentration-dependent killing of pathogenic bacteria.

Enrofloxacin is active against many Gram-negative bacteria, against Gram-positive bacteria and *Mycoplasma* spp.

Pharmacokinetics:

Enrofloxacin administered via drinking water to poultry is rapidly and very well absorbed with high bioavailability. Enrofloxacin possess very good tissue penetration with higher concentrations found in target tissues like lungs, liver, kidney, intestine and muscle tissue than the serum concentration. In poultry enrofloxacin is poorly metabolized to its active metabolite ciprofloxacin. Enrofloxacin is eliminated from the body at a half-life of 6 hours.

Indications:

Treatment of infections caused by the following bacteria susceptible to enrofloxacin:

Chickens: *Mycoplasma gallisepticum*, *Mycoplasma synoviae*, *Avibacterium paragallinarum*, *Pasteurella multocida*, *Escherichia coli*.

Turkeys: *Mycoplasma gallisepticum*, *Mycoplasma synoviae*, *Pasteurella multocida*, *Escherichia coli*.

Rabbits: For the treatment of infectious diseases due to *Pasteurella multocida* and bacterial enteritis due to infection with *E. coli*.

Enrofloxacin should be used where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the active substance of choice.

Recommended Dose:

Chickens and turkeys

10 mg enrofloxacin/kg bodyweight per day for 3–5 consecutive days. Treatment for 3–5 consecutive days; for 5 consecutive days in mixed infections and chronic progressive forms.

Rabbits

10 mg/kg bodyweight per day for 5 consecutive days.

If no clinical improvement is achieved within 2–3 days, alternative antimicrobial therapy should be considered based on susceptibility testing.

Mode of Administration

Oral administration.

Contraindication:

Do not use for prophylaxis.

Do not use when resistance / cross-resistance to (fluoro)quinolones is known to occur in the flock intended for treatment.

Do not use in case of known hypersensitivity to the active substance, other (fluoro)quinolones or to any of the excipients.

Warning and precautions:

Special warnings for each target species

Treatment of *Mycoplasma* spp infections may not eradicate the organism.

Special precautions for use

i) Special precautions for use in animals

Official and local antimicrobial policies should be taken into account when the product is used.

Fluoroquinolones should be reserved for the treatment of clinical conditions which have responded poorly, or are expected to respond poorly, to other classes of antimicrobials. Whenever possible fluoroquinolones should only be used based on susceptibility testing.

ii) Special precautions to be taken by the person administering the veterinary medicinal product to animals

Those with known hypersensitivity to (fluoro)quinolones should avoid contact with this product.

Avoid contact with skin and eyes.

Rinse any splashes from skin or eyes immediately with water.

Wash hands and exposed skin after use.

Do not eat drink or smoke whilst handling the product.

TO BE PRESCRIBED BY REGISTERED VETERINARY SURGEONS ONLY.

Interactions with Other Medicinal Product:

In vitro, an antagonism was shown, when combining fluoroquinolones with bacteriostatic antimicrobial agents such as macrolides or tetracyclines and phenicols. The simultaneous application of substances containing aluminium or magnesium can impair the absorption of enrofloxacin.

Statement on Usage During Pregnancy and Lactation:

Not authorised for use in birds producing eggs for human consumption.

Do not administer to layer replacement birds within 14 days of coming into lay.

Adverse Effects / Undesirable Effects:

None known.

Overdose and Treatment:

No adverse clinical symptoms were observed in chickens and turkeys treated respectively with doses up to 10 and 6 times higher than the therapy dose. The use of fluoroquinolones during the growth phase combined with a marked and prolonged increase in the intake of drinking water, and hence active ingredient, possibly due to high temperatures, may potentially be associated with damage of the articular cartilage.

Withdrawal Period (s):

Chickens: Meat and offal: 7 days

Turkeys: Meat and offal: 13 days

Rabbits: Meat and offal: 15 days.

Not authorised for use in birds producing eggs for human consumption.

Do not administer to layer replacement birds within 14 days of coming into lay.

Storage Condition:

Store below 30°C. Protect from heat and sunlight exposure.

Keep medicine out of reach of children/Jauhkan ubat dari kanak-kanak.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Shelf-life after first opening: 7 days

Shelf-life after reconstitution or dilution: 24 hours

Packaging Available: 1 L

Date of revision: 05-06-2017

Product Holder/ Manufacturer:

NAM PHARMA SDN. BHD.

5, Lebuhr Perusahaan Klebang 11,
Taman Perindustrian Antarabangsa IGB,
31200 Chemor, Perak, Malaysia.

Tel: +605-2931838, 1839 Fax: +605-2931833