

LINCOMYN 110MG/G SOLUBLE POWDER

FOR ANIMAL USE ONLY

Brand or Product Name :

LINCOMYN 110MG/G SOLUBLE POWDER

Name and Strength of Active Substance(s)

Each gram of LINCOMYN 110MG/G SOLUBLE POWDER contains Lincomycin 110 mg which is equivalent to Lincomycin Hydrochloride 124.74 mg.

Product Description

A white or almost white powder.

Target Species

Swine and chicken (broiler).

Pharmacodynamics

Lincomycin is a lincosamide antibiotic produced by *Streptomyces lincolensis*. Lincomycin is bacteriostatic in action inhibiting the protein synthesis predominantly by binding to the 50S ribosomal subunits of bacteria. Depending on the sensitivity of micro-organisms, and on the concentration of the active substance the protein synthesis inhibition antibacterial action can either be bacteriostatic or bactericide. Lincomycin is active against a wide range of Gram-positive micro-organisms, including also anaerobic bacteria such as *Clostridia* and *Brachyspira* spp., as well as *Mycoplasma* spp. Lincomycin has no activity against Gram-negative bacteria, such as *Klebsiella* spp., *Pasteurella* spp. and *Salmonella* spp. The resistance rate against lincomycin is slow, multiple-step type. Plasmid mediated infectious resistance is also described. No cross-resistance has been described with penicillin, ampicillin, cephalosporins, tetracyclines, or novobiocin.

Pharmacokinetics

Lincomycin is rapidly absorbed orally. After oral administration, the absorbed lincomycin is eliminated through the bile and faeces in active form or as metabolites. *Lincomycin* is also excreted in the milk. Lincomycin reaches inflammation site by polymorph neutrophil granulocytes which explains its fast absorption and distribution, efficient penetration and targeted activity in difficult to reach tissues.

Indications

Treatment of infection caused by mycoplasma and bacteria susceptible to lincomycin for example:

Pigs: swine dysentery caused by *Brachyspira hyodysenteriae*, mycoplasma pneumonia associated with *Mycoplasma hyopneumoniae*.

Chicken: necrotic enteritis caused by *Clostridium perfringens*.

Route of Administration

For use in drinking water.

Dosage and Administration

To be administered in drinking water.

Indication	Treatment (mg/kg bodyweight)		Duration of Treatment (day)
	Product	Lincomycin	
Swine dysentery	91	10	10 days "or at least 5 days after clinical signs disappear".
Swine Mycoplasma pneumonia	91	10	21 days.
Poultry necrotic enteritis caused by <i>Clostridium perfringens</i>	45.5	5	7 days.

Administration:

The product should be administered continuously in the drinking water.

Swine: Dissolve 300 g of product in 1000 litres of water to provides 33 mg lincomycin per litre. When using a water proportioner, prepare a stock solution by dissolving 300 g of product in 10 litres of water. Set the proportioner to deliver 10 ml of stock solution per litre of drinking water.

Chicken (broiler): The concentration to be used depends on the actual body weight and the water consumption of the animals and can be calculated as follows:

$$\frac{\text{....mg product}}{\text{per kg body weight and day}} \times \frac{\text{Average chicken body weight (kg)}}{\text{Average daily water intake (litre/animal)}} = \text{....mg product per litre of drinking water}$$

To ensure a correct dosage body weight should be determined as accurately as possible. The intake of medicated water depends on the clinical condition of the animals. In order to obtain the correct dosage the concentration of Lincomycin has to be adjusted accordingly. The medicated water should be the only source of drinking water, throughout the treatment period. Medicated water should be refreshed every 24 hours.

Contraindications

Do not use in case of hypersensitivity to the active substance or to the excipient. Do not use if resistance to lincosamides has been detected. Do not administer to horses, ruminants, guinea pigs, hamsters, chinchilla and rabbits. Ingestion by these species may result in severe gastrointestinal effects.

Warning and Precaution

Special warnings for each target species: None

i. Special precautions for use in animals: Medicated drinking water uptake can be affected by the severity of the disease. In case of insufficient uptake of water, animals should be treated parenterally. Use of the product should be based on susceptibility testing and take into account official and local antimicrobial policies. Use of the product deviating from the instructions given may increase the prevalence of bacteria resistant to lincomycin and may decrease the effectiveness of treatment with other lincosamides, macrolides or streptogramins due to potential for cross-resistance.

ii. Special precautions to be taken by the person administering the veterinary medicinal product to animals: People with known hypersensitivity to lincomycin should avoid contact with the veterinary medicinal product. Care should be taken not to inhale any dust. The wearing of approved dust masks, gloves and safety glasses is recommended whilst reconstituting or administering the solution. Direct contact of the product with the skin, eyes and mucous membranes should be avoided. In case of accidental exposure rinse abundantly with water. In case of allergic reaction (inflammation of the face, lips or eyes or respiratory difficulties) during reconstituting or administering of the product, seek medical advice immediately and show the package leaflet or the label to the physician. Do not smoke, eat or drink when handling the product. Wash hands and any exposed skin with soap and water immediately after use.

iii. Other precautions: When spreading the manure of weaner pigs treated with the veterinary medicinal product at a dose of 10 mg/kg for 21 days, a minimum distance to surface water of 10 m should be applied.

Drug Interactions

Do not use together with macrolide antibiotics.

Pregnancy and Lactation

The safety of the veterinary medicinal product has not been established in pregnant or lactating sows. Use only in accordance with benefit/risk assessment by the responsible veterinarian. Do not use in birds in lay or within 4 weeks before the onset of the laying period.

Adverse Effects

Lincomycin may occasionally cause transient soft stools and/or mild swelling of the anus within the first two days of treatment. Very rarely some pigs may show reddening of skin and mild irritable behaviour. These conditions are usually self-correcting within 5-8 days without discontinuing lincomycin treatment.

Overdose and Treatment

Lincomycin has a good margin of safety, but higher levels of dosage than recommended may cause diarrhoea and loose stools in pigs.

Withdrawal Period(s)

Meat and offal of swine: 0 day

Meat and offal of chicken (broiler): 5 days

Not authorised for use in laying birds producing eggs for human consumption.

Shelf-life

Shelf life as packaged for sale: 3 year

Shelf life after first opening : 24 hours

Shelf life after reconstitution : 24 hours

Maximum Residual Limit (MRL)

Lincomycin

Pig : Muscle 200 µg/kg, Liver 500 µg/kg, Kidney 1500 µg/kg, Fat 100 µg/kg, Skin with adhering fat 300 µg/kg

Chicken : Muscle =200 µg/kg, Liver = 500 µg/kg, Kidney = 500 µg/kg, Fat = 100 µg/kg, Skin with adhering fat = 300 µg/kg [Ref. CODEX ALIMENTARIUS : CAC/MRL 2-2014]

Storage Conditions

Store in a dry place below 30°C. Keep out of reach of children. Jauhi ubat daripada kanak-kanak.

Any opened sachet or unused reconstituted water must be discarded after 24 hours.

Disposal of Containers

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Dosage Forms and Packaging Available

100g, 200g, 500g, 1000g powder in aluminum bag.

Name and Address of Manufacturer

Life Biopharma Sdn. Bhd. (Company No. 546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2017-02-28