

# Feniflor 100 mg/ml Oral Solution

(FOR VETERINARY USE ONLY)

## COMPOSITION

Florfenicol .....100g  
Excipients q.s.....1000ml

## PRODUCT DESCRIPTION

The product is a light yellow clear solution.

## PHARMACODYNAMICS

Florfenicol is a bacteriostatic antibiotic that inhibits protein synthesis by binding to ribosomal subunits of susceptible bacteria, leading to the inhibition of peptidyl transferase and thereby preventing the transfer of amino acids to growing peptide chains and subsequent protein formation. The bacterial receptor that is the site of action for florfenicol is considered to be the same as that for chloramphenicol and thiamphenicol.

## PHARMACOKINETICS

After administration to pigs by gavage at 15 mg/kg under experimental conditions, absorption of florfenicol was variable but peak serum concentrations of approximately 5 µg/ml were reached approximately 2 hours after dosing. The terminal half-life was between 2 and 3 hours. When pigs were given free access, for 5 days, to water medicated with florfenicol at a concentration of 100 mg per litre of water, serum concentrations of florfenicol exceeded 1 µg/mL for the entire 5 day treatment period except for a couple of short excursions below 1 µg/mL. After absorption and distribution, florfenicol is extensively metabolised by pigs and rapidly eliminated, primarily in urine. After parenteral dosing of florfenicol to pigs, it has been shown that lung concentrations are similar to serum concentrations.

For broiler chickens, serum concentrations versus time in chickens after the administration of florfenicol in drinking water at an expected concentration of 100ppm for 18 hours per day over 3 days.

## INDICATIONS

For the treatment of swine respiratory disease associated with *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.  
In broiler chicken for the treatment and control of air sacculitis associated with *E. coli* susceptible to florfenicol.

## CONTRA-INDICATIONS

The medical contraindications have been selected on the basis of the potential significance. This medication should not be used if there is previous allergic or toxic reaction to florfenicol.

## MODE OF ADMINISTRATION

Oral Administration through drinking water

## RECOMMENDED DOSE:

For broiler:

1ml/L of Feniflor or 100 ppm (100 mg/L) for 3 consecutive days in drinking water. This dosage equivalent to 35 mg/kg body weight.

For swine:

1ml/L of Feniflor or 100 ppm (100 mg/L) for 5 consecutive days in drinking water. This dosage rate equivalent to 10 mg/kg body weight for pigs that are drinking 10% of their body weight.

## WARNING AND PRECAUTIONS

Automatic water proportions should not be used if water hardness is greater than 275 parts per million. Galvanized metal watering systems should not be used and chlorinator should not be operated while the medication is being administered.

Keep medication out of reach of children.

## INTERACTION WITH OTHER MEDICAMENTS

No data available.

## PREGNANCY AND LACTATION

For Swine: Not for use in breeding swine. The effects of florfenicol on porcine reproductive performance, pregnancy and lactation have not been assessed.

For Broiler Chickens: Do not feed to birds intended for breeding purposes. The effects of the product on the reproductive function of treated chickens have not been determined.

## ADVERSE EFFECT

For Swine: perianal inflammation may be observed in approximately 40% of the treated pigs. Rectal eversion may occur in about 6% of the treated pigs. These effects should resolve completely within one week. Transient diarrhea may occur with the use of this drug.

For Broiler Chickens: None known.

## OVERDOSE AND TREATMENT

In swine, case of overdosing results in a decrease in weight gain, food and water consumption, peri-anal erythema and oedema and modification of some haematological and biochemical parameters indicative of dehydration may be observed.

In poultry and swine, stop medication and seek for veterinarian advice if there is sign of overdose.

## WITHDRAWAL PERIODS

Swine: 21days

Broiler chicken: 5 days

## STORAGE CONDITION

Store in a cool and dry room ( $\leq 30^{\circ}\text{C}$ ). Keep away from direct sunlight

After first opening : Use within 30 days

After dilution : Finished product is stable after 7days

**Date of Revision** : 27<sup>th</sup> May 2017

**Mal no.** :

**Packaging** : 1 L

**Exp date** :

**Mfg date** :

## Manufactured By:

**Nam Pharma Sdn. Bhd.**

No. 5, Lebuhr Perusahan Klebang 11,  
Taman Perindustrian Antarabangsa IBG,  
31200 Chemor, Perak, Malaysia.

## Marketing By:

**F.E Venture Sdn. Bhd.**

Lot 3 & 5, Jalan PJS 11/8, Bandar Sunway,  
46150 Petaling Jaya, Selangor D.E., Malaysia