

Toltrastop 25 mg/ml Oral Solution

(FOR VETERINARY USE ONLY)

COMPOSITION

Toltrazuril25g
Excipients q.s.1000ml

PRODUCT DESCRIPTION

The product is a clear, colourless to yellowish solution.

PHARMACODYNAMICS

Toltrazuril acts as an anticoccidial, active against all intracellular stages but the mode of action is unknown.

PHARMACOKINETICS

In poultry toltrazuril is absorbed at a rate of at least 50%. The highest equivalent concentrations are found in liver. The active substance is rapidly metabolised in poultry. The main metabolite is characterised as a sulfone-derivative. About 1 week after the last dose this metabolite represents by far the most relevant residue in the animal.

INDICATIONS

For the treatment of Coccidiosis in Broilers and Broiler Breeders. Toltrastop 25 mg/ml Oral Solution is effective against *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. necatrix*, *E. tenella*, *E. mitis*.

CONTRA-INDICATIONS

None

MODE OF ADMINISTRATION

Oral Administration via drinking water

RECOMMENDED DOSE:

The recommend dose rate is 7 mg Toltrazuril per kg bodyweight per day given for 2 consecutive days. This corresponds to:
- 28 ml Toltrastop 25mg/ml oral Solution (equivalent to 700 mg toltrazuril) per 100 kg of bodyweight per day for 2 consecutive days.

As a guide for usage this is normally equivalent to 25 ppm (equivalent to 1 ml Toltrastop 25mg/ml) per litre of drinking water for continuous medication over 48 hours, or to 75 ppm (equivalent to 3 ml Toltrastop 25mg/ml) per litre of drinking water given for 8 hours per day, on 2 consecutive days.

INTERACTION WITH OTHER MEDICAMENTS

None known

Toltrastop 25 mg/ml Oral Solution is compatible with coccidiostats and amoebostats.

PREGNANCY AND LACTATION

None known

ADVERSE EFFECT

None known

OVERDOSE AND TREATMENT

A three - five fold overdose is readily tolerated without any symptoms. If the recommended dose is exceeded beyond 3 - 5 times there is a decrease in water intake.

WITHDRAWAL PERIODS

Chicken meat: 18 days.

Do not use in birds producing eggs for human consumption.

TO BE PRESCRIBED AND TREATED WITH BY REGISTERED VETERINARY SURGEONS ONLY

WARNING AND PRECAUTIONS

Special warnings for each target species

Toltrastop does not interfere with the development of immunity against coccidiosis. As with all anticoccidials, prolonged use may result in the development of resistant strains.

Special precautions for use

i. Special precautions for use in animals

Dilutions more concentrated than 1:1,000 (1 ml Toltrastop 25mg/ml to 1 litre drinking water) may result in precipitation. Predilution is not recommended.

ii. Special precautions to be taken by the person administering the medicinal product to animals

- Toltrastop is an alkaline solution.
- Wear synthetic rubber gloves when handling the product.
- Wash any splashes from skin or eyes immediately with water.
- Wash hands and exposed skin after use.
- Do not eat, drink or smoke whilst using the product.

iii. Other precautions

- Only make up sufficient drinking water for requirements.
- Wash out empty container with water 3 times when medicating birds and add washings to drinking water.
- Store litter from treated houses as long as possible. However, it should only be stored where there is no risk of polluting surface water or groundwater.
- Do not apply to agricultural land at a rate greater than 1.5 tonnes/hectare/year.
- If spreading within an NVZ ensure that the NVZ action programme rules are complied with.
- The principle metabolite of toltrazuril is persistent and mobile in groundwater.

STORAGE CONDITION

Store below 30°C. Protect from sunlight and heat exposure.

Keep medication out of reach of children
(Jauhkan ubat dari kanak-kanak)

Shelf life after first opening : 30 days

Shelf life after dilution : 7 days

Shelf life for sale : 3 years

Date of Revision : 12th April 2017

Mal no. : MAL20172950HAC

Packaging : 1 L

Exp date :

Mfg date :

Manufactured By:

Nam Pharma Sdn. Bhd.

No. 5, Lebuhr Perusahan Klebang 11,
Taman Perindustrian Antarabangsa IBG,
31200 Chemor, Perak, Malaysia.

Marketing By:

F.E Venture Sdn. Bhd.

Lot 3 & 5, Jalan PJS 11/8, Bandar Sunway,
46150 Petaling Jaya, Selangor D.E., Malaysia