

Pharmasin 100 % w/w Granules for Use in Drinking Water

for Pigs, Chickens, Turkeys and Calves

1. MANUFACTURER OF THE VETERINARY MEDICINAL PRODUCT

BIOVET Joint Stock Company
39 Petar Rakov Street, 4550 Peshtera, Bulgaria

2. PRODUCT REGISTRATION HOLDER NAME AND ADDRESS

Pahang Pharmacy Sdn. Bhd. (10826-K)
Lot 5979 Jalan Teratai, 5½ Mile Off Jalan Meru,
41050 Klang, Selangor Darul Ehsan, Malaysia

3. NAME OF THE VETERINARY MEDICINAL PRODUCT

Pharmasin 100 % w/w Granules for Use in Drinking Water for Pigs, Chickens, Turkeys and Calves
Tylosin (as Tylosin tartrate)

4. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

1.1 g of granules contains
1 g of tylosin (corresponding to 1.1 g of tylosin tartrate)

5. DOSAGE FORM

Granules for use in drinking water

6. PRODUCT DESCRIPTION

Pharmasin 100% w/w represent white to light yellow coloured granules intended for use in drinking water.

7. INDICATION(S)

Calves: Treatment and prevention of pneumonia caused by *Mycoplasma spp* when the disease has been established in the herd

Pigs:

- Treatment and prevention of enzootic pneumonia caused by *Mycoplasma hyopneumoniae* and *Mycoplasma hyorhinis* when the disease has been established in the herd.

- Treatment and prevention of Porcine Intestinal Adenomatosis (Ileitis) associated with *Lawsonia intracellularis* when the disease has been established in the herd.

Chickens:

- Treatment and prevention of chronic respiratory diseases (CRD) caused by *Mycoplasma gallisepticum* and *Mycoplasma synoviae* when the disease has been established in the flock

- Treatment and prevention of necrotic enteritis caused by *Clostridium perfringens* when the disease has been established in the flock

Turkeys: Treatment and prevention of infectious sinusitis caused by *Mycoplasma gallisepticum*, when the disease has been established in the flock.

8. CONTRAINDICATIONS

Do not use in animals with known hypersensitivity to tylosin or other macrolides,

Do not use in cases with known resistance to tylosin or cross-resistance to other macrolides (MLS-resistance).

Do not use in animals vaccinated with tylosin-sensitive vaccines either at the same time or within 1 week previously.

Do not use in animals with hepatic disorders.

Do not use in horses because of danger for

inflammation of the caecum.

9. ADVERSE REACTIONS

In pigs, adverse reactions have been observed, including diarrhea, pruritus, erythema of the skin, swelling of the vulva, rectal edema and prolapse. These reversible signs appeared 48-72 hours after start of treatment.

10. TARGET SPECIES

Calves, pigs, chickens, turkeys

11. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Oral administration through the drinking water

In calves the product can also be administered through milk or milk replacer.

1.1 gram of the veterinary medicinal product corresponds to 1 gram of tylosin. The dosages are as follows:

Calves:

10 - 20 mg tylosin per kg BW (corresponding to 11 - 22 mg of the veterinary medicinal product per kg BW), twice daily (corresponding to a daily dose of 20 - 40 mg tylosin per kg BW), for 7 - 14 days.

Turkeys:

75 - 100 mg tylosin per kg BW per day (corresponding to 82.5 - 110 mg of the veterinary medicinal product per kg BW) for 3 - 5 days.

Chickens:

For the treatment of chronic respiratory disease:
75 - 100 mg tylosin per kg BW per day (corresponding to 82.5 - 110 mg of the veterinary medicinal product per kg BW) for 3 - 5 days.

For the treatment of necrotic enteritis:

20 mg tylosin per kg BW per day (corresponding to 22 mg of the veterinary medicinal product) for 3 days.

Pigs:

For the treatment of enzootic pneumonia:
20 mg tylosin per kg BW per day (corresponding to 22 mg of the veterinary medicinal product per kg BW) for 10 days.

For the treatment of ileitis or PLA:

5 - 10 mg tylosin per kg BW per day (corresponding to 5.5 - 11 mg of the veterinary medicinal product per kg BW) for 7 days.

For the preparation of the medicated water/milk/milk-replacer the body weight of the animals to be treated and their actual daily water/milk/milk-replacer consumption should be taken into due account. Consumption may vary depending on factors like age, state of health, breed, husbandry system. To provide the required amount of active substance in mg per litre drinking water/milk/milk-replacer the following calculation should be made:

$$\frac{\text{..... mg tylosin per kg bodyweight per day}}{\text{Average amount of drinking water or milk / animal (L)}} \times \frac{\text{Average bodyweight (kg) of the animals to be treated}}{\text{L of drinking water}} = \text{.....mg tylosin / L of drinking water}$$

If individual animals show signs of a serious infection such as a reduced water or feed intake, then they should be treated individually, such as by injection.

12. WITHDRAWAL PERIOD

Calves (meat and offal): 12 days

Pigs (meat and offal): 1 day

Turkeys (meat and offal): 2 days

Turkey (eggs): Zero days

Chickens (meat and offal): 1 day

Chicken (eggs): Zero days

13. PHARMACODYNAMICS

Tylosin is a macrolide antibiotic produced by a strain of *Streptomyces fradiae*. It exerts its antimicrobial effect by inhibiting protein synthesis of susceptible micro-organisms.

The tylosin spectrum of activity includes amongst others Gram-positive bacteria, some Gram - negative strains such as *Pasteurella*, and *Mycoplasma spp*.

14. PHARMACOKINETICS

In most species peak plasma concentrations have been attained 1 to 2 hours after administration of tylosin. Compared to plasma levels clearly higher tissue concentrations have been observed. Tylosin was extensively metabolised.

15. SPECIAL STORAGE PRECAUTIONS

Store in the original container in order to protect from light. Store below 30°C.

Shelf-life after dilution or reconstitution according to directions:

medicated water: 24 hours

medicated milk or milk replacer: 24 hours

Shelf-life after first opening the immediate

packaging: 3 months

16. SPECIAL WARNING(S)

Due to likely variability (time, geographical) in susceptibility of bacteria to tylosin, bacteriological sampling and susceptibility testing are recommended.

Under-dosing and/or treating for an insufficient length of time are considered to promote the development of resistance in bacteria and should be avoided.

Interactions

Lincosamides and aminoglycoside antibiotics antagonize the activity of tylosin.

Incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

Special precautions for use in animals

Animals with acute infections may have a reduced water and feed intake and should be treated with a suitable injectable veterinary medicinal product first.

Do not leave or dispose of water containing tylosin

may occasionally be serious and therefore direct contact should be avoided.

To avoid exposure during preparation of the medicated drinking water, wear overalls, safety glasses, impervious gloves and wear either a disposable half mask respirator conforming to European Standard EN149 or a non-disposable respirator to European Standard EN140 with a filter to EN143.

Wash hands after use.

In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water.

Do not handle the product if you are allergic to ingredients in the product.

If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Use during pregnancy, lactation or lay

Laboratory studies in mice and rats have not produced any evidence of teratogenic, foetotoxic or maternotoxic effects. No studies have been conducted in the target species. Use only according to the benefit/risk assessment by the responsible veterinarian.

Overdose

There is no evidence of tylosin toxicity in rats, at dose rates of up to 1000 mg/kg by the oral route. There is no evidence of tylosin toxicity in chickens, turkeys, pigs or calves when administered orally at up to three times the recommended dose.

17. PACKAGING

1.1 kg resealable block bottom zipped sachet made of polyethylene /aluminium/polyethylene terephthalate laminate.

110 g high density polyethylene pot with polypropylene cap.

18. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

19. DATE OF REVISION

March 2017



tartrate where it may be accessible to either animals not under treatment or wildlife.

User warnings

Tylosin may induce irritation. Macrolides, such as tylosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tylosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances