

# LINCOCEN 44MG/G PREMIX POWDER

**Composition:**

Each gm contains:

Lincomycin (as lincomycin hydrochloride).....44mg

**Product Description:**

Lincocen 44mg/g Premix Powder is an off white powder.

**Pharmacodynamics:**

Lincomycin is a lincosamide antibiotic and is active against a wide range of Gram-positive aerobic and anaerobic bacteria, anaerobic Gram-negative bacteria and mycoplasma. Lincosamides are generally bacteriostatic.

**Pharmacokinetics:**

Lincomycin has approximately 50% systemic bioavailability from oral administration in pigs and is significantly re-circulated via the biliary route.

**Indications:****Broilers**

For the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to lincomycin in broilers.

**Mode of Administration:**

Orally. Mixed into feed.

**Recommended Dosage:**

To be administered orally via the feed.

**Broilers:**

For the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to lincomycin in broilers

2 grams per ton of feed.

**Contraindications:**

Do not administer to animals with known hypersensitivity to the active ingredient.

Not for use in animals other than the target species. Lincomycin may cause fatal enterocolitis in horses, rodents and rabbits.

**Warnings and Precautions:****Special warnings for each target species**

None.

**Special precautions for use in animals**

Not applicable.

**Special precautions to be taken by the person administering the veterinary medicinal product to animals**

Persons known to be hypersensitive (allergic) to lincomycin and/or soybean millfeed should not handle this product.

Care should be taken not to inhale any dust. The wearing of approved dust masks (either a disposable half-mask respirator conforming to European Standard EN 149 or a non-disposable respirator to European Standard EN 140 with a filter to EN 143) and safety glasses is recommended during the handling and mixing of this product.

Skin contact should be avoided.

Wash hands and any exposed skin with soap and water immediately after use.

**Interactions with Other Medicaments:**

Clinical incompatibility may exist between lincomycin and erythromycin due to competitive binding at the ribosomal site of action.

**Pregnancy and Lactation:**

Not applicable.

**Adverse Effects/Undesirable Effects:**

May occasionally cause transient soft stools and/or mild swelling of the anus within the first two days of treatment. Very rarely some pigs may show reddening of skin and mild irritable behaviour. These conditions are usually self-correcting within 5-8 days without discontinuing lincomycin treatment.

**Overdose and Treatment:**

Not applicable.

**Withdrawal Period(s):**

Broilers:

No drug withdrawal period is required before slaughter of birds fed LINCOCEN 44MG/G PREMIX POWDER.

Not for use in layers and breeders.

**Storage conditions:**

Store in a cool dry place, protected from light. Do not store above 30°C.

Keep out of reach of children/Jauhkan dari kanak-kanak.

**Pack size(s):**

Paper bag of 10kg.

**Product Registration Holder/ Manufacturer**

Osmosis Nutrition Sdn. Bhd. (594876A)  
Lot 16014(P.T. 24341), Jalan Nilam 3,  
Bandar Nilai Utama,  
71800 Nilai, Negeri Sembilan, Malaysia.  
Tel: +606-799 9090 Fax: +606-799 9155

**Date of revision of Package Insert:**

19/06/2017