

DIMECTAL Powder for Oral Suspension 3g

COMPOSITION

Each 3.9g sachet contains:

Diosmectite 3g

PRODUCT DESCRIPTION

Appearance of powder:

Creamy grey powder

Appearance of Suspension:

Creamy grey suspension having orange flavour.

PHARMACODYNAMIC

DIMECTAL retains a powerful coating property on the gastrointestinal mucosa due to its leaflet structure and high plastic viscosity. DIMECTAL increases the resistance of the mucosal gel in response to aggressive agents and by interacting with glycoproteins of mucus. By its action on the gastrointestinal mucous barrier and its high-binding capacity, DIMECTAL protects the gastrointestinal mucosa. DIMECTAL is radiolucent, does not colour the stools and, at usual doses, does not modify the physiological intestinal transit time.

DIMECTAL is an anti-diarrhoeal that absorbs and flushes out toxins, bacteria, viruses and parasites from the intestinal wall as well as excess water in the gut to reduce loose and watery stools. This product also repairs, regenerates, and heals the intestinal mucus lining which has been damaged by diarrhoea. By creating a barrier on the intestinal lining, it protects the intestines against toxins, bacteria, viruses and parasites, and reduces gut inflammation. Hence, the peristaltic movement of the gastrointestinal (GI) tract (including the bowels) is normalised - restoring and regulating a healthy gut environment, conducive for intestinal microflora to recolonise and populate.

PHARMACOKINETIC

DIMECTAL is neither absorbed nor metabolized. It is eliminated through the faeces.

INDICATION

- Treatment of acute diarrhoea in adults and children above 2 years, in addition to oral rehydration
- Symptomatic treatment of chronic functional diarrhoea in adults.
- Symptomatic treatment of pain associated with functional bowel diseases in adults.

RECOMMENDED DOSAGE

Treatment of Acute Diarrhoea

Children:

In children from 2 years old:

4 sachets a day for 3 days, then 2 sachets per day. (Up to maximum 7 days)

Adults: Average of 3 sachets per day.

In practice, the daily dose can be doubled at the start of treatment. (Up to maximum 7 days)

Other Indications:

Adults: Average of 3 sachets per day.

DIMECTAL Powder for Oral Suspension 3g

Method of Administration:

Children: The contents of sachet can be added and thoroughly stirred in the feeding bottle with 50ml of water, to be given during the day, or thoroughly mixed with a semi liquid food: broth, stewed fruit, mashed vegetables or baby food.

Adults: Contents of one sachet should be thoroughly stirred in half a glass of water before taking.

ROUTE OF ADMINISTRATION

Oral

CONTRAINDICATIONS

Not to administer to children under 2 years of age.

WARNING AND PRECAUTIONS

DIMECTAL is not a rehydration therapy. A proper rehydration should always be used in case of dehydration.

In children above 2 years, should be used only in the treatment of acute diarrhoea (7 days maximum), in association with early administration of an oral rehydration solution (ORS). Any chronic use should be avoided.

In adult: Prolonged or repeated use is not recommended without medical advice.

Consult your doctor or pharmacist:

- If you have an intolerance to particular sugars(s).

This medicine contains glucose and sucrose. Its use is not recommended in patients with fructose intolerance, glucose or galactose malabsorption syndrome or sucrose/ isomaltase deficiency (rare hereditary disease).

- In case of history of severe constipation.

- In case of acute diarrhoea, if your symptoms do not improve or worsen within 3 days of treatment.

- If your digestive pain is associated with fever and vomiting.

PREGNANCY AND LACTATION

DIMECTAL is not recommended during pregnancy and lactation

If you are pregnant, think you might be pregnant or are planning to become pregnant or are breast feeding, you should seek the advice from your doctor or pharmacist before taking this medicine.

DRUG INTERACTIONS

As the adsorbent properties of DIMECTAL may interfere with the rates and/or levels of absorption of other substances, it is recommended not to administer any other drugs at the same time as DIMECTAL.

ADVERSE EFFECTS

Very rare cases of development or aggravation of constipation have been reported; treatment was able to be continued after decreasing the dosage.

DIMECTAL Powder for Oral Suspension 3g

OVERDOSGE AND TREATMENT

Taking too much DIMECTAL may cause constipation. The constipation usually disappears when treatment is stopped or when the dose is reduced. Ask your doctor or pharmacist.

STORAGE CONDITIONS

Store below 30°C.

PRESENTATION

Powder for Oral Suspension, 3.9g per sachet. 10, 12, 15, 20, 30, 60 and 100 sachets per pack

MANUFACTURED BY

Noripharma Sdn Bhd. 200701034604 (792633-A)
Lot 5030 Jalan Teratai,
5½ Mile Off Jalan Meru,
41050 Klang, Selangor,
Malaysia.

PRODUCT REGISTRATION HOLDER

Pahang Pharmacy Sdn Bhd. 197101000448 (10826-K)
Lot 5979 Jalan Teratai,
5½ Mile Off Jalan Meru,
41050 Klang, Selangor,
Malaysia.

DATE OF REVISION

April 2024