

AVICOL (FLORFENICOL 15% W/V ORAL SOLUTION)

CONTROLLED MEDICINE

Composition:

Each 1L contains:

Florfenicol.....150 gm

Product description:

Solution for use in Drinking Water.

Clear, colourless to yellow solution.

Target species:

Poultry (Chicken) and Swine (pig)

Pharmacokinetics

Swine:

After administration to pigs by gavage at 15 mg/kg under experimental conditions, Florfenicol was detectable in the serum of all animals through 12 hours after gavage. After gavage of florfenicol on the fifth day, the maximum serum florfenicol concentrations were between 2.78 µg/ml and 7.34 µg/ml and the median terminal half life was 2.0 hours. The extent of bioavailability ranged from 24% to 97%.

Poultry:

Serum concentrations versus time in chickens after the administration of florfenicol in drinking water at an expected concentration of 100 ppm for 18 hours per day over 3 days.

Pharmacodynamics

Swine:

Florfenicol is a synthetic broad spectrum antibiotic active against many gram-negative and gram-positive bacteria isolated from domestic animals. It acts by inhibiting bacterial protein synthesis. In vivo activity has been demonstrated against bacterial pathogens commonly associated with swine respiratory disease, including *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.

Poultry :

Florfenicol is a synthetic broad spectrum antibiotic active against many gram-negative and gram-positive bacteria isolated from domestic animals. It acts by inhibiting bacterial protein synthesis.

Indication:

Treatment of bacterial diseases susceptible to florfenicol.

Chicken:

Effective against species: *E.Coli*

Indication: Colibacillosis

Pig:

Effective against species: *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Streptococcus suis*, *Mycoplasma Hyopneumoniae*

Indication: Pleuropneumonia, Pneumonic pasteurellosis, Streptococcal infection, *Mycoplasma pneumonia*

Dosage and Administration

Oral administration through drinking water.

Chicken:

0.13 ml (equivalent to 20 mg florfenicol) per kg body weight for 5 days.

Pig:

0.067 ml (equivalent to 10 mg florfenicol) per kg body weight for 5 days

Contraindication:

Do not use in case of hypersensitivity to the active substances or excipient

Warning and Precautions:

- a. Do not administer to the following sick animals
 - Do not use on animals that have had shock and hypersensitivity to this drug and oxacephem based antibiotics.
- b. General precautions
 - In continuous administration, temporary inflammation may occur in the cloaca of chicken and the anus of pigs.
- c. Application precautions
 - 1. If mixed with drinking water, mix evenly. If it is not homogenously mixed, it may result in weakening and can't achieve the purpose of administration.
 - 2. Do not use chlorinated water during administration.
 - 3. Do not use galvanized water containers.
 - 4. Precipitation may occur if other dilution ratios are used.

Interactions with other medications

No data available

Pregnancy and lactation:

The effect of florfenicol on porcine reproductive performance, pregnancy and lactation have not been assessed.

The effect of florfenicol on the reproductive functions of treated chickens have not been determined.

Adverse Effect:

In continuous administration, temporary inflammation may occur in the cloaca of the chickens and the anus of pigs.

Overdose and Treatment:

In case of overdosing, a decrease in weight gain, water intake, peri-anal erythema and oedema and modification of some haematological and biochemical parameters indicative of dehydration may be observed

Withdrawal period:

Chicken : 2 days

Pigs : 16 days

Do not use in layer chicken

Proposed Shelf Life

Shelf-life of the veterinary medicinal product as packaged for sale: 12 months

Shelf-life after first opening the immediate packaging: 28 days

Shelf-life after reconstitution or dilution: The solution is discarded after 24 hours

Storage Condition Storage below 30°C

Protect from light and moisture.

For veterinary use only

Keep away from children

Jauhi daripada kanak-kanak

BATCH NUMBER**MANUFACTURED DATE****EXPIRY DATE****MAL NUMBER****MARKET AUTHORIZATION HOLDER**

FIJILIAM ENTERPRISE SDN.BHD

100 & 101, Taman A.S.T, Jalan Sungai Ujong

70200 Seremban, Negeri Sembilan,
Malaysia

MANUFACTURER:

Nam Pharma Sdn Bhd

5, Lebuhr Perusahaan

Klebang 11, Taman
Perindustrian

Antarabangsa IGB

31200,

CHEMOR PERAK

MALAYSIA

Date of Revision: January 2021