

BAPID 100 mg Tablet

ANTI-GASTRITIS AND ANTI-GASTRIC ULCER DRUG

Each film coated tablet contains:

Rebamipide 100 mg◇ **PRODUCT DESCRIPTION:**

White, round, biconvex, bevelled on both sides, film coated tablet with engraved REB on one side and 100 on the other

◇ **MECHANISM OF ACTION:****Pharmacology**Prostaglandin-increasing effect

The drug has the effect of increasing the PGE2 content in the gastric mucosa and protected the gastric mucosa from injury caused by Ethanol loading.

Cytoprotective effect

The drug inhibits gastric mucosal injury induced by Aspirin, Ethanol, or HCl-ethanol loading. Effect on inflammatory cytokine release (interleukin-8) in the gastric mucosa

Rebamipide, taken by the oral route, suppressed the increase production of interleukin-8 in the mucosa of patients with *Helicobacter pylori*. The drug also inhibited the activation of NF-κB and suppressed the expression of interleukin-8 mRNA in epithelial cells cocultured with *Helicobacter pylori* (*in vitro*).

PharmacokineticsPlasma Concentration

The table below shows the pharmacokinetic parameters of Rebamipide following single oral administration of Rebamipide tablets at the dose 100 mg in a fasted state.

Pharmacokinetic Parameters of Rebamipide

	t _{max} (hr)	C _{max} (ng/mL)	t _{1/2} (hr)	AUC _{24h} (ng•hr/mL)
Rebamipide 100 mg Tablet	2.4 ± 1.2	216 ± 79	1.9 ± 0.7	874 ± 209

Mean value ± SD, n = 27, t_{1/2} calculated from values up to 12 hr.

The absorption of Rebamipide following single oral administration at a dose of 150 in a fed state tended to be slower than that in a fasted stage. However, food did not affect bioavailability of the drug in humans. Pharmacokinetic parameters obtained from patients with renal impairment after single oral administration of Rebamipide at 100 mg revealed higher plasma concentrations and a longer elimination half-life compared with those in healthy subjects. At steady-state, Rebamipide plasma concentrations observed in dialyzed renal patients following repeated administration were very close to the values simulated from single administration. Therefore, the drug was not considered to accumulate.

-1-

Metabolism

Rebamipide was primarily excreted as the unchanged compound in the urine after single oral administration to healthy adult males at a dose of 600 mg. A metabolite with a hydroxyl group at the 8th position was identified in the urine.

However, the excretion of this metabolite was only 0.03% of the administered dose. The enzyme involved in the formation of the metabolite was CYP3A4.

The usual dosage in adults is 100 mg three times daily.

Excretion

Approximately 10% of the administered dose was excreted in the urine when Rebamipide was administered as a single oral dose to healthy adult males at 100 mg.

Protein Binding

Rebamipide at 0.05-5 µg/mL was added to human plasma, *in vitro*, and 98.4%-98.6% of the drug was bound to plasma proteins.

◇ **INDICATION:**

- Gastric ulcers
- Treatment of gastric mucosal lesions (erosion, bleeding, redness, and edema) in the following conditions; acute gastritis and acute exacerbation of chronic gastritis.

◇ **DOSAGE AND ADMINISTRATION:**

Oral

Gastric ulcers

The usual adult dosage of Rebamipide is one tablet (100 mg of Rebamipide) taken by the oral route three times daily, in the morning, in the evening, and before at bedtime.

Treatment of gastric mucosal lesions (erosion, bleeding, redness, and edema) in the following conditions; acute gastritis and acute exacerbation of chronic gastritis

The usual adult dosage of Rebamipide is one tablet (100 mg of Rebamipide) three times daily taken by the oral route.

◇ **PRECAUTION:**Elderly

Special care is required in elderly patients to minimize the risk of gastrointestinal disorders, because these patients may be physiologically more sensitive to this drug than younger patients.

Pediatric use

The safety of this drug in children has not been established.

Patient's Instruction for Use

However, patients should be instructed not ingest any portion of the blister pack. There have been reports that the sharp edges of the sheet can cut or penetrate the esophageal mucosa if accidentally ingested, resulting in mediastinitis or other serious complications.

◇ **PREGNANCY AND LACTATION:****Pregnancy**

This drug should be administered to pregnant or possibly pregnant women only if the anticipated therapeutic benefit is thought to outweigh any potential risk. The safety of this drug in pregnant women has not been established.

Breastfeeding

Nursing should be interrupted when this drug is administered to.

-2-

PM Specification	Product	Code No.	Dimension	Packaging Type	Thickness
	BAPID 100	IBMY 0051	W 15 x L 20 cm.	Wood Free Paper (กระดาษปลอดไม้)	60 g (0.08 mm.)
Designed by: (PDM, ASPD, PDC-M, PDC-A, PDC-M, PDC-A)					
Checked by: (PDM/ASPD)					
Approved by: (MD (Only Pattern and Colors))					
Approved by: (Date)	PLC: (Date)	LRA: (Date)	IRA: (Date)	QCC-PM: (Date)	CUSTOMER/SALES DEPT.: (Date)

BAPID 100_Insert_Front,Back_IBMY 0051 By_DA 12/05/2023

❖ **SIDE EFFECT:**

Clinically significant Side Effects/

- Shock or anaphylactoid reactions may occur. Patients should therefore be closely monitored. If abnormal findings are observed, the drug should be discontinued and appropriate measures taken.
- Leukopenia and thrombocytopenia may occur. Patients should therefore be closely monitored. If abnormal findings are observed, the drug should be discontinued and appropriate measures taken.
- Hepatic dysfunction and jaundice, as indicated by increases in AST (GOT), ALT (GPT), γ -GTP, and alkaline phosphatase levels, have been reported in patients receiving Rebamipide tablets. If abnormal laboratory findings are observed, the drug should be discontinued and appropriate measures taken.

Other Side Effects

System Organ Class	Adverse Reaction
Hypersensitivity ¹	Rash, pruritus, drug-eruption-like eczema, and other symptoms of hypersensitivity
	Urticaria
Neuropsychiatric	Numbness, dizziness, and sleepiness
Gastrointestinal	Constipation, feeling of abdomen enlarged, diarrhea, nausea, vomiting, heartburn, abdominal pain, belching, taste abnormality, etc.
	Thirst
Hepatic ²	Increased AST (GOT), ALT (GPT), γ -GTP and alkaline phosphate levels
Hematologic	Leukopenia, granulocytopenia, etc.
	Thrombocytopenia
Other	Menstrual disorders, increased BUN levels, edema, and feeling of a foreign body in the pharynx
	Breast swelling and pain, gynecomastia, induction of lactation, palpitation, fever, facial flushing, numbness of tongue, cough, respiratory distress, and alopecia

¹ If such symptoms of hypersensitivity occur, the drug should be discontinued.

² If transaminase levels are markedly increased or fever and rash develop, the drug should be discontinued and appropriate measures should be taken.

❖ **EFFECTS ON ABILITY TO DRIVE AND USE MACHINE:**

No specific studies have been conducted to assess the direct effect of Rebamipide on the ability to drive and use machines.

❖ **CONTRAINDICATION:**

Patients with a history of hypersensitivity to any ingredient of this drug.

❖ **DRUG INTERACTION:**

Rebamipide had little inhibitory effect on cytochrome P450 enzymes *in vitro*. Rebamipide is hardly metabolized in the liver and excreted in urine as unchanged compound; thus, the drug has not much influence on metabolism of other drugs. No drug interactions between Rebamipide and other drugs are reported.

❖ **OVERDOSE AND TREATMENT:**

There is limited information on overdose. Treatment of overdose should be symptomatic. Close medical supervision and monitoring should continue until the patient recovering.

❖ **STORAGE:**

Store at temperature of not more than 30°C.

❖ **DOSAGE FORM AND PACKAGING AVAILABLE:**

100 mg Tablet, Blister 10x10's

❖ **DATE OF REVISION:**

May 10, 2023

Manufactured by:
UNISON LABORATORIES CO., LTD.
39 Moo 4, Klong Udomcholjorn, Muang Chachoengsao,
Chachoengsao 24000 Thailand

C-MY/100523-04 (AR)
(Mucosta Tab 100 PI – MY, 7/18; ¹1st corres. 19/8/20)

IBMY 0051

Product	Code No.	Dimension	Packaging Type	Thickness
BAPID 100	IBMY 0051	W 15 x L 20 cm.	Wood Free Paper (กระดาษปลอดไม้)	60 g (0.08 mm.)
Designed by: (PDM, ASPD, PDC-M, PDC-A, PDC-M, PDC-A) Checked by: (PDM/ASPD) Approved by: (MD (Only Pattern and Colors))				
Approved by: PLC: (Date) LRA: (Date) IRA: (Date) QCC-PM: (Date) CUSTOMER/SALES DEPT.: (Date)				