

SPACE FOR
PHARMACODE

SPACE FOR
PHARMACODE

SPACE FOR
PHARMACODE

PE00000 MY

® Registered Trade Mark

PHARMACODE:

Co-existent tinea pedis should be treated with an appropriate antimycotic cream.

Older people

There are no specific dosage recommendations for use in elderly patients.

Paediatric population

There are no specific dosage recommendations for children owing to the lack of clinical experience available to date.

Contraindications

Hypersensitivity to the active substance or to any of the excipients listed under section List of excipients

Special Warnings and Precautions for Use

Amorolfine 5% Nail Lacquer should not be applied on the skin around the nail.

Avoid contact of the lacquer with eyes, ears and mucous membranes.

Owing to the lack of clinical experience available to date, children should not be treated with amorolfine 5% nail lacquer.

During the application of amorolfine no artificial nails shall be used.

After applying amorolfine 5% nail lacquer, an interval of at least 10 min should be respected before application of any cosmetic nail lacquer.

Before repeat application of amorolfine 5% nail lacquer, the cosmetic nail lacquer should be removed carefully.

When organic solvents are used impermeable gloves shall be used otherwise amorolfine nail lacquer will be removed.

A systemic or local allergic reaction could possibly occur after use of this product. If this happens, the product should be stopped immediately and medical advice should be sought.

Remove the product carefully by using a nail remover solution. The product should not be reapplied.

Interaction with other medicinal products and other forms of interaction

No interaction studies have been performed.

Use of artificial nails should be avoided during treatment.

Fertility, Pregnancy and lactation

Experience with amorolfine use during pregnancy and/or lactation is limited. Only a few cases of exposure to topical amorolfine use in pregnant women have been reported in the post- authorisation setting, therefore the potential risk is unknown. Studies in animals have shown reproductive toxicity at high oral doses; it is unknown whether amorolfine is excreted in human milk. Amorolfine should not be used during pregnancy and/or lactation unless clearly necessary.

Effects on ability to drive and use machines

Not relevant

Undesirable effects

Adverse drug reactions are rare. Nail disorders (e.g. nail discoloration, broken nails, brittle nails) may occur. These reactions can also be linked to the onychomycosis itself.

System Organ Class	Frequency	Adverse drug reaction
Immune system disorders	Unknown frequency*	Hypersensitivity (systemic allergic reaction)*

Skin and subcutaneous tissue disorders	Rare ($\geq 1/10000$, $< 1/1000$)	Nail disorder, nail discoloration, onychoclasia (broken nails), onychorrhexis (brittle nails)
	Very rare ($< 1/10000$)	Skin burning sensation
	Unknown frequency*	Erythema*, pruritus*, contact dermatitis*, urticaria*, blister*

*post-marketing experience

Reporting of suspected adverse reactions

Reporting suspected adverse reaction after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions.

Overdose

No systemic signs of overdose are expected following topical application of amorolfine 5% nail lacquer.

In case of accidental oral ingestion, appropriate symptomatic measures should be taken if needed.

PHARMACOLOGICAL PROPERTIES

Pharmacodynamic properties

Pharmacotherapeutic group: Other antifungals for topical use
ATC code: D01AE16

Mechanism of action

Amorolfine nail lacquer is a topical antimycotic. Amorolfine belongs to a new chemical class, and its fungicidal action is based on an alteration of the fungal cell membrane targeted primarily on sterol biosynthesis. The ergosterol content is reduced, and at the same time unusual sterically nonplanar sterols accumulate.

Microbiology

Amorolfine is a broad spectrum antimycotic. It is highly active (MIC < 2mcg/ml) *in vitro* against

yeasts:	<i>Candida</i> , <i>Cryptococcus</i> , <i>Malassezia</i>
dermatophytes:	<i>Trichophyton</i> , <i>Microsporum</i> , <i>Epidermophyton</i>
moulds:	<i>Hendersonula</i> , <i>Alternaria</i> , <i>Scopulariopsis</i>
dematiacea:	<i>Cladosporium</i> , <i>Fonsecaea</i> , <i>Wangiella</i>
dimorphic fungi:	<i>Coccidioides</i> , <i>Histoplasma</i> , <i>Sporothrix</i>

With the exception of *Actinomyces*, bacteria are not sensitive to amorolfine. *Propionibacterium acnes* is only slightly sensitive.

Pharmacokinetic properties

Amorolfine from nail lacquer penetrates into and diffuses through the nail plate and is thus able to eradicate poorly accessible fungi in the nail bed.

Systemic absorption of the active ingredient is very low with this type of application. Following prolonged use of Amorolfine Nail Lacquer, there is no indication of drug accumulation in the body.

Preclinical safety data

None stated.

PHARMACEUTICAL PARTICULARS

List of excipients

Absolute Alcohol (Ethanol 99% v/v), N-butyl Acetate Extra Pure, Ethyl Acetate, Ammonio Methacrylate Copolymer (Type A), Triacetin.

PE00000 MY

ICONGRAPHICS CODE:

PANTONE SHADE
 PANTONE BLACK PROCESS C
Supersedes Artwork Code

PHARMACODE: