

Dated 24th of April, 2021

PACKAGE INSERT

ZURIL EMV 2.5% w/v Oral Solution

FOR TREATMENT POULTRY COCCIDIOSIS

**TO BE PRESCRIBED AND TREATED WITH BY REGISTERED VETERINARY
SURGEONS ONLY**

a) Brand Name: ZURIL EMV 2.5% w/v Oral Solution

Active ingredient name: Toltrazuril base

Strength of active ingredient:

Each ml contains:

Toltrazuril.....25 mg/ml

Preservatives:

Propylene glycol.... 2.00 % w/v

Methylparaben.... 0.18 % w/v

Propylparaben.... 0.02 % w/v

Dosage form: Oral solution for diluting in drinking water

Product description: colorless to yellowish solution with characteristic odor.

b) Indication:

For the treatment of coccidiosis in chickens and turkeys, caused by infections with various species of *Eimeria*:

Chickens: *E. acervulina*, *E. brunetti*, *E. maxima*, *E. mitis*, *E. necatrix*, *E. tenella*.

Turkeys: *E. adenoides* and *E. meleagrimitis*.

c) **Amounts to be administered and administration route:**

For oral administration via the drinking water

In order to ensure administration of a correct dose, accurately determine the body weight of the animals.

The dose is 7 mg toltrazuril per kg body weight (bw) per day (equivalent to 0.28 ml of product per kg bw per day). Treatment is carried out on two consecutive days.

The medicinal product should be administered either continuously over 24 hours or for one period of 8 hours per day – for 2 consecutive days.

Medicated drinking water should be refreshed every 24 hours.

The dosage should be based on the current, actual drinking water intake of the birds, because this varies depending on the bird species, on the age, state of health and intended use of the birds, and depending on the housing conditions (e.g. different ambient temperature, different lighting regime).

In the case of continuous treatment over 24 hours, the volume of the product to be mixed into the drinking water for the animals to be treated is calculated according to the following formula:

Volume of the product required per liter drinking water:

$$\frac{0.28 \text{ ml product per kg bw per day} \times \text{Average bw (kg) of the birds to be treated}}{\text{Average drinking water intake in liters over 24 hours per bird}} = \text{x ml of product per liter drinking water}$$

Total volume of the product required per day (24 h):

The calculated volume (x ml of product per liter) must be multiplied by the total consumption of drinking water per day (24 h).

In the case of treatment for a period of 8 hours per day, the volume of the product to be mixed into the drinking water for the animals to be treated is calculated according to the following formula:

Volume of the product required per liter drinking water:

$$\frac{0.28 \text{ ml product per kg bw per day} \times \text{Average bw (kg) of the birds to be treated}}{\text{Average drinking water intake in liters over 8 hours per bird}} = \text{y ml of product per liter drinking water}$$

Total volume of the product required for a treatment period of 8 hours:

The calculated volume (y ml of product of per liter) must be multiplied by the total consumption of drinking water per 8-hour period.

The appropriate volume of solution must be added daily to the drinking water while stirring.

If volumes of between 1 and 4 ml product are added per liter drinking water, solubility is guaranteed over the period of treatment.

In order to ensure that all the animals drink water evenly, sufficient space must be made available at the waterer. Free-range animals must be kept indoors during treatment.

After the end of the treatment, the watering system must be cleaned in an appropriate manner in order to prevent any exposure to residual subtherapeutic doses, particularly if liable to promote the development of resistance.

Predilution and the administration through a dosing pump (proportioner) are not recommended. Use preferably a bulk tank.

d) Contraindication:

Do not use in cases of known hypersensitivity to the active substance or any of the excipients.

e) Pharmacodynamics:

Toltrazuril is an anticoccidial of the triazinetrione group, active against *Eimeria* spp. Toltrazuril induces changes in the fine structure of the developmental stages of coccidia. These are caused primarily by swelling of the endoplasmic reticulum and of the Golgi apparatus, abnormal changes to the perinuclear space and disturbances in cell division. Toltrazuril causes a decrease in the activity of respiratory chain enzymes in the parasites

f) Pharmacokinetics:

After oral administration, toltrazuril undergoes at least 50% absorption in poultry. The highest concentrations are to be found in the liver and kidneys of the poultry. The active substance is broken down rapidly. The main metabolite is toltrazuril sulfone.

g) Environmental Properties: Not applicable

h) Target species:

Chickens (broilers, pullets and breeders), Turkeys

i) Route of Administration: by orally.

j) Warnings and precautions:

Special warnings for each target species

Good hygiene can reduce the risk of coccidiosis. It is therefore recommended any deficiencies in husbandry should be addressed in addition to treatment. Poultry houses should be kept clean and dry. It is recommended to treat all animals in a pen. For best results, treatment should be initiated before the clinical signs of disease have spread throughout the whole group

Special precautions for use

Special precautions for use in animals

As with any anticoccidials, frequent and prolonged use of an antiprotozoal of the same class may result in the development of resistances. It is important to keep to the recommended dose in order to minimise the risk of resistance.

If resistance is present it should be considered to use another antiprotozoal from another class/mechanism of action.

This VMP should not be used together with feed additives/or other veterinary medicinal products that might interfere with the efficacy of the product, like "coccidiostats" and "histomonostats".

For best results, treatment should be initiated before the clinical signs of disease have spread throughout the whole group.

The veterinary medicinal product is a strongly alkaline solution and should not be administered undiluted.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

People with known hypersensitivity to toltrazuril should avoid contact with the veterinary medicinal product.

The veterinary medicinal product is an alkaline solution.

Wear synthetic rubber gloves when handling the product.

Contact with skin and mucous membranes should be avoided.

In case of direct contact with the eyes or skin, wash immediately and thoroughly with water.

Do not ingest. In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Do not eat, drink or smoke during use.

Wash hands after use.

k) Interaction of other medicaments:

Combination of the product with antibiotics may result in reduced water intake in turkeys. The concomitant administration of other substances to the drinking water should be avoided

l) Pregnancy and lactation or lay

Not applicable, see Withdrawal periods.

m) Side effects

None known

n) Symptoms and Treatment of overdose:

A reduction in drinking water intake may be the first sign of an overdose. This is observed only after an overdose with more than 10 times the recommended dose.

o) Storage condition:

Store at temperature below 30°C.

Once opened, the medicinal product should be stored at temperatures below 25 °C in the original packaging with tightly closed cap

p) Shelf life:

The product packaged for sales has a shelf-life of 2 years.

Shelf-life after first opening the immediate packaging: 1 month

Shelf-life after dilution or reconstitution according to directions: 24 hours

q) Withdrawal Period:

Chickens:

Meat and offal: 16 days

Turkeys:

Meat and offal: 16 days

Eggs: Not authorised for use in laying birds producing eggs for human consumption. Do not use within 4 weeks of onset of laying.

r) Disposal container

All Hazardous chemicals must be disposed of in accordance with the Environmental Quality (Scheduled Waste) Regulation 2005 and by the authorized waste management companies.

s) Manufacturer:

Emivest Feedmill Vietnam Company Limited

Address: No. A-11C-CN Bau Bang Industrial Park, Lai Uyen Ward, Bau Bang District, Binh Duong Province, Vietnam

t) Importer and Distributor:

F.E VENTURE Sdn Bhd

Lot 3 & 5, Jalan PJS 11/8, Bandar Sunway, 46150 Petaling Jaya, Selangor Darul Ehsan, Malaysia.