

I-Dectin 20mg Chewable Tablet

I-Dectin 20mg Chewable Tablet appears to be a white to off-white flat beveled edge round tablet with a score line on one side.

Composition:

Each chewable tablet of 500 mg contains
Ivermectin

20 mg

Pharmacodynamics:

Ivermectin is a macrocyclic-lactone derivative which has a broad antiparasitic activity against nematodes and arthropods. It acts by inhibiting nerve impulses. Its mode of action includes the glutamate-gated chloride ion channels. Ivermectin binds selectively and with high affinity to glutamate-gated chloride ion channels which occur in invertebrate nerve and muscle cells. This leads to an increase in the permeability of the cell membrane to chloride ions with hyperpolarisation of the nerve or muscle cell, resulting in paralysis and death of the relevant parasites. Compounds of this class may also interact with other ligand-gated chloride channels, such as those gated by the neurotransmitter gamma-aminobutyric acid (GABA). The high margin of safety for compounds of this class is attributable to the fact that, in mammals, glutamate receptors in chloride ion channels do not occur; the macrocyclic lactones have a low affinity to other mammalian ligand-gated chloride ion channels and they do not readily cross the blood-brain barrier.

Pharmacokinetics:

The terminal half-life varies between 2 and 5 days after various routes of administrations. Ivermectin is a poorly metabolised compound. Due to its lipophilic nature, ivermectin is excreted in bile and ultimately eliminated from the body via the faeces. In horses, about 75% of the administered dose is excreted via the faeces after an oral administration of ivermectin at the recommended dose. Moreover 90% of the total drug is excreted within 4 days post-administration. Approximately 2% of unchanged ivermectin and metabolites are excreted in urine.

Indications:

For the treatment of nematode and arthropod infestations, due to adult and immature roundworms and bots in horses:

• **Nematodes**

Large-strongyles:

Strongylus vulgaris (adult and arterial larvae)
Strongylus edentatus (adult and L4 tissue larval stages)
Strongylus equinus (adult and L4 larval stage)
Triodontophorus spp. (adult)

Small-strongyles:

Cyathostomum (adult and non-encysted mucosal larvae): Cylicocyclus spp.,
Cylicostephanus spp., Gyalocephalus spp.
Parascaris: Parascaris equorum (adult and larvae).
Oxyuris: Oxyuris equi (adult and larvae).
Trichostrongylus: Trichostrongylus axei (adult).

• **Dipteran insects:** Gasterophilus spp. (larvae).

Recommended Dosage:

Single oral administration.

200 µg of ivermectin per kg of bodyweight corresponding to 1 tablet per 100 kg bodyweight.

Weight	Dosage	Weight	Dosage
Up to 100 kg	1 tablet	401-500 kg	5 tablets
101-200 kg	2 tablets	501-600 kg	6 tablets
201-300 kg	3 tablets	601-700 kg	7 tablets
301-400 kg	4 tablets	701-800 kg	8 tablets

Warning and Precautions:

Special warnings for each target species

Care should be taken to avoid the following practices because they increase the risk of development of resistance and could ultimately result in ineffective therapy:

- too frequent and repeated use of anthelmintics from the same class over an extended period of time,
- underdosing, which may be due to underestimation of body weight, misadministration of the product, or lack of calibration of the dosing device (if any).

Suspected clinical cases of resistance to anthelmintics should be further investigated using appropriate tests (e.g. Faecal Egg Count Reduction Test). Where the results of the test(s) strongly suggest resistance to a particular anthelmintic, an anthelmintic belonging to another pharmacological class and having a different mode of action should be used.

Resistance to ivermectin has been reported in *Parascaris equorum* in horses. Therefore the use of this product should be based on local (regional, farm) epidemiological information about susceptibility of nematodes and recommendations on how to limit further selection for resistance to anthelmintics.

The product is safe for use in stallions.

Young foals, miniature horses and toy breeds weighing less than 50 kg may be unable to ingest tablets. Seek the advice of your veterinary surgeon.

Special precautions for use in animals

The product has been formulated for use in horses only. Cats, dogs (especially Collies, Old English Sheepdogs and related breeds or crosses) and also turtles and tortoises may be adversely affected by the concentration of ivermectin in this product if they are allowed to ingest spilled tablets or have access to used packaging.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Wash hands after use.

Avoid contact with the eyes. In the event of accidental contact with the eyes, rinse immediately with plenty of water. In case of eye irritation, seek medical attention. Do not eat, drink or smoke while handling this product. In the event of accidental ingestion, seek medical advice and show the package leaflet to the physician.

Interactions with Other Medicaments:

None known.

Use during pregnancy, lactation or lay:

Can be used during pregnancy and lactation.

Side effects:

Colic, diarrhoea and anorexia have been reported in very rare occasions post treatment, in particular when there is heavy worm burden. In very rare occasions, allergic reactions such as hypersalivation, lingual oedema, urticaria, tachycardia, congested mucus membranes, and subcutaneous oedema have been reported following treatment with the product.

Overdose and Treatment:

A tolerance study performed in adult horses with doses up to 5 times the recommended dosage did not show any adverse reactions.

Administration to mares at 3 times the recommended dosage at 14-day intervals during the whole gestation and lactation periods did not result in any abortion, nor any adverse effect during gestation, at parturition or on the mares general health, nor any abnormality in the foals.

To ensure a correct dosage, body weight should be determined as accurately as possible.

Once the correct dose has been determined, it should be administered in the following way :

Present the tablet in the palm of your hand.

Presenting one tablet at a time makes it easier for the horse to accept it, however the administration of multiple tablets at once is also possible.

Repeat this gesture until the complete dose has been administered.

During the initial administration, the tablet can be combined with a small amount of food or a treat to increase the acceptance by the horse.

In the event that the required dose is not ingested an alternative treatment should be administered. Seek the advice of your veterinary surgeon.

The veterinary surgeon should give advice regarding appropriate dosing programmes and stock management to achieve adequate parasite control for both roundworm and bot infestations.

Contraindication:

Do not use in foals under 2 weeks of age.

Do not use in horses known to be hypersensitive to the active ingredient or any of the other ingredients.

Do not use in dogs or cats as severe adverse reactions may occur.

Administration to stallions at 3 times the recommended dosage did not show any adverse effect in particular on the reproductive performances.

Administration to foals with doses up to 5 times the recommended dosage did not show any adverse reaction.

Withdrawal Period (s):

Meat and offal: 35 days.

Not permitted for use in horses producing milk for human consumption.

Storage Condition:

Store below 30°C. Protect from heat and sunlight exposure.

Keep out of reach of children/ Jauhkan dari kanak-kanak.

Any unused product or packaging should be disposed of in accordance with local and national legislation.

Shelf Life as packaged for Sale: 3 years.

Packaging Available: 1 x 10Tablets , 2 x 10Tablets

Date of revision: 13-07-2021

Product Holder/ Manufacturer:

NAM PHARMA SDN. BHD. (768370-H)

5, Lebuhr Perusahaan Klebang 11,

Taman Perindustrian Antarabangsa IGB,

31200 Chemor, Perak, Malaysia.

Tel: +605-2931838, 1839 Fax: +605-2931833