

Aetos Ivy Leaves Extract Syrup 35mg/5mL

Aetos Ivy Leaves Extract Syrup 35mg/5mL which consists of 35 mg of Ivy Leaves extract powder (*Hedera Helix*), is a herbal cough expectorant with no added alcohol and artificial colouring agent. It has been traditionally used for relief of cough, throat irritation and reducing phlegm.

No added:

- Alcohol
- Artificial colouring agent

Description: Light brown to brown, slightly cloudy syrup with vanilla flavour and aroma.

Composition: Each 5 ml contains: Ivy Leaves Extract Powder (*Hedera Helix*) 35 mg

Indication: Traditionally used for relief of cough, throat irritation and reducing phlegm.

Contraindication: Hypersensitivity to the active substance or to plants of the Araliaceae family.

Children under 2 years of age because of the general risk of aggravation of respiratory symptoms through secretolytic drugs.

Special warnings and precautions for use: Persistent or recurrent cough in children between 2 – 4 years of age required medical diagnosis before treatment.

When dyspnea, fever or purulent sputum occurs, a doctor or a pharmacist should be consulted.

Caution is recommended in patients with gastritis or gastric ulcer.

This is a traditional medicine.

Please consult your pharmacist/ doctor if symptoms persist/ worsen.

Keep out reach of children & *Jauhkan daripada capaian kanak-kanak.*

Fertility, Pregnancy and Lactation: Safety during pregnancy and lactation has not been established. In the absence of sufficient data, the use during pregnancy and lactation is not recommended.

No fertility data are available.

Undesirable effects: Gastrointestinal reactions (nausea, vomiting, diarrhoea) have been reported. The frequency is not known.

Allergic reactions (urticarial, skin rash, dyspnea, anaphylactic reaction) have been reported. The frequency is not known.

If other adverse reactions not mentioned above occur, a doctor or a pharmacist should be consulted.

Effect on ability to drive and use machines: No studies on the effect on the ability to drive and use machines have been performed.

Drug Interactions: Unknown

Overdose: Overdose can provoke nausea, vomiting and diarrhoea and agitation.

Dosage and administration:

Adults: 5 - 7.5mL 3 times daily

6 - 9 years old and adolescents (10 - 12 years old): 5mL 3 times daily.

2 - 5 years old: 2.5mL 3 times daily.

This medicine should be taken after meal.

Use measuring cup provided to measure the recommended volume.

Duration of use: If the symptoms persist longer than one week during the use of the medicinal product, a doctor or a pharmacist should be consulted.

Storage: Store below 30°C. Protect from light and moisture.

Presentation / Packing: Bottle of 100 mL.

Product Registration Holder: Abiotide Sdn. Bhd.

121, Jalan Tunku Abdul Rahman, 30010, Ipoh, Perak, Malaysia.

Manufactured by: HOVID Bhd.

121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar), 30010 Ipoh, Perak, Malaysia.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Date of revision: May 2021

3

8

5