

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory only OR for Specialist Use only

BORTECIP

Bortezomib for Injection 3.5mg/vial

COMPOSITION

Each vial contains

Bortezomib (as monohydrate) 3.5 mg

Mannitol Ph.Eur.....35 mg

PRODUCT DESCRIPTION

Before reconstitution: A white to off white lyophilized mass or powder filled in 10 ml USP Type 1 clear tubular glass vial.

After reconstitution: clear and colourless solution.

DOSAGE FORM

Powder for Solution for Injection

PHARMACOLOGY

Pharmacodynamics

Pharmacotherapeutic Group: Antineoplastic agents, other antineoplastic agents

ATC code: L01XX32

Mechanism of Action and Pharmacodynamic Effects

Bortezomib is a proteasome inhibitor. It is specifically designed to inhibit the chymotrypsin-like activity of the 26S proteasome in mammalian cells. The 26S proteasome is a large protein complex that degrades ubiquitinated proteins. The ubiquitin-proteasome pathway plays an essential role in regulating the turnover of specific proteins, thereby maintaining homeostasis within cells. Inhibition of the 26S proteasome prevents this targeted proteolysis and affects multiple signaling cascades within the cell, ultimately resulting in cancer cell death.

Bortezomib is highly selective for the proteasome. At 10 μM concentrations, bortezomib does not inhibit any of a wide variety of receptors and proteases screened and is more than 1,500-fold more selective for the proteasome than for its next preferable enzyme. The kinetics of proteasome inhibition was evaluated *in vitro*, and bortezomib was shown to dissociate from the proteasome with a $t_{1/2}$ of 20 minutes, thus demonstrating that proteasome inhibition by bortezomib is reversible.

Bortezomib mediated proteasome inhibition affects cancer cells in a number of ways, including, but not limited to, altering regulatory proteins, which control cell cycle progression and nuclear factor kappa B (NF- κ B) activation. Inhibition of the proteasome results in cell cycle arrest and apoptosis. NF- κ B is a transcription factor whose activation is required for many aspects of tumourigenesis, including cell growth and survival, angiogenesis, cell-cell interactions, and metastasis. In myeloma, bortezomib affects the ability of myeloma cells to interact with the bone marrow microenvironment.

Experiments have demonstrated that bortezomib is cytotoxic to a variety of cancer cell types and that cancer cells are more sensitive to the pro-apoptotic effects of proteasome inhibition than normal cells. Bortezomib causes reduction of tumour growth including multiple myeloma.

Data from *in vitro*, *ex-vivo*, and animal models with bortezomib suggest that it increases osteoblast differentiation and activity and inhibits osteoclast function. These effects have been observed in patients with multiple myeloma affected by an advanced osteolytic disease and treated with bortezomib

Pharmacokinetics

Absorption

Following intravenous bolus administration of a 1.0 mg/m² and 1.3 mg/m² dose to 11 patients with multiple myeloma and creatinine clearance values greater than 50 ml/min, the mean first-dose maximum plasma concentrations of bortezomib were 57 and 112 ng/ml, respectively. In subsequent doses, mean maximum observed plasma concentrations ranged from 67 to 106 ng/ml for the 1.0 mg/m² dose and 89 to 120 ng/ml for the 1.3 mg/m² dose.

Following an intravenous bolus or subcutaneous injection of a 1.3 mg/m² dose to patients with multiple myeloma, the total systemic exposure after repeat dose administration (AUC_{last}) was equivalent for subcutaneous and intravenous administrations. The C_{max} after subcutaneous administration (20.4 ng/ml) was lower than intravenous (223 ng/ml). The AUC_{last} geometric mean ratio was 0.99 and 90% confidence intervals were 80.18%-122.80%.

Distribution

The mean distribution volume (V_d) of bortezomib ranged from 1,659 l to 3,294 l following single-or repeated-dose intravenous administration of 1.0 mg/m² or 1.3 mg/m² to patients with multiple myeloma. This suggests that bortezomib distributes widely to peripheral tissues. Over a bortezomib concentration range of 0.01 to 1.0 µg/ml, the *in vitro* protein binding averaged 82.9% in human plasma. The fraction of bortezomib bound to plasma proteins was not concentration-dependent.

Biotransformation

Bortezomib is primarily oxidatively metabolized via cytochrome P450 enzymes, 3A4, 2C19, and 1A2. The major metabolic pathway is deboronation to form two deboronated metabolites that subsequently undergo hydroxylation to several metabolites. Deboronated-bortezomib metabolites are inactive as 26S proteasome inhibitors.

Elimination

The mean elimination half-life ($t_{1/2}$) of bortezomib upon multiple dosing ranged from 40-193 hours. Bortezomib is eliminated more rapidly following the first dose compared to subsequent doses. Mean total body clearances were 102 and 112 l/h following the first dose for doses of 1.0 mg/m² and 1.3 mg/m², respectively, and ranged from 15 to 32 l/h and 18 to 32 l/h following subsequent doses for doses of 1.0 mg/m² and 1.3 mg/m², respectively.

Special populations

Hepatic impairment

Mild hepatic impairment did not alter dose-normalized bortezomib AUC. However, the dose-normalized mean AUC values were increased by approximately 60% in patients with moderate or severe hepatic impairment. A lower starting dose is recommended in patients with moderate or severe hepatic impairment, and those patients should be closely monitored.

Renal impairment

Patients were administered intravenous doses of 0.7 to 1.3 mg/m² of Bortezomib twice weekly. Exposure of BORTEZOMIB (dose-normalized AUC and C_{max}) was comparable among all the groups of patients regardless of degrees of renal impairment [Normal (CrCL ≥ 60 ml/min/1.73 m², n=12), Mild (CrCL=40-59 ml/min/1.73 m², n=10), Moderate (CrCL=20-39 ml/min/1.73 m², n=9), and Severe (CrCL < 20 ml/min/1.73 m², n=3)].

Age

The pharmacokinetics of bortezomib were characterized following twice weekly intravenous bolus administration of 1.3 mg/m² doses to pediatric patients (2-16 years old) with acute lymphoblastic leukemia (ALL) or acute myeloid leukemia (AML). Based on a population pharmacokinetic analysis, clearance of bortezomib increased with increasing body surface area (BSA). Geometric mean (%CV) clearance was 7.79 (25%) L/hr/m², volume of distribution at steady-state was 834 (39%) L/m², and the elimination half-life were 100 (44%) hours. After correcting for the BSA effect, other demographics such as age, body weight, and sex did not have significant effects on bortezomib clearance. BSA-normalized clearance of bortezomib in pediatric patients was similar to that observed in adults.

INDICATIONS

BORTECIP is indicated for the treatment of patients with multiple myeloma.

BORTECIP is indicated for the treatment of patients with mantle cell lymphoma.

DOSAGE AND METHOD OF ADMINISTRATION

Important Dosing Guidelines

BORTECIP is for intravenous or subcutaneous use only. Do not administer BORTECIP by any other route.

Because each route of administration has a different reconstituted concentration, use caution when calculating the volume to be administered.

The recommended starting dose of BORTECIP is 1.3 mg/m^2 . BORTECIP may be administered intravenously at a concentration of 1 mg/ml, or subcutaneously at a concentration of 2.5 mg/ml.

BORTECIP retreatment may be considered for patients with multiple myeloma who had previously responded to treatment with BORTECIP and who have relapsed at least 6 months after completing prior BORTECIP treatment. Treatment may be started at the last tolerated dose.

When administered intravenously, administer BORTECIP as a 3 to 5 second bolus intravenous injection.

Dosage in Previously Untreated Multiple Myeloma

BORTECIP is administered in combination with oral melphalan and oral prednisone for 9, six- week treatment cycles as shown in **Table 1**. In Cycles 1 to 4, BORTECIP is administered twice weekly (Days 1, 4, 8, 11, 22, 25, 29 and 32). In Cycles 5 to 9, BORTECIP is administered once weekly (Days 1, 8, 22 and 29). At least 72 hours should elapse between consecutive doses of BORTECIP.

Table 1: Dosage Regimen for Patients with Previously Untreated Multiple Myeloma

Twice Weekly BORTECIP (Cycles 1 to 4)												
Week	1				2		3	4		5		6
BORTECIP (1.3 mg / m ²)	Day 1	-	-	Day 4	Day 8	Day 11	Rest period	Day 22	Day 25	Day 29	Day 32	Rest Period
Melphalan (9 mg / m ²) Prednisone (60 mg / m ²)	Day 1	Day 2	Day 3	Day 4	-	-	Rest period	-	-	-	-	Rest period
Once Weekly BORTECIP (Cycles 5 to 9 when used in combination with Melphalan and Prednisone)												
Week	1				2		3	4		5		6
BORTECIP (1.3 mg / m ²)	Day 1	-	-	-	Day 8	-	Rest period	Day 22	-	Day 29	-	Rest Period
Melphalan (9 mg / m ²) Prednisone (60 mg / m ²)	Day 1	Day 2	Day 3	Day 4	-	-	Rest period	-	-	-	-	Rest period

Dose Modification Guidelines for BORTECIP when given in Combination with Melphalan and Prednisone

Prior to initiating any cycle of therapy with BORTECIP in combination with melphalan and prednisone:

- Platelet count should be at least $70 \times 10^9/L$ and the absolute neutrophil count (ANC) should be at least $1.0 \times 10^9/L$
- Nonhematological toxicities should have resolved to Grade 1 or baseline

Table 2: Dose Modifications during Cycles of Combination BORTECIP, Melphalan and Prednisone Therapy

Toxicity	Dose Modification or Delay
Hematological toxicity during a cycle: If prolonged Grade 4 neutropenia or thrombocytopenia, or thrombocytopenia with bleeding is observed in the previous cycle	Consider reduction of the melphalan dose by 25% in the next cycle
If platelet count is not above $30 \times 10^9/L$ or ANC is not above $0.75 \times 10^9/L$ on a BORTECIP dosing day (other than day 1)	Withhold BORTECIP dose
If several BORTECIP doses in consecutive cycles are withheld due to toxicity	Reduce BORTECIP dose by 1 dose level (from 1.3 mg/m^2 to 1 mg/m^2 , or from 1 mg/m^2 to 0.7 mg/m^2)
Grade 3 or higher non-hematological toxicities	Withhold BORTECIP therapy until symptoms of toxicity have resolved to Grade 1 or baseline. Then, BORTECIP may be reinitiated with one dose level reduction (from 1.3 mg/m^2 to 1 mg/m^2 , or from 1 mg/m^2 to 0.7 mg/m^2). For BORTECIP-related neuropathic pain and/or peripheral neuropathy, hold or modify BORTECIP as outlined in Table 6 .

For information concerning melphalan and prednisone, see manufacturer's prescribing information.

Dose modifications guidelines for peripheral neuropathy are provided.

Posology for previously untreated multiple myeloma patients eligible for haematopoietic stem cell transplantation (induction therapy)

Combination therapy with dexamethasone

BORTECIP 3.5 mg powder for solution for injection is administered via intravenous or subcutaneous injection at the recommended dose of 1.3 mg/m² body surface area twice weekly for two weeks on days 1, 4, 8, and 11 in a 21-day treatment cycle. This 3-week period is considered a treatment cycle. At least 72 hours should elapse between consecutive doses of BORTECIP. Dexamethasone is administered orally at 40 mg on days 1, 2, 3, 4, 8, 9, 10 and 11 of the BORTECIP treatment cycle.

Four treatment cycles of this combination therapy are administered.

Combination therapy with dexamethasone and thalidomide

BORTECIP 3.5 mg powder for solution for injection is administered via intravenous or subcutaneous injection at the recommended dose of 1.3 mg/m² body surface area twice weekly for two weeks on days 1, 4, 8, and 11 in a 28-day treatment cycle. This 4-week period is considered a treatment cycle. At least 72 hours should elapse between consecutive doses of BORTECIP. Dexamethasone is administered orally at 40 mg on days 1, 2, 3, 4, 8, 9, 10 and 11 of the BORTECIP treatment cycle.

Thalidomide is administered orally at 50 mg daily on days 1-14 and if tolerated the dose is increased to 100 mg on days 15-28, and thereafter may be further increased to 200 mg daily from cycle 2 (see Table 3).

Four treatment cycles of this combination are administered. It is recommended that patients with at least partial response receive 2 additional cycles.

Table 3: Posology for BORTECIP combination therapy for patients with previously untreated multiple myeloma eligible for haematopoietic stem cell transplantation

Vc+Dx	Cycles 1 to 4			
	Week	1	2	3
	Vc (1.3 mg/m ²)	Day 1,4	Day 8,11	Rest Period
	Dx 40 mg	Day 1,2,3,4	Day 8,9,10,11	-

Vc+Dx+T	Cycle 1				
	Week	1	2	3	4
	Vc (1.3 mg/m ²)	Day 1,4	Day 8,11	Rest Period	Rest Period
	T 50 mg	Daily	Daily	-	-
	T 100 mg ^a	-	-	Daily	Daily
	Dx 40 mg	Day 1,2,3,4	Day 8,9,10,11	-	-
	Cycles 2 to 4^b				
	Vc (1.3 mg/m ²)	Day 1,4	Day 8,11	Rest Period	Rest Period
	T 200 mg	Daily	Daily	Daily	Daily
	Dx 40 mg	Day 1,2,3,4	Day 8,9,10,11	-	-

Vc = BORTECIP; Dx = Dexamethasone; T=Thalidomide

^a Thalidomide dose is increased to 100 mg from week 3 of Cycle 1 only if 50 mg is tolerated and to 200 mg from cycle 2 onwards if 100 mg is tolerated

^b Up to 6 cycles may be given to patients who achieve at least a partial response after 4 cycles

Dosage adjustments for transplant eligible patients

For BORTECIP dosage adjustments, as described under “***Dosage and Dose Modifications for Relapsed Multiple Myeloma and Relapsed Mantle Cell Lymphoma***” and “***Dose Modifications for Peripheral Neuropathy***” should be followed.

In addition, when BORTECIP is given in combination with other chemotherapeutic medicinal products, appropriate dose reductions for these products should be considered in the event of toxicities according to the recommendations in the local package inserts.

Dosage in Previously Untreated Mantle Cell Lymphoma

BORTECIP (1.3 mg/m²) is administered intravenously in combination with intravenous rituximab, cyclophosphamide, doxorubicin and oral prednisone (VcR-CAP) for 6 three-week treatment cycles as shown in **Table 4**. BORTECIP is administered first followed by rituximab. BORTECIP is administered twice weekly for two weeks (Days 1, 4, 8, and 11) followed by a ten-day rest period on Days 12 to 21. For patients with a response first documented at cycle 6, two additional VcR-CAP cycles are recommended. At least 72 hours should elapse between consecutive doses of BORTECIP.

Table 4: Dosage Regimen for Patients with Previously Untreated Mantle Cell Lymphoma

Twice Weekly BORTECIP (6, Three Week Cycles)*								
Week	1					2		3
BORTECIP (1.3 mg/m ²)	Day 1	--	--	Day 4	--	Day 8	Day 11	Rest period
Rituximab (375 mg/m ²) Cyclophosphamide (750 mg/m ²) Doxorubicin (50 mg/m ²)	Day 1	--	--			--	--	Rest period
Prednisone (100 mg/m ²)	Day 1	Day 2	Day 3	Day 4	Day 5	-	-	Rest Period

*Dosing may continue for 2 more cycles (for a total of 8 cycles) if response is first seen at cycle 6.

Dose Modification Guidelines for BORTECIP When Given in Combination with Rituximab, Cyclophosphamide, Doxorubicin and Prednisone

Prior to the first day of each cycle (other than Cycle 1):

- Platelet count should be at least $100 \times 10^9/\text{L}$ and absolute neutrophil count (ANC) should be at least $1.5 \times 10^9/\text{L}$
- Hemoglobin should be at least 8 g/dL (at least 4.96 mmol/L)
- Non-hematologic toxicity should have recovered to Grade 1 or baseline

Interrupt BORTECIP treatment at the onset of any Grade 3 hematologic or non-hematological toxicities, excluding neuropathy [see **Table 6** and **Special warnings and special precautions for use**]. For dose adjustments, see **Table 5** below.

Table 5: Dose Modifications on Days 4, 8, and 11 during Cycles of Combination BORTECIP, Rituximab, Cyclophosphamide, Doxorubicin and Prednisone Therapy

Toxicity	Dose modification or delay
Hematological toxicity	
Grade 3 or higher neutropenia, or a platelet count not at or above $25 \times 10^9/\text{L}$	Withhold BORTECIP therapy for up to 2 weeks until the patient has an ANC at or above $0.75 \times 10^9/\text{L}$ and a platelet count at or above $25 \times 10^9/\text{L}$. • If, after BORTECIP has been withheld, the toxicity does not resolve, discontinue BORTECIP. • If toxicity resolves such that the patient has an ANC at or above $0.75 \times 10^9/\text{L}$ and a platelet count at or above $25 \times 10^9/\text{L}$, BORTECIP dose should be reduced by 1 dose level (from $1.3 \text{ mg}/\text{m}^2$ to $1 \text{ mg}/\text{m}^2$, or from $1 \text{ mg}/\text{m}^2$ to $0.7 \text{ mg}/\text{m}^2$).

Grade 3 or higher non-hematological toxicities	Withhold BORTECIP therapy until symptoms of the toxicity have resolved to Grade 2 or better. Then, BORTECIP may be reinitiated with one dose level reduction (from 1.3 mg/m ² to 1 mg/m ² , or from 1 mg/m ² to 0.7 mg/m ²). For BORTECIP-related neuropathic pain and/or peripheral neuropathy, hold or modify BORTECIP as outlined in Table 6.
--	---

For information concerning rituximab, cyclophosphamide, doxorubicin and prednisone, see manufacturer's prescribing information.

Dosage and Dose Modifications for Relapsed Multiple Myeloma and Relapsed Mantle Cell Lymphoma

BORTECIP (1.3 mg/m²/dose) is administered twice weekly for two weeks (Days 1, 4, 8, and 11) followed by a tenday rest period (Days 12 to 21). For extended therapy of more than eight cycles, BORTECIP may be administered on the standard schedule or, for relapsed multiple myeloma, on a maintenance schedule of once weekly for four weeks (Days 1, 8, 15, and 22) followed by a 13 day rest period (Days 23 to 35). At least 72 hours should elapse between consecutive doses of BORTECIP.

Patients with multiple myeloma who have previously responded to treatment with BORTECIP (either alone or in combination) and who have relapsed at least six months after their prior BORTECIP therapy may be started on BORTECIP at the last tolerated dose. Retreated patients are administered BORTECIP twice weekly (Days 1, 4, 8, and 11) every three weeks for a maximum of eight cycles. At least 72 hours should elapse between consecutive doses of BORTECIP. BORTECIP may be administered either as a single agent or in combination with dexamethasone.

BORTECIP therapy should be withheld at the onset of any Grade 3 non-hematological or Grade 4 hematological toxicities excluding neuropathy as discussed below (**see Special warnings and special precautions for use**). Once the symptoms of the toxicity have resolved, BORTECIP therapy may be reinitiated at a 25% reduced dose (1.3 mg/m²/dose reduced to 1 mg/m²/dose; 1 mg/m²/dose reduced to 0.7 mg/m²/dose).

For dose modifications guidelines for peripheral neuropathy, see Dose modifications for peripheral neuropathy below.

Dose Modifications for Peripheral Neuropathy

Starting BORTECIP subcutaneously may be considered for patients with pre-existing or at high risk of peripheral neuropathy. Patients with pre-existing severe neuropathy should be treated with BORTECIP only after careful risk-benefit assessment.

Patients experiencing new or worsening peripheral neuropathy during BORTECIP therapy may require a decrease in the dose and/or a less dose-intense schedule.

For dose or schedule modification guidelines for patients who experience BORTECIP-related neuropathic pain and/or peripheral neuropathy see **Table 6**.

Table 6: Recommended Dose Modification for BORTECIP related Neuropathic Pain and/or Peripheral Sensory or Motor Neuropathy

Severity of Peripheral Neuropathy Signs and Symptoms*	Modification of Dose and Regimen
Grade 1 (asymptomatic; loss of deep tendon reflexes or paresthesias) without pain or loss of function	No action
Grade 1 with pain or Grade 2 (moderate symptoms; limiting instrumental Activities of Daily Living (ADL)**)	Reduce BORTECIP to 1 mg/m ²
Grade 2 with pain or Grade 3 (severe symptoms; limiting self care ADL***)	Withhold BORTECIP therapy until toxicity resolves. When toxicity resolves reinitiate with a reduced dose of BORTECIP at 0.7 mg/m ² once per week.
Grade 4 (life- threatening consequences; Urgent intervention indicated)	Discontinue BORTECIP

*Grading based on NCI Common Toxicity Criteria CTCAE v4.0.

**Instrumental ADL: refers to preparing meals, shopping for groceries or clothes, using telephone, managing money etc.

***Self care ADL: refers to bathing, dressing and undressing, feeding self, using the toilet, taking medications, and not bedridden

Special Populations

Patients with Renal Impairment

The pharmacokinetics of BORTECIP are not influenced by the degree of renal impairment. Therefore, dosing adjustments of BORTECIP are not necessary for patients with renal insufficiency. Since dialysis may reduce BORTECIP concentrations, the drug should be administered after the dialysis procedure (see ***Pharmacokinetic Properties***).

Patients with Hepatic Impairment

Patients with mild hepatic impairment do not require a starting dose adjustment and should be treated per the recommended BORTECIP dose. For patients with moderate or severe hepatic impairment, see **Table 7** below, (also, see ***Pharmacokinetic Properties***):

Table 7: Recommended Starting Dose Modification for BORTECIP in Patients with Hepatic Impairment

	Bilirubin Level	SGOT (AST) Levels	Modification of Starting Dose in Multiple Myeloma and Relapsed Mantle Cell Lymphoma (1.3 mg/m² twice weekly)
Mild	≤ 1.0x ULN	> ULN	None
	> 1.0x–1.5x ULN	Any	None
Moderate	> 1.5x–3x ULN	Any	Reduce BORTECIP to 0.7 mg/m ² in the first cycle. Consider dose escalation to 1.0 mg/m ² or further dose reduction to 0.5 mg/m ² in subsequent cycles based on patient tolerability.
Severe	> 3x ULN	Any	

Abbreviations: SGOT = serum glutamic oxaloacetic transaminase; AST = aspartate aminotransferase; ULN = upper limit of the normal range

Administration

BORTECIP is administered intravenously or subcutaneously. When administered intravenously, BORTECIP is administered as a 3-5 second bolus intravenous injection through a peripheral or central intravenous catheter followed by a flush with 0.9% sodium chloride solution for injection. For subcutaneous administration, the reconstituted solution is injected into the thighs (right or left) or abdomen (right or left). Injection sites should be rotated for successive injections.

If local injection site reactions occur following BORTECIP injection subcutaneously, a less concentrated BORTECIP solution (1 mg/ml instead of 2.5 mg/ml) may be administered subcutaneously or changed to IV injection.

CONTRAINDICATIONS

- Hypersensitivity to the active substance, to boron or to any of the excipients.
- Acute diffuse infiltrative pulmonary and pericardial disease.
- When Bortezomib is given in combination with other medicinal products, refer to their Summaries of Product Characteristics for additional contraindications

WARNINGS AND PRECAUTIONS

When BORTECIP is given in combination with other medicinal products, the Summary of Product Characteristics of these other medicinal products must be consulted prior to initiation of treatment with BORTECIP. When thalidomide is used, particular attention to pregnancy testing and prevention requirements is needed.

Intrathecal administration

There have been fatal cases of inadvertent intrathecal administration of BORTECIP. BORTECIP 3.5 mg powder for solution for injection is for intravenous or subcutaneous use. BORTECIP should not be administered intrathecally.

Gastrointestinal toxicity

Gastrointestinal toxicity, including nausea, diarrhea, vomiting and constipation are very common with BORTECIP treatment. Cases of ileus have been uncommonly reported. Therefore, patients who experience constipation should be closely monitored.

Haematological toxicity

BORTECIP treatment is very commonly associated with hematological toxicities (thrombocytopenia, neutropenia and anaemia).

Gastrointestinal and intracerebral haemorrhage, have been reported in association with BORTECIP treatment. Therefore, platelet counts should be monitored prior to each dose of BORTECIP. BORTECIP therapy should be withheld when the platelet count is $< 25,000/\mu\text{l}$ or, in the case of combination with melphalan and prednisone, when the platelet count is $\leq 30,000/\mu\text{l}$. Potential benefit of the treatment should be carefully weighed against the risks, particularly in case of moderate to severe thrombocytopenia and risk factors for bleeding.

Complete blood counts (CBC) with differential and including platelet counts should be frequently monitored throughout treatment with BORTECIP. Platelet transfusion should be considered when clinically appropriate.

In patients with MCL, transient neutropenia that was reversible between cycles was observed, with no evidence of cumulative neutropenia. Since patients with neutropenia are at increased risk of infections, they should be monitored for signs and symptoms of infection and treated promptly. Granulocyte colony stimulating factors may be administered for haematologic toxicity according to local standard practice. Prophylactic use of granulocyte colony stimulating factors should be considered in case of repeated delays in cycle administration.

Herpes zoster virus reactivation

Antiviral prophylaxis is recommended in patients being treated with BORTECIP.

Hepatitis B Virus (HBV) reactivation and infection

When rituximab is used in combination with BORTECIP, HBV screening must always be performed in patients at risk of infection with HBV before initiation of treatment. Carriers of hepatitis B and patients with a history of hepatitis B must be closely monitored for signs of active HBV infection during and following rituximab combination treatment with BORTECIP. Antiviral prophylaxis should be considered.

Progressive multifocal leukoencephalopathy (PML)

Very rare cases with unknown causality of John Cunningham (JC) virus infection, resulting in PML and death, have been reported in patients treated with Bortezomib. Patients diagnosed with PML had prior or concurrent immunosuppressive therapy. Most cases of PML were diagnosed within 12 months of their first dose of BORTECIP. Patients should be monitored at regular intervals for any new or worsening neurological symptoms or signs that may be suggestive of PML as part of the differential diagnosis of CNS problems. If a diagnosis of PML is suspected, patients should be referred to a specialist in PML and appropriate diagnostic measures for PML should be initiated. Discontinue BORTECIP if PML is diagnosed.

Peripheral neuropathy

Treatment with BORTECIP is very commonly associated with peripheral neuropathy, which is predominantly sensory. However, cases of severe motor neuropathy with or without sensory peripheral neuropathy have been reported. The incidence of peripheral neuropathy increases early in the treatment and has been observed to peak during cycle 5.

It is recommended that patients be carefully monitored for symptoms of neuropathy such as a burning sensation, hyperesthesia, hypoesthesia, paraesthesia, discomfort, neuropathic pain or weakness.

Patients experiencing new or worsening peripheral neuropathy should undergo neurological evaluation and may require a change in the dose, schedule or route of administration to subcutaneous. Neuropathy has been managed with supportive care and other therapies.

Early and regular monitoring for symptoms of treatment-emergent neuropathy with neurological evaluation should be considered in patients receiving Bortezomib in combination with medicinal products known to be associated with neuropathy (e.g. thalidomide) and appropriate dose reduction or treatment discontinuation should be considered.

In addition to peripheral neuropathy, there may be a contribution of autonomic neuropathy to some adverse reactions such as postural hypotension and severe constipation with ileus. Information on autonomic neuropathy and its contribution to these undesirable effects is limited.

Seizures

Seizures have been uncommonly reported in patients without previous history of seizures or epilepsy. Special care is required when treating patients with any risk factors for seizures.

Hypotension

BORTECIP treatment is commonly associated with orthostatic/postural hypotension. Most adverse reactions are mild to moderate in nature and are observed throughout treatment. Patients who developed orthostatic hypotension on Bortezomib (injected intravenously) did not have evidence of orthostatic hypotension prior to treatment with Bortezomib. Most patients required treatment for their orthostatic hypotension. A minority of patients with orthostatic hypotension experienced syncopal events. Orthostatic/postural hypotension was not acutely related to bolus infusion of Bortezomib. The mechanism of this event is unknown although a component may be due to autonomic neuropathy. Autonomic neuropathy may be related to bortezomib or bortezomib may aggravate an underlying condition such as diabetic or amyloidotic neuropathy. Caution is advised when treating patients with a history of syncope receiving medicinal products known to be associated with hypotension; or who are dehydrated due to recurrent diarrhoea or vomiting. Management of orthostatic/postural hypotension may include adjustment of antihypertensive medicinal products, rehydration or administration of mineralocorticosteroids and/or

sympathomimetics. Patients should be instructed to seek medical advice if they experience symptoms of dizziness, light-headedness or fainting spells.

Posterior Reversible Encephalopathy Syndrome (PRES)

PRES is a rare, often reversible, rapidly evolving neurological condition, which can present with seizure, hypertension, headache, lethargy, confusion, blindness, and other visual and neurological disturbances. Brain imaging, preferably Magnetic Resonance Imaging (MRI), is used to confirm the diagnosis. In patients developing PRES, BORTECIP should be discontinued.

Heart failure

Acute development or exacerbation of congestive heart failure, and/or new onset of decreased left ventricular ejection fraction has been reported during bortezomib treatment. Fluid retention may be a predisposing factor for signs and symptoms of heart failure. Patients with risk factors for or existing heart disease should be closely monitored.

Electrocardiogram investigations

There have been isolated cases of QT-interval prolongation in clinical studies, causality has not been established.

Pulmonary disorders

There have been rare reports of acute diffuse infiltrative pulmonary disease of unknown aetiology such as pneumonitis, interstitial pneumonia, lung infiltration, and acute respiratory distress syndrome (ARDS) in patients receiving BORTECIP. Some of these events have been fatal. A pre-treatment chest radiograph is recommended to serve as a baseline for potential post-treatment pulmonary changes.

In the event of new or worsening pulmonary symptoms (e.g., cough, dyspnoea), a prompt diagnostic evaluation should be performed and patients treated appropriately. The benefit/risk ratio should be considered prior to continuing BORTECIP therapy.

High-dose cytarabine (2 g/m² per day) by continuous infusion over 24 hours with daunorubicin and BORTECIP for relapsed acute myelogenous leukaemia is not recommended.

Renal impairment

Renal complications are frequent in patients with multiple myeloma. Patients with renal impairment should be monitored closely.

Hepatic impairment

Bortezomib is metabolized by liver enzymes. Bortezomib exposure is increased in patients with moderate or severe hepatic impairment; these patients should be treated with Bortezomib at reduced doses and closely monitored for toxicities.

Hepatic reactions

Rare cases of hepatic failure have been reported in patients receiving BORTECIP and concomitant medicinal products and with serious underlying medical conditions. Other reported hepatic reactions include increases in liver enzymes, hyperbilirubinaemia, and hepatitis. Such changes may be reversible upon discontinuation of bortezomib.

Tumour lysis syndrome

Because bortezomib is a cytotoxic agent and can rapidly kill malignant plasma cells and MCL cells, the complications of tumour lysis syndrome may occur. The patients at risk of tumour lysis syndrome are those with high tumour burden prior to treatment. These patients should be monitored closely and appropriate precautions taken.

Concomitant medicinal products

Patients should be closely monitored when given bortezomib in combination with potent CYP3A4-inhibitors. Caution should be exercised when bortezomib is combined with CYP3A4- or CYP2C19 substrates.

Normal liver function should be confirmed and caution should be exercised in patients receiving oral hypoglycemics.

Potentially immunocomplex-mediated reactions

Potentially immunocomplex-mediated reactions, such as serum-sickness-type reaction, polyarthrititis with rash and proliferative glomerulonephritis have been reported uncommonly. Bortezomib should be discontinued if serious reactions occur.

DRUG INTERACTIONS

Bortezomib is a weak inhibitor of the cytochrome P450 (CYP) isozymes 1A2, 2C9, 2C19, 2D6 and 3A4. Based on the limited contribution (7%) of CYP2D6 to the metabolism of bortezomib, the CYP2D6 poor metaboliser phenotype is not expected to affect the overall disposition of bortezomib.

Patients should be closely monitored when given bortezomib in combination with potent CYP3A4 inhibitors (e.g. ketoconazole, ritonavir).

There was no significant effect on bortezomib (injected intravenously) in combination with omeprazole, a potent CYP2C19 inhibitor.

Concomitant use of bortezomib with strong CYP3A4 inducers (e.g., rifampicin, carbamazepine, phenytoin, phenobarbital and St. John's Wort) is not recommended, as efficacy may be reduced.

There was no significant effect on bortezomib in combination with dexamethasone, a weaker CYP3A4 inducer.

Hypoglycemia and hyperglycemia were uncommonly and commonly reported in diabetic patients receiving oral hypoglycemics. Patients on oral antidiabetic agents receiving BORTECIP treatment may require close monitoring of their blood glucose levels and adjustments of the dose of their antidiabetics.

FERTILITY, PREGNANCY AND LACTATION

Contraception in males and females

Male and female patients of childbearing potential must use effective contraceptive measures during and for 3 months following treatment.

Pregnancy

If Bortezomib is used during pregnancy, or if the patient becomes pregnant while receiving this medicinal product, the patient should be informed of potential for hazard to the foetus.

Thalidomide is a known human teratogenic active substance that causes severe life-threatening birth defects. Thalidomide is contraindicated during pregnancy and in women of childbearing potential unless all the conditions of the thalidomide pregnancy prevention programme are met. Patients receiving Bortezomib in combination with thalidomide should adhere to the pregnancy prevention programme of thalidomide. Refer to the Summary of Product Characteristics of thalidomide for additional information.

Breast-feeding

It is not known whether bortezomib is excreted in human milk. Because of the potential for serious adverse reactions in breast-fed infants, breast-feeding should be discontinued during treatment with Bortezomib.

Fertility

Fertility studies were not conducted with Bortezomib.

UNDESIRABLE EFFECTS

Summary of the safety profile

Serious adverse reactions uncommonly reported during treatment with BORTECIP include cardiac failure, tumour lysis syndrome, pulmonary hypertension, posterior reversible encephalopathy syndrome, acute diffuse infiltrative pulmonary disorders and rarely autonomic neuropathy.

The most commonly reported adverse reactions during treatment with BORTECIP are nausea, diarrhoea, constipation, vomiting, fatigue, pyrexia, thrombocytopenia, anaemia, neutropenia, peripheral neuropathy (including sensory), headache, paraesthesia, decreased appetite, dyspnoea, rash, herpes zoster and myalgia.

Tabulated summary of adverse reactions

Multiple Myeloma

Adverse reactions are listed below by system organ class and frequency grouping. Frequencies are defined as: Very common; common; uncommon; rare; very rare; not known.

Adverse reactions in patients with Multiple Myeloma treated with BORTECIP

System Class	Organ	Incidence	Adverse reaction
Infections and infestations		Common	Herpes zoster (inc disseminated & ophthalmic), Pneumonia*, Herpes simplex*, Fungal infection*
		Uncommon	Infection*, Bacterial infections*, Viral infections*, Sepsis (inc septic shock)*, Bronchopneumonia, Herpes virus infection*, Meningoencephalitis herpetic#, Bacteraemia (inc staphylococcal), Hordeolum, Influenza, Cellulitis, Device related infection, Skin infection*, Ear infection*, Staphylococcal infection, Tooth infection*
		Rare	Meningitis (inc bacterial), Epstein-Barr virus infection, Genital herpes, Tonsillitis, Mastoiditis, Post viral fatigue syndrome
Neoplasms benign, malignant and unspecified (incl cysts and polyps)		Rare	Neoplasm malignant, Leukaemia plasmacytic, Renal cell carcinoma, Mass, Mycosis fungoides, Neoplasm benign*
Blood and lymphatic disorders	system	Very Common	Thrombocytopenia*, Neutropenia*, Anaemia*
		Common	Leukopenia*, Lymphopenia*

	Uncommon	Pancytopenia*, Febrile neutropenia, Coagulopathy*, Leukocytosis*, Lymphadenopathy, Haemolytic anaemia#
	Rare	Disseminated intravascular coagulation, Thrombocytosis*, Hyperviscosity syndrome, Platelet disorder NOS, Thrombotic microangiopathy (inc Thrombocytopenic purpura)#, Blood disorder NOS, Haemorrhagic diathesis, Lymphocytic infiltration
Immune system disorders	Uncommon	Angioedema#, Hypersensitivity*
	Rare	Anaphylactic shock, Amyloidosis, Type III immune complex mediated reaction
Endocrine disorders	Uncommon	Cushing's syndrome*, Hyperthyroidism*, Inappropriate antidiuretic hormone secretion
	Rare	Hypothyroidism
Metabolism and nutrition disorders	Very Common	Decreased appetite
	Common	Dehydration, Hypokalaemia*, Hyponatraemia*, Blood glucose abnormal*, Hypocalcaemia*, Enzyme abnormality*
	Uncommon	Tumour lysis syndrome, Failure to thrive*, Hypomagnesaemia*, Hypophosphataemia*, Hyperkalaemia*, Hypercalcaemia*, Hyponatraemia*, Uric acid abnormal*, Diabetes mellitus*, Fluid retention
	Rare	Hypermagnesaemia*, Acidosis, Electrolyte imbalance*, Fluid overload, Hypochloraemia*, Hypovolaemia, Hyperchloraemia*, Hyperphosphataemia*, Metabolic disorder, Vitamin B complex deficiency, Vitamin B12 deficiency, Gout, Increased appetite, Alcohol intolerance
Psychiatric disorders	Common	Mood disorders and disturbances*, Anxiety disorder*, Sleep disorders and disturbances*
	Uncommon	Mental disorder*, Hallucination*, Psychotic disorder*, Confusion*, Restlessness
	Rare	Suicidal ideation*, Adjustment disorder, Delirium, Libido decreased
Nervous system disorders	Very Common	Neuropathies*, Peripheral sensory neuropathy, Dysaesthesia*, Neuralgia*
	Common	Motor neuropathy*, Loss of consciousness (inc syncope), Dizziness*, Dysgeusia*, Lethargy, Headache*
	Uncommon	Tremor, Peripheral sensorimotor neuropathy, Dyskinesia*, Cerebellar

		coordination and balance disturbances*, Memory loss (exc dementia)*, Encephalopathy*, Posterior Reversible Encephalopathy Syndrome#, Neurotoxicity, Seizure disorders*, Post herpetic neuralgia, Speech disorder*, Restless legs syndrome, Migraine, Sciatica, Disturbance in attention, Reflexes abnormal*, Parosmia
	Rare	Cerebral haemorrhage*, Haemorrhage intracranial (inc subarachnoid)*, Brain oedema, Transient ischaemic attack, Coma, Autonomic nervous system imbalance, Autonomic neuropathy, Cranial palsy*, Paralysis*, Paresis*, Presyncope, Brain stem syndrome, Cerebrovascular disorder, Nerve root lesion, Psychomotor hyperactivity, Spinal cord compression, Cognitive disorder NOS, Motor dysfunction, Nervous system disorder NOS, Radiculitis, Drooling, Hypotonia, Guillain-Barre syndrome, Demyelinating polyneuropathy
Eye disorders	Common	Eye swelling*, Vision abnormal*, Conjunctivitis*
	Uncommon	Eye haemorrhage*, Eyelid infection*, Chalazion#, Blepharitis#, Eye inflammation*, Diplopia, Dry eye*, Eye irritation*, Eye pain, Lacrimation increased, Eye discharge
	Rare	Corneal lesion*, Exophthalmos, Retinitis, Scotoma, Eye disorder (inc. eyelid) NOS, Dacryoadenitis acquired, Photophobia, Photopsia, Optic neuropathy#, Different degrees of visual impairment (up to blindness)*
Ear and labyrinth disorders	Common	Vertigo*
	Uncommon	Dysacusis (inc tinnitus)*, Hearing impaired (up to and inc deafness), Ear discomfort*
	Rare	Ear haemorrhage, Vestibular neuronitis, Ear disorder NOS
Cardiac disorders	Uncommon	Cardiac tamponade#, Cardio-pulmonary arrest*, Cardiac fibrillation (inc atrial), Cardiac failure (inc left and right ventricular)*, Arrhythmia*, Tachycardia*, Palpitations, Angina pectoris, Pericarditis (inc pericardial effusion)*, Cardiomyopathy*, Ventricular dysfunction*, Bradycardia
	Rare	Atrial flutter, Myocardial infarction*, Atrioventricular block*, Cardiovascular disorder (inc cardiogenic shock), Torsade de pointes, Angina unstable, Cardiac valve disorders*, Coronary artery insufficiency, Sinus arrest
Vascular disorders	Common	Hypotension*, Orthostatic hypotension, Hypertension*
	Uncommon	Cerebrovascular accident#, Deep vein thrombosis*, Haemorrhage*, Thrombophlebitis (inc superficial), Circulatory collapse (inc hypovolaemic shock), Phlebitis, Flushing*, Haematoma (inc perirenal)*,

		Poor peripheral circulation*, Vasculitis, Hyperaemia (inc ocular)*
	Rare	Peripheral embolism, Lymphoedema, Pallor, Erythromelalgia, Vasodilatation, Vein discolouration, Venous insufficiency
Respiratory, thoracic and mediastinal disorders	Common	Dyspnoea*, Epistaxis, Upper/lower respiratory tract infection*, Cough*
	Uncommon	Pulmonary embolism, Pleural effusion, Pulmonary oedema (inc acute), Pulmonary alveolar haemorrhage#, Bronchospasm, Chronic obstructive pulmonary disease*, Hypoxaemia*, Respiratory tract congestion*, Hypoxia, Pleurisy*, Hiccups, Rhinorrhoea, Dysphonia, Wheezing
	Rare	Respiratory failure, Acute respiratory distress syndrome, Apnoea, Pneumothorax, Atelectasis, Pulmonary hypertension, Haemoptysis, Hyperventilation, Orthopnoea, Pneumonitis, Respiratory alkalosis, Tachypnoea, Pulmonary fibrosis, Bronchial disorder*, Hypocapnia*, Interstitial lung disease, Lung infiltration, Throat tightness, Dry throat, Increased upper airway secretion, Throat irritation, Upper-airway cough syndrome
Gastrointestinal disorders	Very Common	Nausea and vomiting symptoms*, Diarrhoea*, Constipation
	Common	Gastrointestinal haemorrhage (inc mucosal)*, Dyspepsia, Stomatitis*, Abdominal distension, Oropharyngeal pain*, Abdominal pain (inc gastrointestinal and splenic pain)*, Oral disorder*, Flatulence
	Uncommon	Pancreatitis (inc chronic)*, Haematemesis, Lip swelling*, Gastrointestinal obstruction (inc small intestinal obstruction, ileus)*, Abdominal discomfort, Oral ulceration*, Enteritis*, Gastritis*, Gingival bleeding, Gastrooesophageal reflux disease*, Colitis (inc clostridium difficile)*, Colitis ischaemic#, Gastrointestinal inflammation*, Dysphagia, Irritable bowel syndrome, Gastrointestinal disorder NOS, Tongue coated, Gastrointestinal motility disorder*, Salivary gland disorder*
	Rare	Pancreatitis acute, Peritonitis*, Tongue oedema*, Ascites, Oesophagitis, Cheilitis, Faecal incontinence, Anal sphincter atony, Faecaloma*, Gastrointestinal ulceration and perforation*, Gingival hypertrophy, Megacolon, Rectal discharge, Oropharyngeal blistering*, Lip pain, Periodontitis, Anal fissure, Change of bowel habit, Proctalgia, Abnormal faeces
Hepatobiliary disorders	Common	Hepatic enzyme abnormality*
	Uncommon	Hepatotoxicity (inc liver disorder), Hepatitis*, Cholestasis

	Rare	Hepatic failure, Hepatomegaly, Budd-Chiari syndrome, Cytomegalovirus hepatitis, Hepatic haemorrhage, Cholelithiasis
Skin and subcutaneous tissue disorders	Common	Rash*, Pruritus*, Erythema, Dry skin
	Uncommon	Erythema multiforme, Urticaria, Acute febrile neutrophilic dermatosis, Toxic skin eruption, Toxic epidermal necrolysis#, Stevens-Johnson syndrome#, Dermatitis*, Hair disorder*, Petechiae, Ecchymosis, Skin lesion, Purpura, Skin mass*, Psoriasis, Hyperhidrosis, Night sweats, Decubitus ulcer#, Acne*, Blister*, Pigmentation disorder*
	Rare	Skin reaction, Jessner's lymphocytic infiltration, Palmar-plantar erythrodysesthesia syndrome, Haemorrhage subcutaneous, Livedo reticularis, Skin induration, Papule, Photosensitivity reaction, Seborrhoea, Cold sweat, Skin disorder NOS, Erythrodermia, Skin ulcer, Nail disorder
Musculoskeletal and connective tissue disorders	Very Common	Musculoskeletal pain*
	Common	Muscle spasms*, Pain in extremity, Muscular weakness
	Uncommon	Muscle twitching, Joint swelling, Arthritis*, Joint stiffness, Myopathies*, Sensation of heaviness
	Rare	Rhabdomyolysis, Temporomandibular joint syndrome, Fistula, Joint effusion, Pain in jaw, Bone disorder, Musculoskeletal and connective tissue infections and inflammations*, Synovial cyst
Renal and urinary disorders	Common	Renal impairment*
	Uncommon	Renal failure acute, Renal failure chronic*, Urinary tract infection*, Urinary tract signs and symptoms*, Haematuria*, Urinary retention, Micturition disorder*, Proteinuria, Azotaemia, Oliguria*, Pollakiuria
	Rare	Bladder irritation
Reproductive system and breast disorders	Uncommon	Vaginal haemorrhage, Genital pain*, Erectile dysfunction,
	Rare	Testicular disorder*, Prostatitis, Breast disorder female, Epididymal tenderness, Epididymitis, Pelvic pain, Vulval ulceration
Congenital, familial and genetic disorders	Rare	Aplasia, Gastrointestinal malformation, Ichthyosis
General disorders and administration site conditions	Very Common	Pyrexia*, Fatigue, Asthenia
	Common	Oedema (inc peripheral), Chills, Pain*, Malaise*

	Uncommon	General physical health deterioration*, Face oedema*, Injection site reaction*, Mucosal disorder*, Chest pain, Gait disturbance, Feeling cold, Extravasation*, Catheter related complication*, Change in thirst*, Chest discomfort, Feeling of body temperature change*, Injection site pain*
	Rare	Death (inc sudden), Multi-organ failure, Injection site haemorrhage*, Hernia (inc hiatus)*, Impaired healing*, Inflammation, Injection site phlebitis*, Tenderness, Ulcer, Irritability, Non-cardiac chest pain, Catheter site pain, Sensation of foreign body
Investigations	Common	Weight decreased
	Uncommon	Hyperbilirubinaemia*, Protein analyses abnormal*, Weight increased, Blood test abnormal*, C-reactive protein increased
	Rare	Blood gases abnormal*, Electrocardiogram abnormalities (inc QT prolongation)*, International normalized ratio abnormal*, Gastric pH decreased, Platelet aggregation increased, Troponin I increased, Virus identification and serology*, Urine analysis abnormal*
Injury, poisoning and procedural complications	Uncommon	Fall, Contusion
	Rare	Transfusion reaction, Fractures*, Rigors*, Face injury, Joint injury*, Burns, Laceration, Procedural pain, Radiation injuries*
Surgical and medical procedures	Rare	Macrophage activation

NOS = not otherwise specified

*Grouping of more than one MedDRA preferred term.

Post-marketing adverse reaction regardless of indication

Mantle Cell Lymphoma (MCL)

Additional adverse drug reactions identified associated with the use of the combination therapy [Bortezomib, Rituximab, cyclophosphamide, doxorubicin and prednisone (VcR-CAP)] were hepatitis B infection and myocardial ischaemia. Higher incidence of the haematological adverse reactions (neutropenia, thrombocytopenia, leukopenia, anemia, lymphopenia), peripheral sensory neuropathy, hypertension, pyrexia, pneumonia, stomatitis, and hair disorders in MCL patient population as compared to patient with multiple myeloma.

Adverse reactions are listed below by system organ class and frequency grouping. Frequencies are defined as: Very common; common; uncommon; rare; very rare, not known.

Adverse reactions in patients with Mantle Cell Lymphoma treated with VcR-CAP

System Class	Organ	Incidence	Adverse reaction
Infections and infestations		Very Common	Pneumonia*
		Common	Sepsis (inc septic shock)*, Herpes zoster (inc disseminated & ophthalmic), Herpes virus infection*, Bacterial infections*, Upper / lower respiratory tract infections*, Fungal infection*, Herpes simplex*
		Uncommon	Hepatitis B Infection*, Bronchopneumonia
Blood and lymphatic disorders		Very Common	Thrombocytopenia*, Febrile neutropenia, Neutropenia*, Anaemia*, Leukopenia*, Lymphopenia*
		Uncommon	Pancytopenia*
Immune system disorders		Common	Hypersensitivity*
		Uncommon	Anaphylactic reaction
Metabolism and nutrition disorders		Very Common	Decreased appetite
		Common	Hypokalaemia*, Hyponatraemia*, Blood glucose abnormal*, Diabetes mellitus*, Fluid retention
		Uncommon	Tumour lysis syndrome
Psychiatric disorders		Common	Sleep disorders and disturbances*
Nervous system disorders		Very Common	Peripheral sensory neuropathy, Dysaesthesia*, Neuralgia*
		Common	Neuropathies*, Motor neuropathy*, Loss of consciousness (inc syncope), Encephalopathy*, Peripheral sensorimotor neuropathy, Dizziness*, Dysgeusia*, Autonomic neuropathy
		Uncommon	Autonomic nervous system imbalance
		Rare	Guillain-Barre syndrome, Demyelinating polyneuropathy
Eye disorders		Common	Vision abnormal*,
Ear and labyrinth disorders		Common	Dysacusis (inc tinnitus)*
		Uncommon	Vertigo*, Hearing impaired (up to and inc deafness)
Cardiac disorders		Common	Cardiac fibrillation (inc atrial), Cardiac failure (inc left and right ventricular)*, Arrhythmia*, Ventricular dysfunction*, Myocardial

		ischaemia
	Uncommon	Cardiovascular disorder (inc cardiogenic shock)
Vascular disorders	Common	Hypotension*, Orthostatic hypotension, Hypertension*
Respiratory, thoracic and mediastinal disorders	Common	Dyspnoea*, Cough*, Hiccups
	Uncommon	Acute respiratory distress syndrome, Pulmonary embolism, Pneumonitis, Pulmonary hypertension, Pulmonary oedema (inc acute)
Gastrointestinal disorders	Very Common	Nausea and vomiting symptoms*, Diarrhoea*, Stomatitis*, Constipation
	Common	Gastrointestinal haemorrhage (inc mucosal)*, Dyspepsia, Abdominal distension, Oropharyngeal pain*, Gastritis*, Oral ulceration*, Abdominal discomfort, Dysphagia, Gastrointestinal inflammation*, Abdominal pain (inc gastrointestinal and splenic pain)*, Oral disorder*
	Uncommon	Colitis (inc clostridium difficile)*
	Common	Hepatotoxicity (inc liver disorder), Hepatitis*, Cholestasis
	Uncommon	Hepatic failure
Skin and subcutaneous tissue disorders	Very Common	Hair disorder*
	Common	Rash*, Pruritus*, Dermatitis*
Musculoskeletal and connective tissue disorders	Common	Musculoskeletal pain*, Muscle spasms*, Pain in extremity,
Renal and urinary disorders	Common	Urinary tract infection*
General disorders and administration site conditions	Very Common	Pyrexia*, Fatigue, Asthenia
	Common	Oedema (inc peripheral), Chills, Injection site reaction*, Malaise*
	Common	Hyperbilirubinaemia*, Protein analyses abnormal*, Weight decreased, Weight increased

*Grouping of more than one MedDRA preferred term.

OVERDOSE

In patients, overdose more than twice the recommended dose has been associated with the acute onset of symptomatic hypotension and thrombocytopenia with fatal outcomes.

There is no known specific antidote for bortezomib overdose. In the event of an overdose, the patient's vital signs should be monitored and appropriate supportive care given to maintain blood pressure (such as fluids, pressors, and/or inotropic agents) and body temperature.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

Bortezomib may have a moderate influence on the ability to drive and use machines. Bortezomib may be associated with fatigue very commonly, dizziness commonly, syncope uncommonly and orthostatic / postural hypotension or blurred vision commonly. Therefore, patients must be cautious when driving or using machines and should be advised not to drive or operate machinery if they experience these symptoms (see *Undesirable Effects*).

INSTRUCTIONS FOR USE

Instructions for reconstitution

BORTEZOMIB must be reconstituted by a healthcare professional.

Intravenous injection

Add 3.5 ml of sodium chloride 9 mg/ml (0.9%) solution for injection, by using a syringe of the appropriate size, without removing the vial stopper. Dissolution of the lyophilised powder is completed in less than 2 minutes.

After reconstitution, each ml solution contains 1 mg bortezomib (final concentration = 1mg/mL). The reconstituted solution is clear and colourless, with a final pH of 4 to 7.

The reconstituted solution must be inspected visually for particulate matter and discolouration prior to administration. If any discolouration or particulate matter is observed, the reconstituted solution must be discarded.

Subcutaneous injection

Add 1.4 ml of sodium chloride 9 mg/ml (0.9%) solution for injection, by using a syringe of the appropriate size, without removing the vial stopper. Dissolution of the lyophilised powder is completed in less than 2 minutes.

After reconstitution, each ml solution contains 2.5 mg bortezomib (final concentration = 2.5mg/mL). The reconstituted solution is clear and colourless, with a final pH of 4 to 7.

The reconstituted solution must be inspected visually for particulate matter and discolouration prior to administration. If any discolouration or particulate matter is observed, the reconstituted solution must be discarded.

INCOMPATIBILITY

Not applicable

STORAGE AND HANDLING INSTRUCTION

Unopened vial: Do not store above 30° C. Protect from light.

Reconstituted Solution: Do not store above 25 °C. Use within 8 hours of reconstitution.

SHELF LIFE

24 months

PACKAGING INFORMATION

Carton containing 10 ml glass vial.

PRODUCT REGISTRATION HOLDER

Cipla Malaysia Sdn Bhd
Suite 1101, Amcorp Tower,
Amcorp Trade Centre,
18 Persiaran Barat,
46050 Petaling Jaya,
Selangor, Malaysia

MANUFACTURED BY

Panacea Biotec Pharma Ltd
Malpur, Baddi, Tehsil Nalagarh,
District Solan, Himachal Pradesh, 173205, India.

DATE OF REVISION

January 2022