

PACKAGE INSERTS

AMPRO EMV 300mg/g Water Soluble Powder

For the Prevention and treatment of coccidiosis in Broilers, Broiler Breeders, Replacement Pullets, Turkeys, Ducks and Pigeons.

Brand Name: **AMPRO EMV 300 mg/g Water Soluble Powder**

Dosage form: **Water Soluble Powder**

Qualitative and quantitative composition

Each gram of product contains:

Active substance:

Amprolium hydrochloride.....300 mg

Excipients q.s to.....1 gram

Product description: white to ivory-white powder with characteristic odor

Pharmacodynamics:

By mimicking its structure, amprolium competitively inhibits thiamine utilization by the parasite. Prolonged high dosages can cause thiamine deficiency in the host; excessive thiamine in the diet can reduce or reverse the anticoccidial activity of the drug.

Amprolium is thought to act primarily upon the first generation schizont in the cells of the intestinal wall, preventing differentiation of the metrozoites. It may suppress the sexual stages and sporulation of the oocysts.

Pharmacokinetics:

After oral administration absorption is low, reaching the maximum concentration 4 hours later. It is excreted mainly through faeces.

Environmental Properties: Not applicable

Indication:

AMPRO EMV 300 mg/g Water Soluble Powder is administered through the drinking water as an aid in the prevention and treatment of economically important strains of coccidiosis in poultry including *Eimeria tenella*, *Eimeria necatrix* and most strains of *E.acervulina*.

Target species:

Poultry (Broilers, Broiler Breeders, Replacement Pullets, Turkeys, Ducks and Pigeons)

Recommended dose:

Prevention:

22.50 g /25 Liters of drinking water (240 mg Amprolium per liter) for 24 hours every fourth day, during periods of likely infection. This action helps control the coccidia by interrupting the life cycle and encourages immunity to develop.

For treatment:

22.50 g /25 Liters of drinking water (240 mg Amprolium per liter) continuously for 5-7 days, reduce this to 11.35 g/ liter for a further 5-7 days, then to 6 g/ liter for another 5-7 days. This gradual reduction in the level of treatment assists the birds to develop immunity while maintaining some level of protection.

Instruction for use:

Administration:

Premix the amount of product required in 1/2 liters of water first and when completely dissolved add to drinking water, mixing thoroughly.

A fresh solution should be prepared daily.

Route of Administration: by orally.

Contraindication:

Do not administer vitamin B1 (thiamine) supplements during treatment with AMPRO EMV 300 mg/g Water Soluble Powder. Because Vitamin B1 will reduce the efficacy of the treatment.

Warning and precautions:

Special precautions for use:

Special precautions for use in target animals

The product is intended for preventive and treatment use,

Special precaution to be taken by the person administering the veterinary medicine product to animals

This active ingredient was not classified as non-hazardous substance according to Regulation (EC) No.1272/2008

This product should be handled in places with good ventilation, to avoid dust forming

Should wear impervious gloves and protective masks when handling the product to avoid directly contact with and inhale the product

Side effects:

None known

Interaction of other medicaments:

THIAMINE: Exogenously administered thiamine in high doses may reverse or reduce the efficacy of amprolium

Pregnancy and lactation:

Amprolium is the only active pharmaceutical ingredient approved for prevention and treatment in laying chicken.

Symptoms and Treatment of overdose:

Overdosage of amprolium can suppress weight gain in broilers and cause polyneuritis.

To treat amprolium overdose, thiamine should be administered parenterally or orally.

Storage condition:

Store in cold place with room temperature below 30°C

Shelf life:

Shelf-life of product packaged for sale: 2 years.

Shelf-life of product, after the first opening: should be used up within 7 days

Shelf-life after dilution according to directions: 12 hours.

Withdrawal Period:

Poultry

- Meat and offal : zero days

- Eggs: zero days

Manufacturer:**Emivest Feedmill Vietnam Company Limited**

Address: No. A-11C-CN Bau Bang Industrial Park, Lai Uyen Ward, Bau Bang District, Binh Duong Province, Vietnam

Importer and Distributor:**FE VENTURE SDN. BHD**

Lot 3 & 5, Jalan PJS 11/8, Bandar Sunway, 46150 Petaling Jaya, Selangor Darul Ehsan, Malaysia.