

For the use only of a Veterinary Medical Practitioner or a Hospital or a Laboratory or a farm

Praziquantel, Pyrantel Pamoate & Febantel Tablets

Prazivet Plus

Composition:

Each uncoated tablet contains

Praziquantel USP	50mg
Pyrantel Pamoate USP	144mg
Febantel	150mg

Product Description

Pale Yellow Coloured Uncoated Tablets

Pharmacodynamics

This product contains anthelmintics active against gastrointestinal roundworms and tapeworms. The product contains three active substances:

1. Febantel, a probenzimidazole,
2. Pyrantel embonate (pamoate) a tetrahydropyrimidine derivative,
3. Praziquantel, a partially hydrogenated pyrazinoisoquinoline derivative

ATC VetCode: QP52AF30

In this fixed combination pyrantel and febantel act against all relevant nematodes (ascarids, hookworms, and whipworms) in dogs. In particular the activity spectrum covers *Toxocara canis*, *Toxascaris leonina*, *Uncinaria stenocephala*, *Ancylostoma caninum* and *Trichuris vulpis*. This combination shows synergistic activity in the case of hookworms and febantel is effective against *T. vulpis*.

The spectrum of activity of praziquantel covers all important cestode species in dogs, in particular *Taenia* spp; *Dipylidium caninum*; *Echinococcus granulosus* and *Echinococcus multilocularis*. Praziquantel acts against all adult and immature forms of these parasites.

Praziquantel is very rapidly absorbed through the parasite's surface and distributed throughout the parasite. Both in vitro and in vivo studies have shown that praziquantel causes severe damage to the parasite integument, resulting in the contraction and paralysis of the parasites. There is an almost instantaneous tetanic contraction of the parasite musculature and a rapid vacuolisation of the syncytial tegument. This rapid contraction has been explained by changes in divalent cation fluxes, especially calcium.

Pyrantel acts as a cholinergic agonist. Its mode of action is to stimulate nicotinic cholinergic receptors of the parasite, induce spastic paralysis of the nematodes and thereby allow removal from the gastro intestinal (GI) system by peristalsis.

Within the mammalian system febantel undergoes ring closure forming fenbendazole and oxfendazole. It is these chemical entities which exert the anthelmintic effect by inhibition of tubulin polymerisation. Formation of microtubules is thereby prevented, resulting in disruption of structures vital to the normal functioning of the helminth. Glucose uptake, in particular is affected, leading to a depletion in cell ATP. The parasite dies upon exhaustion of its energy reserves, which occurs 2-3 days later.

Pharmacokinetics

No data available.

Environmental Properties (for product used directly in the environment e.g medicine for fish)

Non- hazardous

Indication

For the control of the following gastrointestinal tapeworms and roundworms in dogs and puppies.

Ascarids: Toxocara canis, Toxascaris leonina (adult and late immature forms).

Hookworms: Uncinaria stenocephala, Ancylostoma caninum (adults)

Whipworms: Trichuris vulpis (adults)

Tapeworms: Echinococcus spp., Taenia spp., Dipylidium caninum (adult and immature forms)

Target Species

Dogs

Recommended Dose

The recommended dose rates are: 15 mg/kg bodyweight febantel, 14.4 mg/kg pyrantel and 5 mg/kg praziquantel. This is equivalent to 1 tablet per 10 kg (22 lbs) bodyweight.

Puppies and Small Dogs:

3-5 kg bodyweight = ½ tablet

>5-10 kg bodyweight 1 tablet

Medium Dogs:

>10-15 kg bodyweight = 1 ½ tablets

>15-20 kg bodyweight = 2 tablets

>20-25 kg bodyweight = 2 ½ tablets

>25-30 kg bodyweight = 3 tablets

Large Dogs:

>30-35 kg bodyweight = 3 ½ tablets

>35-40 kg bodyweight = 4 tablets

For oral administration, the tablets can be given to the dog or disguised in food. No starvation is needed before, or after, treatment.

Puppies should be treated at 2 weeks of age and every 2 weeks until 12 weeks of age. Thereafter they should be treated at 3 month intervals. It is advisable to treat the bitch at the same time as the puppies. Not for use in dogs weighing less than 3 kg.

For the control of *Toxocara*, nursing bitches should be dosed 2 weeks after giving birth and every two weeks until weaning.

For routine worm control adult dogs should be treated every 3 months.

For routine treatment a single dose is recommended.

In the event of heavy roundworm infestation a repeat dose should be given after 14 days.

Route of Administration

Oral

Contraindications

Do not use simultaneously with piperazine compounds.

Warnings and Precautions

As a precautionary measure to prevent the establishment of *Echinococcus multilocularis* in the Malaysia, it is recommended that all dogs and cats entering the country be treated with praziquantel.

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*. Tapeworm infestation is certain to reoccur unless control of intermediate hosts such as fleas, mice etc is undertaken.

i) Special precautions for use in animals

Any part used tablet should be discarded

Consult a veterinary surgeon before treating pregnant animals for roundworms.

Do not exceed the stated dose when treating pregnant bitches.

ii) Special precautions to be taken by the person administering the medicinal product to animals

In the interests of good hygiene, persons administering the tablet directly to the dog or by adding it to the dog's food, should wash their hands afterwards

Side Effects

In very rare cases slight and transient digestive tract disorders such as vomiting and/or diarrhoea may occur. In individual cases these signs can be accompanied by nonspecific signs such as lethargy, anorexia or hyperactivity.

Interaction with other medicaments

Do not use simultaneously with piperazine compounds.

Pregnancy & Lactation

Consult a veterinary surgeon before treating pregnant animals for roundworms. The product may be used during lactation.

Symptoms & Treatment of Overdose

The product is well tolerated in dogs. In safety studies doses of 5 x or greater gave rise to occasional vomiting.

Storage Condition

Store below 30 Celcius (Room Temperature)

Shelf life

36 months

Manufacturer

Smith and Kenner Pharmaceuticals Pvt Ltd, Sy. No 72/1, Tattiannaram Village, Hayathnagar Mandal Ranga Reddy, Telangana, INDIA

Product Owner

Tradeon Band, Samrudhi, Plot No. RE 22, Gymkhana Road, MIDC, DOMBIVLI (East) District Thane, Maharashtra 421203 India

Product Registration Holder

KM Vet Pharm Sdn Bhd, No 1, Jalan Perindustrian Batu Caves 1/1, Taman Perindustrian Batu Caves, 68100 Batu Caves, Selangor Malaysia

Date of Revision

November 2020