

Veterinary Package Insert

Product Name:

Yen-AnTil 20% W/W Powder

Product Description:

White and odourless powder.

Target Species:

Swine, Others: Pigs

Pharmacodynamics:

Tilmicosin is a semi-synthetic antibiotic of the macrolide group, and is believed to affect protein synthesis. It has bacteriostatic action but at high concentrations it may be bactericidal. This antibacterial activity is predominantly against Gram-positive microorganism with activity against certain gram-negative ones and Mycoplasma of a bovine, porcine, ovine and avian origin. In particular its activity has been demonstrated against the following micro-organism:

Pigs: *Mycoplasma hyopneumoniae*, *Pasteurella multocida*, *Actinobacillus pleuropneumoniae*.

Scientific evidence suggests that macrolides act synergistically with the host immune system. Macrolides appear to enhance phagocyte killing of bacteria. Tilmicosin has been shown to inhibit in vitro the replication of the Porcine Reproductive and Respiratory Syndrome virus in alveolar macrophages in a dose dependent fashion.

Cross resistance between tilmicosin and other macrolides and lincomycin has been observed.

Pharmacokinetics:

Pigs:

Absorption: When administered to pigs via the oral route at a dose of 400 mg tilmicosin/kg feed (equivalent to approximately 21.3 mg tilmicosin/kg bodyweight/day), tilmicosin moves rapidly out of the serum into areas of low pH. The highest concentration in the serum was recorded on day 10 of medication, but concentrations above the limit of quantification were not found in 3 out of 20 animals examined. Lung concentrations increased rapidly between days 2 and 4 but no significant changes were obtained following four days of dosing. The maximum concentration in lung tissue was recorded on day 10 of medication.

When administered at a dose of 200 mg tilmicosin/kg feed (equivalent to approximately 11.0 mg/kg/day), plasma concentrations above the limit of quantification were found in 3 out of 20 animals examined. Quantifiable levels of tilmicosin were found in lung tissue with the maximum concentration being recorded on day 10 of medication.

Distribution: Following oral administration, tilmicosin is distributed throughout the body with especially high levels found in the lung and in lung tissue macrophages. It is also distributed in the liver and kidney tissues.

Biotransformation: Several metabolites are formed, the predominant one being identified as T1. However the bulk of tilmicosin is excreted unchanged.

Elimination: Following oral administration, tilmicosin is excreted mainly via the bile into the faeces, but a small proportion is excreted via the urine.

Indications:

For the control of pneumonia in swine, caused by *Actinobacillus pleuropneumoniae*, *Mycoplasma hyopneumoniae*, *Pasteurella multocida* and other organisms sensitive to tilmicosin.

Recommended Dosage:

Mixing Directions:

Yen-AnTil 20% W/W Powder must be thoroughly mixed in feed before use.

Thoroughly mix Yen-AnTil 20% W/W Powder into ratio as directed.

To ensure adequate mixing, it is recommended that a pre-blend of Yen-AnTil 20% W/W Powder be made before incorporation into the total amount of complete feed.

A dose rate of 8 to 24mg Tilmicosin / kg bodyweight / day is to be provided.

Dose Rate Table

Tilmicosin Concentration in Complete Feed (ppm)	Kg of Yen-AnTil 20% W/W Powder per 1000kg of Complete Feed
200	1.113kg
400	2.226kg

Feeding Directions:

Feed continuously as the sole ration for a 14 day period, beginning approximately seven (7) days before an anticipated disease outbreak. If clinical signs of pneumonia have already been observed, clinically ill pigs may receive supplementary treatment by injection of an approved antibiotics.

Mode of Administration:

Oral

Environmental Properties:

Sludge from waste treatment ponds must be disposed of in approved landfills for contaminated wastes. This product may accumulate in soils. DO NOT spread effluent containing Tilmicosin on the same pastures or croplands in successive years. DO NOT allow wastes from pigs treated with Tilmicosin to enter waterways. These wastes contain Tilmicosin which is highly toxic to algae.

Contraindications:

This product should not be used in any feed (supplement, concentrate or complete) containing bentonite, as bentonite in feeds may affect the efficacy of Tilmicosin.

Avoid allowing horses or other equines access to feeds containing Tilmicosin.

Warning and Precautions:

Precautions: The safety of Tilmicosin in boards intended for breeding purposes has not been established.

Safety Directions: Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with skin and eyes. Do not inhale dust. When mixing wear elbow-length PVC gloves and goggles and if dust is present half face respirator with dust cartridge or canister. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arm and face thoroughly with soap and water. After each day's use, wash gloves, goggles, respirator, and if rubber wash with detergent and warm water, and contaminated clothing.

Interactions with Other Medicaments

None known

Pregnancy and Lactation:

The safety of tilmicosin has not been established in boars used for breeding purposes.

Adverse Effects:

In very rare cases, feed intake may decrease (including feed refusal) in animals receiving medicated feed. This effect is transient.

Symptoms and Treatment of Overdose:

No symptoms of overdose have been seen in pigs fed a ration containing levels of tilmicosin up to 80 mg/kg bodyweight (equivalent to 2000 ppm in the feed or ten times the recommended dose) for 15 days.

Withdrawal Period:

REMOVE ALL MEDICATED FEED 14 days before slaughter for human consumption.

Any variation to the registered dose, frequency, duration, route, disease or target species may result in the need to extend the registered withholding period.

Packaging:

Packed in 100gm aluminium foil, 1kg aluminium foil or 25kg HDPE plastic bag

Storage Condition:

Store below 30°C in dry place, protected from light.

Shelf Life

Two years from date of manufacturing

To be used within 24 hours after first opening of container.

To be used within 24 hours after dilution.

Product Registration Holder:

YENHER AGRO-PRODUCTS SDN.BHD

(Company No.: 222455-P)

1628, Jalan IKS Simpang Ampat 1, Taman IKS
Simpang Ampat, 14100 Simpang Ampat, Seberang
Perai Selatan, Pulau Pinang, Malaysia.

Manufactured by:

DEXCHEM INDUSTRIES SDN. BHD.

(Company No.: 555717-U)

No. 6 & 8, Jalan Industri Cherok To'kun2, Taman
Industri Cherok To'kun, 14000 Bukit Mertajam, Pulau
Pinang, Malaysia.

March 2021