

Avensin Granular 40

BROAD SPECTRUM ANTICOCCIDIAL PREMIX

FORMULA:

Each 100g of premix contains

Monensin40g
Excipients up to100g

PRODUCT DESCRIPTION:

Grey or taupe granular.

PHARMACODYNAMICS:

Ionophores act as alkali metal ionophores. They complex with Na⁺ in the cell membrane to produce passive extracellular transport of K⁺ and intracellular influx of H⁺, which kills bacteria and coccidian by lowering intracellular pH. In the rumen, ionophores selectively affect Gram(+) organisms resulting in a shift to Gram(-) populations in the rumen microflora. This increases the production of propionic acid and decreases the production of acetic and butyric acids by rumen bacteria. This change in volatile acids (VFA) increases feed efficiency by reducing bacterial energy losses to CO₂ and methane, thereby increasing the energy content per unit of feed.

PHARMACOKINETICS:

Ionophores are absorbed orally. Monensin absorption is 50% in ruminants. They are rapidly and extensively metabolized by the liver and the numerous metabolites are excreted by bile and eliminated in the feces. Absorption is more complete and metabolism is slower in monogastric animals, especially horses, which may explain the greater toxicity in this species.

INDICATION:

Broiler and Layer Replacement Chickens: As an aid in the prevention of coccidiosis caused by *E. necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

MIXING DIRECTION AND RECOMMENDED DOSE:

Thoroughly mix 250-300 g of Avensin Granular 40% w/w with a sufficient quantity of feed to make one metric ton (1,000 kg) of complete chicken feed to provide 100-120 g Monensin per metric ton (1,000 kg). Feed continuously as the sole ration.

MODE OF ADMINISTRATION:

Route of Administration: ORAL

RESTRAINTS:

Do not use Avensin Granular 40% w/w in chickens producing eggs for human consumption.

CONTRAINDICATIONS:

Dogs, horses or other equines must not be allowed access to feeds containing Avensin Granular 40% w/w. Ingestion by dogs and horses may be fatal.

PRECAUTIONS:

The recommended levels of Avensin Granular 40% w/w must not be exceeded. High levels may cause decreased feed intake, poor growth, diarrhoea and death. Responses to Avensin Granular 40% w/w are unlikely if the nutrient intake is severely restricted. Feeding Avensin Granular 40% w/w to lactating cows may result in decreased milk fat percentage but there is no effect on total fat yield. Tiamulin should not be administered in poultry when Avensin Granular 40% w/w is being fed in the ration – severe growth depression or death may occur.

INTERACTIONS WITH OTHER MEDICAMENTS:

Poultry consuming Monensin should not be treated with products containing tiamulin. Severe growth depression may occur.

SAFETY DIRECTIONS:

Poisonous if swallowed. Will irritate the eyes, nose and throat and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. Do not inhale dust. When opening the container and mixing into feed wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves, goggles and half face piece respirator with dust cartridge or canister. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, contaminated clothing, goggles and respirator and if rubber wash with detergent and warm water. Additional information is contained in the Safety Data Sheet

SIDE EFFECTS:

Toxicity of ionophores when used in species for which they are approved is uncommon, unless mixing errors occur. Ionophore toxicity is due to cellular electrolyte imbalances, increased extracellular K⁺, and intracellular Na⁺ and Ca²⁺ concentrations, resulting in cellular damage and death. The increased intracellular Ca²⁺ concentration is due to the exchange of Na⁺ for Ca²⁺ by Na⁺ --Ca²⁺ exchanger; this exchange is particularly prominent in cardiac and skeletal muscles and these are usually the most severely affected. Horses are the most susceptible species to toxic effects when accidentally exposed to ionophore containing feeds.

SYMPTOM OF OVERDOSE:

The birds refused feed, had decreased water consumption and suffered severe paralysis ranging from an abnormal gait to a complete inability to move.

TREATMENT OF OVERDOSE:

If overdose is confirmed, remove the feed containing Monensin and provide proper access to drinking water to the birds.

WITHHOLDING PERIOD(S):

Poultry: Meat: Nil. Eggs: DO NOT USE in birds producing eggs for human consumption.

STORAGE CONDITIONS:

Store in a cool and dry place below 30°C.

SHELF LIFE:

2 years.

SHELF LIFE AFTER RECONSTITUTION OF PRODUCT INTO FEED:

24 hours.

DOSAGE FORMS AND PACKAGING AVAILABLE:

Medicated Premix
25 kg bag

Manufactured by
SHANDONG QILU KING-PHAR PHARMACEUTICAL CO. LTD.
21 Qinglong Road, Pingyin, Jinan 250400 China

Registration holder:
Phibro Corporation (M) Sdn Bhd
311, Level 3, Menara Axis,
2, Jalan 51A/223, Seksyen 51A,
46100 Petaling Jaya, Selangor, Malaysia

Revision Date of Package Insert: Aug 15, 2025