

NEO-MYC 500mg/g Water Soluble Powder

NEO-MYC 500mg/g Water Soluble Powder appears to be a white to light yellow powder.

Composition:

Each g contains

Neomycin (as Neomycin Sulfate)

500 mg

Pharmacodynamics:

Aminoglycosides enter susceptible bacteria by oxygen-dependent active transport (making anaerobes impervious to them) and by passive diffusion. Once the antibiotic has gained access, it binds irreversibly to a receptor protein on the 30S ribosomal subunit and blocks the formation of a complex that includes mRNA, formylmethionine, and tRNA. As a result, the tRNA is translated incorrectly, producing a nonfunctional protein. Aminoglycosides also disrupt protein synthesis by disruption of polysomes and may prevent the initiation of DNA replication.

Pharmacokinetics:

In general, aminoglycosides and apramycin are very poorly absorbed from oral administration in adult animals, including cattle, chickens, and pigs. Aminoglycosides are distributed primarily into the extracellular space and over time accumulate in tissues. The amount of antibiotic in most tissues appears to be dependent on the total dose administered over time rather than the size of each individual dose. Aminoglycosides do not distribute well across membrane barriers and, therefore, are not found at high concentrations in brain tissue, cerebrospinal fluid, ocular fluid, or respiratory secretions.

Indication:

Goats, Cattle (excluding veal calves), Swine and Sheep: For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate.

Turkey (growing): For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate.

Recommended Dosage:

For the treatment and control of colibacillosis (bacterial enteritis) caused by E.Coli susceptible to neomycin sulfate in goats, cattle (excluding veal calves), swine and sheep: Administer 10 mg neomycin sulfate per 0.45 kg of body weight per day (22mg per kg) in divided doses for a maximum of 14 days.

For the control of mortality associated with E. coli organisms susceptible to neomycin sulfate in growing turkey: Administer 10 mg neomycin sulfate per 0.45 kg of body weight per day (22mg per kg) for a maximum of 5 days.

Add to drinking water. Not for use in liquid supplements. Prepare a fresh solution daily.

Mode of Administration:

Oral Administration.

Contraindications:

The medical contraindications have been selected on the basis of their potential clinical significance. Except under special circumstances, this medication should not be used when the following medical problems exist:

- Dehydration, hypovolemic
- Hypersensitivity to aminoglycosides
- Renal dysfunction

Warnings and Precautions:

If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. Treatment should continue 24-48 hours beyond remission of disease symptoms, but not to exceed a total of 14 days for goats, cattle (excluding veal calves), swine and sheep, and 5 days for turkeys. Discontinue treatment 1 day prior to slaughter in cattle (excluding veal calves), 2 days in sheep and 3 days in goats and swine. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

Interactions with Other Medicaments:

The following drug interactions have been selected on the basis of their potential clinical significance: aminoglycosides (two or more concurrently), calcium channel blockers, calcium, halothane anesthesia, iron (supplemental), ketorolac, phenylbutazone, nonsteroidal anti-inflammatory drugs (Nsaids), loop diuretics, nephrotoxic medications, ototoxic medications and neuromuscular blocking agents.

Statement on Usage During Pregnancy and Lactation:

Because of poor lipid solubility, aminoglycosides have relatively poor penetration from plasma into milk.

Adverse Effects / Undesirable Effects:

The following side/adverse effects have been selected on the basis of their potential clinical significance.

Side effects needing medical attention: Nephrotoxicity, ototoxicity, auditory; ototoxicity, vestibular and neuromuscular blockade.

Side effects needing medical attention only if they continue or are bothersome:

Diarrhea in calves and pigs given oral doses of neomycin that are higher than the label dose.

Overdose and Treatment:

Symptoms of overdose includes nephrotoxicity, neuromuscular blockade, ototoxicity (auditory), and ototoxicity (vestibular).

For neuromuscular blockade

• Administration of edrophonium, 0.5 mg/kg, will reverse neuromuscular blocking effects. Administration of calcium chloride at 10 to 20 mg /kg, calcium gluconate at 30 to 60 mg/kg, or neostigmine at a dose of 100 to 200 mcg per kg of body weight can also reverse muscle response depression and associated dyspnea.

For renal toxicity

- Aminoglycoside administration should be immediately discontinued.
- Polyionic electrolyte fluid therapy should be initiated to stimulate diuresis.

Withdrawal Period (s):

Cattle excluding veal calves: 1 day before slaughter.

Sheep: 2 days before slaughter.

Swine and goats: 3 days before slaughter.

Turkey: zero days before slaughter.

Storage Conditions:

Store below 30°C. Protect from heat and sunlight exposure.

Keep medicine out of reach of children. (Jauhkan ubat dari kanak-kanak).

Any unused product or packaging should be disposed of in accordance with local and national legislation.

Shelf-life (after first opening of container): 7 days.

Shelf-life (after reconstitution/ dilution): 24 hours.

Packaging Available:

500g

Date of revision:

12-06-2021

Manufacturer:

NAM PHARMA SDN BHD

5, Lebuhr Perusahan Klebang 11,

Taman Perindustrian Antarabangsa IGB,

31200 Chemor, Perak, Malaysia.

Product Registration Holder

F.E VENTURE SDN BHD

Lot 3&5 Jalan PJS 11/8,

Bandar Sunway,

46150 Petaling Jaya, Selangor, Malaysia.