

HEMOCARE NOFLUX[®] LOZENGES

Ingredient(s):

Each lozenge contains:

Lysozyme Chloride (derived from egg white)	80mg
Pyridoxine Hydrochloride	40mg

Product Description:

A white color round lozenge.



Pharmacodynamics

1. Lysozyme is an enzyme having a mucopolysaccharide lysing action. It enhances the effect of antibiotics to destroy or weaken cell bacteria.
2. Lysozyme also accelerates immuno-bacteriolysis and activates the phagocytosis activity of leukocytes to destroy bacteria and viruses.
3. Lysozyme is widely distributed in tears, nasal discharge, saliva and also breast milk.
4. Lysozyme extracted from egg-white is used clinically for the intensification of natural defense mechanisms, restoration of damaged tissue and anti-inflammation.
5. The body needs pyridoxine hydrochloride in order to make cells of the immune system. Pyridoxine hydrochloride helps to improve immune response to the increase in production of antibodies and also helps in communicative interactions between cytokines and chemokines.

Pharmacokinetics

1. With a single 900mg (potency) oral dose of lysozyme chloride, the peak plasma concentration occurred about 30 minutes to 1 hour after administration.
2. Pyridoxine hydrochloride is absorbed rapidly from the upper intestine regardless of the size of the dose given.
3. Absorption may also occur from the ileum and to a small extent from the colon. These observations suggest the possibility that pyridoxine may be absorbed by diffusion.
4. Pyridoxine is rapidly converted in the liver to pyridoxine phosphate, pyridoxal phosphate and pyridoxamine phosphate via oxidation, this causes the release of pyridoxal and so pyridoxal phosphate to the general circulation where it reaches other organs chiefly as circulating pyridoxal.
5. In urine, pyridoxine was excreted primarily unchanged with a small amount of only one metabolite, most likely 4-pyridoxic acid.
6. Oral liquid: Absorption and Distribution: following administration of [1]-labelled egg white lysozyme (2mg/kg) into the intestinal tract of rats, peak serum and lymphatic concentration occurred at 30 minutes.

Indication:

For the adjunctive treatment of sore throat.

Recommended Dose:

Adult: 1 lozenge to be sucked or dissolved in the mouth once or twice daily or according to doctor's prescription.

Route of Administration:

For oral use.

Contraindication:

Patients hypersensitive to egg-white. Patients with a history of hypersensitivity to any ingredients of Homecare Noflux[®] Lozenges.

Warning and Precautions:

1. Patients with an allergic predisposition manifested as atopic dermatitis, bronchial asthma, drug allergy or food allergy to egg or pyridoxine hydrochloride (Vitamin B6).
2. Abuse of pyridoxine may be responsible for sensory neuropathy that occurs with high doses of pyridoxine (> 200 mg/day) for a long period of time (months to years). This is usually reversible when the treatment is stopped.
3. Please consult the doctor if condition does not improve after 5 days of treatment.

Pregnancy and Lactation

It is best not to use this medicine during pregnancy and breastfeeding.

Side Effects:

1. May cause diarrhoea, anorexia, stomach discomfort, nausea and vomiting.
2. May cause hypersensitivity reactions such as skin rashes.
3. In rare case, shock may occur with administration of this drug. Therefore, patients should be carefully monitored during Homecare Noflux[®] Lozenges.

Symptoms and Treatment of Overdose:

Symptoms: Diarrhoea, anorexia, gastric discomfort.

Treatment: Stop administration.

Storage Condition:

Store at temperature below 30°C. Protect from light and moisture.

Pack Size:

10's x 50

Shelf Life:

2 years from the date of manufacture.

HEMOCARE NOFLUX[®] LOZENGE

Bahan aktif:

Setiap lozenge mengandungi:

Lysozyme Chloride (berasal dari putih telur)	80mg
Pyridoxine Hydrochloride	40mg

Indikasi:

Untuk rawatan tambahan sakit tekak.

Dos:

Dewasa: 1 lozeng untuk dihisap atau dilarutkan di mulut sekali atau dua kali sehari atau berdasarkan preskripsi doktor.

HEMOCARE NOFLUX[®] 消炎錠

成分:

每錠含:

Lysozyme Chloride (萃取自蛋白)	80mg
Pyridoxine Hydrochloride	40mg

適應症:

適用於咽喉痛的治療。

劑量:

成人: 1片, 每天一次至兩次, 或根據醫生的處方將其溶化在口腔中。



Y. S. P.

Manufacturer and Product Registration Holder:

Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,

Kawasan Perindustrian Bandar Baru Bangi,

43000 Kajang, Selangor Darul Ehsan, Malaysia.

Ordering Line: 1 800 88 3027

Product Info: 1 800 88 3679