

QUINOFEED 200G/KG GRANULES

Description and Composition

QUINOFEED 200G/KG GRANULES is a brown granules. Each kg contains 200g of Enrofloxacin as its active ingredient.

Pharmacodynamics

Enrofloxacin, or 1-Cyclopropyl-6-fluoro-7-(4-ethyl-1-piperazinyl)-1,4-dihydro-4-oxo-3-quinolin-2-carboxylic acid, belongs to fluoroquinolone family which is a subfamily of quinolone. Enrofloxacin is the first fluoroquinolone patented in 1984. Quinolones have an action on bacterial topoisomerase. Enrofloxacin is efficient on most gram-negative and gram-positive bacteria but not efficient on anaerobic bacteria.

Pharmacokinetics

Trials in all species showed that enrofloxacin is absorbed readily and distributes to all organs and tissues following oral or parenteral administration. In pigs, when the same dose was administered by the oral or i.m. route, similar rates of absorption and subsequent concentrations of drug in tissues were observed. The highest drug concentration initially occurred in the liver in all species. In all species studied, elimination was primarily via the urine and bile/faeces. In food animal species, ciprofloxacin was the main metabolite recovered. The liver is considered the primary site of enrofloxacin metabolism in these species, forming ciprofloxacin through oxidative dealkylation. Additional metabolites occurred but comprised less than 10% of the total residue.

Indication

Pigs: Treatment of diarrhoea, enterotoxigenic edema and colibacillosis caused by *E.coli* susceptible to Enrofloxacin. Treatment of respiratory disease caused by *Mycoplasma hyopneumoniae*, *Pasteurella multocida* and *Pleural pneumonia (Actinobacillus pleuropneumoniae)* susceptible to Enrofloxacin.

Recommended Dose

Pigs: Diarrhoea, enterotoxigenic edema and colibacillosis: 0.25kg of product per tonne of feed per day which is equivalent to 50ppm of Enrofloxacin uniformly mixed for 5 days. May be given up to 10 days depending on symptoms. Respiratory disease: 0.75kg of product per tonne of feed per day which is equivalent to 150ppm of Enrofloxacin uniformly mixed for 7 days. May be given up to 10 days depending on symptoms.

Route of Administration

For oral administration via feed.

Contraindication

Do not use on animals that have had a shock and hypersensitivity to the drug. Do not administer to animals with liver or kidney disorders.

Warning and Precautions

Special warning for each target species: None known.

Special precautions for use in animal:

Do not use on animals that have had a shock and hypersensitivity to the drug. Do not administer to animals with liver or kidney disorders. Use as directed by your veterinarian. Please read the label carefully before use. Do not use arbitrarily because safety and efficacy have not been established other than the specified breed. In order to prevent misuse and the development of tolerance, please record the history of use of veterinary drugs without omission. Do not administer drugs of the same ingredient simultaneously while administering this drug. If the withdrawal period is not observed, drugs may remain in livestock foods, such as meat, so make sure to accurately count and follow the withdrawal period.

Special precautions to be taken by the person administering the drug to the animals:

The operator should wear protective equipment such as gloves, masks, and protective clothing to avoid direct skin contact or inhalation. In case of contact with skin or eyes when handling the drug, wash it off immediately and seek medical attention if any abnormal symptoms are found.

Interaction with Other Medicaments

Do not use with macrolide or tetracycline antibiotics. The absorption rate *in vivo* may decrease when mixed with magnesium, aluminum, or calcium ion-containing preparations. When administered together with theophylline and caffeine, the blood concentration of theophylline and caffeine can be increased. Probenecid can increase the concentration in the blood by preventing the discharge of tubules of this drug. When combined with Cyclosporine, Cyclosporine may exacerbate renal toxicity. When used in combination with nonsteroidal anti-inflammatory drugs, convulsions may rarely develop.

Pregnancy and Lactation

The attributes of fluoroquinolones make them likely to cross the placenta in many species; however, adverse effects have not yet been reported when fluoroquinolones have been administered to pregnant animals. Because of the risk of producing arthropathies in immature animals, it has been recommended that significant levels of fluoroquinolones in the milk of nursing animals be avoided. Fluoroquinolones can be distributed into milk, sometimes at a higher concentration than in plasma, but it is not known under what conditions significant amounts might be absorbed by nursing animals. The effects of enrofloxacin on swine reproductive performance, pregnancy and lactation have not been adequately determined.

Side Effects

When administered to animals in the growing season, joint abnormalities (lameness, pain, cartilage failure) may appear. Gastrointestinal disorders (vomiting, loss of appetite, diarrhoea, abdominal pain, etc.) may rarely occur. Central nervous system disorders (vertigo, anxiety, headache, depression, ataxia, seizures, etc.) appear and do not use in animals with central nervous system disorders (seizures, etc.) as symptoms may worsen. Hypersensitivity reaction and crystal urine may appear.

Symptoms and Treatment of Overdose

When overdose (more than 10 times), abnormalities such as vomiting and poor feeding may occur.

Storage Condition

Store below 30°C.

Shelf life

3 years. To be used within 24 hours after mixing into feed.

Withdrawal Period

Pigs: 10 days

Packing

10kg and 25kg.

Revision date: 9.7.2021

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No.1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 E-mail: info@rangepharma.com