

MK essentia healthcare series Pain Relief Cream



Active ingredients: Menthol 6%w/w, Methyl Salicylate 13%w/w

Preservative: Chlorocresol 0.2%w/w

Product Description:

White colour, smooth application, sticky and characteristic odor cream.

Indication:

For the symptomatic relief of muscular aches and pains, and stiffness including backache, lumbago and fibrositis.

Direction of Use:

Gently massage a sufficient amount of the cream onto affected area. Not recommended for children below 12 years old.

Route of Administration: Topical

Pharmacodynamics

Menthol

Menthol dilates the blood vessels causing a sensation of coldness, followed by an analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetration of drugs when applied on the skin, to give a faster onset of action.

Methyl Salicylate

Methyl salicylate is a salicylic acid derivative. Salicylates inhibit cyclooxygenase, thereby reducing the formation of prostaglandins and cause platelet dysfunction. Methyl salicylate is used topically as a counter irritant. Upon application, it is absorbed through the skin and is applied for the relief of the pain in rheumatic conditions and painful muscle or joints.

Pharmacokinetics

Menthol

After absorption, menthol is excreted in the urine and bile as a glucuronide.

Methyl Salicylate

May be absorbed through intact skin. The absorption of topical salicylates is proportional to the surface area involved, duration of exposure, concentration and skin integrity. Absorption characteristics of salicylates vary with the dose, formulation and route of administration. Percutaneous absorption is enhanced by exercise, heat, occlusion or disruption of the integrity of the skin or application to large areas of skin. Both the rate and extent of absorption increases after repeated application; increasing the bioavailability. Methyl salicylate is extensively metabolised to salicylic acid in the dermal and subcutaneous tissues after topical application. At therapeutic levels, the half-life of salicylates is 2–4 hours. As salicylate level increase into the toxic range, the half-life can be greater than 18 hours.

Contraindications:

Do not use in patients with aspirin or salicylate idiosyncrasy. Patient with known hypersensitivity to any of the ingredients in the product.

Warnings and Precautions:

This product contains methyl salicylate and when applied or rub on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding.

Initially apply a small amount to a limited skin area and observe whether there is discomfort or itch before applying to larger skin area. Discontinues use if irritation develops. Do not apply to open wounds or damaged skin. Avoid excessive use during pregnancy and lactation. Do not apply over large areas of the body or bandage or use with heating pads. If symptoms persist, consult your doctor.

For external use only. Keep out of reach of children.
Jauhkan daripada capaian kanak-kanak.

Front Dimension: 190mm(W) x 150mm (H)

Effects on Ability to Drive and Use Machines: Not known

Interactions with Other Medicaments:
Warfarin.

Statement on Usage During Pregnancy and Lactation:
Safety for use in pregnancy has not been established; therefore the potential benefit of the product should be weighed against the possible risks to the mother and child.

Adverse Effects/ Undesirable Effects:
Skin irritation, contact dermatitis, rash, itching, redness or swelling, burning or stinging sensation may occur.

May cause hypersensitivity/ allergic reactions in some individuals with sensitive skin.

Symptoms and Treatment of Overdose:
Overdose is unlikely when used as recommended.
If applied to a large area of skin, or in the unlikely event of oral ingestion, the product may cause systematic effects depending on the amount absorbed.
Treatment of overdose should be supportive and symptomatic.

Storage Conditions:
Store below 30°C. Protect from direct sunlight.

Shelf Life: Please refer to the packaging. Do not use beyond the expiry date.

Dosage Form: Cream

Packing: 30g in aluminium tube

MAL XXXXXC

Product Registration Holder:
MegaLive BioSciences Sdn Bhd
1st Floor, No. 13, Jalan Selasih, Taman Semarak,
09000 Kulim, Kedah, Malaysia

Manufacturer:
Syarikat Wen Ken Drug Sdn Bhd
24, Jalan Lambak, Taman Johor,
81200 Johor Bahru, Johor, Malaysia.

Date of Revision: 12.08.2021