



SNAKE BRAND SPORT OIL

Product Description Name & Strength of Active Substance(s)

Bright green emulsion with characteristic odor	
Methyl Salicylate	31% v/v
Menthol	1% w/v
Camphor	1% w/v
Eucalyptus Oil	1% v/v

Pharmacodynamics

Camphor

Camphor is a stimulant, rubefacient, antipruritic and feebly antiseptic. When in combination with other ingredient like menthol, methyl salicylate, it becomes ideal for neuralgia and other painful areas.

Menthol

Menthol dilates the blood vessels causing a sensation of coldness, followed by an analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetration of drugs when applied on the skin, to give a faster onset of action.

Methyl Salicylate

Methyl salicylate is a salicylic acid derivative. Salicylates inhibit cyclooxygenase, thereby reducing the formation of prostaglandins, and cause platelet dysfunction. Methyl salicylate is used topically as a counter-irritant. Upon application, it is absorbed through the skin and is applied for the relief of pain in rheumatic conditions and painful muscle or joints.

Pharmacokinetics

Camphor

- Camphor is readily absorbed from all administration sites. Camphor is well absorbed after dermal exposure.
- It is hydroxylated in the liver to yield hydroxycamphor metabolites which are then conjugated with glucuronic acid and excreted in the urine.
- Camphor crosses the placenta.

Menthol

- After absorption, menthol is excreted in the urine and bile as a glucuronide.

Methyl Salicylate

- May be absorbed through intact skin.
- The absorption of topical salicylates is proportional to the surface area involved, duration of exposure, concentration and skin integrity.
- Absorption characteristics of salicylates vary with the dose, formulation and route of administration.
- Percutaneous absorption is enhanced by exercise, heat, occlusion, or disruption of the integrity of the skin or application to large areas of skin.
- Both the rate and extent of absorption increases after repeated application; increasing the bioavailability.
- Methyl salicylate is extensively metabolized to salicylic acid in the dermal and subcutaneous tissues after topical application.
- At therapeutic levels, the half-life of salicylates is 2-4 hours. As salicylate level increase into the toxic range, the half-life can be greater than 18 hours.

Indication

Provide symptomatic relief for mild to moderate muscular and joint aches and pains, muscle cramps, shoulder aches and stiff neck.

Recommended Dosage Adults & children above 2 years old : Use before or after exercise. Apply to the affected area not more than 3 times daily.Rub/Massage gently. For external application only.

Mode of Administration Topical

Contraindications

- Do not use in patients with aspirin or salicylate idiosyncrasy.
- Hypersensitivity to any of the ingredients in the product.
- Contraindicated in infants below 2 years of age.

Warnings and Precautions

This product contains methyl salicylate and when applied or rub on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding. Topical analgesic preparations containing methyl salicylate should be used with caution in patients at increase risk of developing salicylate adverse effects. Children suffering from flu, chickenpox or fever should avoid using this product because salicylates may induce Reyes Syndrome.

WARNING : THIS PRODUCT CONTAINS CAMPHOR. CAN CAUSE CONVULSION. CONTRAINDICATED IN CHILDREN BELOW 2 YEARS OF AGE. CAUTION MUST BE EXERCISED WHEN OLDER CHILDREN ARE TREATED. AVOID DIRECT APPLICATION INTO THE NOSTRILS.

It is dangerous to place any camphor-containing product into the nostril of children. A small amount applied this way may cause immediate collapse.

Initially apply a small amount to a limited skin area and observe whether there is discomfort or itch before applying to a larger skin area. Discontinue use if irritation develops.

Avoid contact with the eyes and mucous membranes.

Do not apply to open wounds or damaged skin.

Avoid excessive use during pregnancy and lactation.

Do not apply over large areas of the body or bandage or use with heating pads.

If symptoms persist, consult your doctor.

For external use only.

Effects on Ability to Drive and Use Machines None known

Interactions with Other Medicaments

This product contains methyl salicylate and when applied or rubbed on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joints pains may increase the chances of bleeding.

Statement on Usage during Pregnancy and Lactation

Safety for use in pregnancy has not been established; therefore the potential benefit of the product should be weighed against the possible risks to the mother and child.

Adverse Effects / Undesirable Effects

Headache, dizziness, nausea and vomiting, skin irritation, contact dermatitis, rash, itching, redness or swelling, burning or stinging sensation may occur. May cause hypersensitivity/allergic reactions in some individuals with sensitive skin.

Overdose and Treatment

Symptoms of methyl salicylate overdose

Salicylate intoxication can occur after ingestion or topical application of methyl salicylate mild chronic salicylate intoxication, or salicylism, usually occurs only after repeated use of large doses. Salicylism can also occur following excessive topical application of salicylates. Symptoms include dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache, and confusion, and may be controlled by reducing the dosage. Symptoms of more severe intoxication or of acute poisoning following overdose include hyperventilation, fever, restlessness, ketosis, and respiratory alkalosis and metabolic acidosis. Depression of the CNS may lead to coma; cardiovascular collapse and respiratory failure may also occur.

Symptoms of camphor/menthol overdose

Ingestion of significant quantities is reported to cause symptom: severe abdominal pain, nausea, vomiting, vertigo, ataxia, drowsiness, and coma; instant collapse in infants after the local application of menthol/camphor to their nostrils. Seizures may be the first clinical sign of severe toxicity of camphor; however, seizures are usually self-limited. Severe toxicity of camphor can result in delirium, visual hallucinations, cerebral edema, and status epilepticus. Systemic toxicity may include hypotension, tachycardia, respiratory failure, and death.

Treatment

The stomach should be emptied by gastric lavage or administration of oral activated charcoal fluid and electrolyte management is the mainstay of treatment with the immediate aim of correction of acidosis, hyperpyrexia, hypokalemia and dehydration if present. Any convulsions must be controlled first through supportive care including anticonvulsant therapy.

Storage Conditions

Store below 30°C.
KEEP OUT OF REACH OF CHILDREN.
JAUHI DARIPADA CAPAIAN KANAK-KANAK.

Dosage Forms and Packaging Available

30ml, 60ml
Clear glass bottle with aluminum screw cap packed in unit box



Name & Address of Manufacturer

The British Dispensary (L.P.) Co., Ltd.
272 Teparuk Road, Samut Prakan 10270, Thailand.
Tel. (662) 6451333 Fax. (662) 6450700
Email: customerservice@britishdispensary.com

Product registration holder in Malaysia

Carepro Marketing Sdn. Bhd. (735191-H)
No. 54, Jalan Bp 5/11, Bandar Bukit Puchong, 47100
Puchong, Selangor, Malaysia.
Tel: 03-80628889 Email: info@carepro.com.my

Date of Revision of Leaflet
06 / Sep / 2021

Front : 11x18.5

Back : 11x18.5