

PACKAGE INSERT

Brand or product name	WeissNicarb 25% w/w Premix Powder
Name and Strength of Active Substance	Nicarbazine.....250 mg/g
Product Description	Green Yellow Powder
Pharmacodynamics	Nicarbazin is a broad-spectrum chemical anticoccidial agent used to prevent disease and treat infection in food-producing poultry. The drug residue is composed of equimolar quantities of 4,4-dinitrocarbanilide (DNC) and 2-hydroxy-4,6-dimethyl pyrimidine (HDP), with DNC serving as the primary component for tracking of the residue in edible tissues, including fat, muscle, liver, and kidney, and in eggs.
Pharmacokinetics	Nicarbazin, when ingested, is rapidly split in its two components 2-hydroxy-4,6-dimethylpyrimidine (HDP) and dinitrocarbanilide (DNC) which behave independently. Liver is the target tissue. Dinitrocarbanilide (DNC) residues decline rapidly from tissues following nicarbazin withdrawal. Dinitrocarbanilide (DNC) appears as the marker residue. 2-hydroxy-4,6-dimethylpyrimidine (HDP) -related residues are much lower (by two orders of magnitude) than those derived from dinitrocarbanilide (DNC). The metabolic pathways in the turkey are similar to those in the chicken. Nicarbazin as such cannot appear as residue in tissues; only its two individual components may generate residues.
Environmental Properties (for product used directly in the environment e.g. medicine for fish)	N/A
Indication	Aid in preventing outbreaks of cecal (<i>Eimeria tenella</i>) and intestinal (<i>E. acervulina</i> , <i>E. maxima</i> , <i>E. necatrix</i> and <i>E. brunetti</i>) coccidiosis.
Recommended Dosage	125 grams Nicarbazine per tonne of feed (500 grams WeissNicarb 25% w/w Premix Powder).
Mode of Administration	Oral Administration
Contra-indications	Do not use in know cases of hypersensitivity to active ingredient.
Warnings and Precautions	Feed continuously as sole ration from time chicks are placed on litter until past the time when coccidiosis is ordinarily a hazard. Do not use as a treatment for coccidiosis. Do not use in flushing mashers. Do not feed to laying hens. It is forbidden for layers or breeders.

Interactions with Other Medicaments	No incompatibilities or interactions of nicarbazin with feedingstuffs, carriers or other approved additives are to be expected.
Treatment on usage during pregnancy and lactation	There were isolated occurrences of slightly reduced litter size at birth or depressed body weight gain during lactation at the highest dose level. At 600 mg/kg b.w. per day some rats died (7/25), maternal food intake and body weights were reduced, foetal body weights were decreased and there was reduced ossification suggesting retarded foetal development. The NOEL for nicarbazin was considered to be 200 mg/kg b.w. per day for maternal and foetal toxicity.
MRL	<u>Chicken</u> Muscle, liver, kidney and fat/skin - is 200 µg/kg.
Adverse Effects/Undesirable Effects	Nicarbazin has adverse effects on the hatchability and the quality of the eggs. At 125 mg/kg of feed, nicarbazin caused a reduction in egg production of 65% and completely inhibited the hatchability of fertilised eggs. At high dietary concentrations of nicarbazin (1600 mg/kg feed), testis development was retarded. Nicarbazin was fed at 0, 20, 50, 100 and 125 mg/kg feed to broiler breeders and no depression in egg production (with the exception of 125 mg/kg), egg weight or fertility from feeding these levels were observed. However, a linear decrease in hatchability was observed together with an increase in dinitrocarbanilide (DNC) amount in egg yolks and egg-shell pigmentation as the level of nicarbazin increased. At 125 mg/kg nicarbazin reduced egg production, and produced a more severe decrease in hatchability and a complete bleaching of brown-shell eggs by the third day of treatment. However, after seven days of nicarbazin withdrawal from the feed, pigmentation returned to pre-treatment level.
Overdose and Treatment	The primary toxic effects of high oral doses of nicarbazin are decreased body weight gain and reduced feed conversion efficiency. 400 mg nicarbazin/kg feed, there was decreased body weight gain and lower feed efficiency. Nicarbazin fed to chickens at 600 mg per kg feed for 12 weeks produced no signs of toxicity or abnormal behaviour and did not increase mortality, but the treatment decreased body weight gain and lowered feed efficiency. Increased mortality occurred at doses of 1500 and 2000 mg nicarbazin/kg feed. Other toxic effects at 800 mg/kg feed included anaemia and fatty change and haemosiderin deposits in the liver.
Incompatibilities(for injections only)	N/A

Withdrawal period(s)	4 days before slaughter.
Storage conditions (may be omitted if the information is stated on the label or outer carton labels)/(should be supported by stability data)	Store below 30°C (room temperature).
Dosage forms and packaging available	Dosage form: Powder Packaging: 25 kg
Name and address of manufacturer	Weissen Company (M) Sdn. Bhd. 15 & 17, Jalan Metro Perdana Barat 12, Kepong Sri Edaran Ind. Park, 52100 Kuala Lumpur, Malaysia.
Name and address of registration holder	Weissen Company (M) Sdn. Bhd. Lot 105, No. 84, Jalan Velox 23, Taman Velox, 48000 Rawang, Selangor, Malaysia.
Date of revision of Package insert	02.10.2024
Shelf-life of the veterinary product as packaged for sale	24 months
Shelf-life after reconstitution in feed	24 hours