

187 mm

19 mm

L-R

4 mm

In healthy young cats aged 7- 8 months, oral robenacoxib (Onsior tablets) administered at overdoses of up to 5 times the maximum recommended dose (2.4 mg, 7.2 mg, 12 mg robenacoxib/kg bodyweight) for 6 months was well tolerated. A reduction in body weight gain was observed in treated animals. In the high dose group kidney weights were decreased and sporadically associated with renal tubular degeneration/regeneration but not correlated with evidence of renal dysfunction on clinical pathology parameters.

The interchangeable use of Onsior tablets and Onsior solution for injection in 4-month old cats at overdoses of up to 3 times the maximum recommended dose (2.4 mg, 4.8 mg, 7.2 mg robenacoxib/kg orally and 2.0 mg, 4.0 mg and 6.0 mg robenacoxib/kg subcutaneously) resulted in a dose-dependent increase of sporadic oedema at the injection site and minimal to mild subacute/chronic inflammation of the subcutaneous tissue. A dose-dependent increase in the QT interval, a decreased heart rate and corresponding increased respiratory rate were observed in laboratory studies. No relevant effects on body weight, bleeding time or evidence of any gastrointestinal, kidney or liver toxicity were observed.

In overdose studies conducted in cats, there was a dose-dependent increase in the QT interval. The biological relevance of increased QT intervals outside of normal variations observed following overdose of robenacoxib is unknown. No changes in the QT interval were observed after single intravenous administration of 2 or 4 mg/kg robenacoxib to anaesthetised healthy cats.

As with any NSAID, overdose may cause gastrointestinal, kidney, or liver toxicity in sensitive or compromised cats. There is no specific antidote. Symptomatic supportive therapy is recommended and should consist of administration of gastrointestinal protective agents and infusion of isotonic saline.

WITHDRAWAL PERIOD
Not applicable.

SPECIAL PRECAUTIONS FOR STORAGE
Store below 25°C

PACKAGE QUANTITY
Cardboard box containing 6 x 1, 12 x 1, 30 x 1 or 60 x 1 tablets in Alu/Alu perforated unit dose blisters.
Not all pack sizes may be marketed.

MANUFACTURED BY
Elanco France
26, rue de la chapelle
68330 Huringue
France.

MARKETING AUTHORISATION HOLDER
Elanco Malaysia Sdn Bhd
Unit 5.04, Level 5 & 6, Tower Block, The Bousteador,
No. 10 Jalan PJU 7/6, Mutiara Damansara,
47800 Petaling Jaya, Selangor, Malaysia.

DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED
March 2022

MAL21126017HA
CONTROLLED MEDICINE

Onsior, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.
© 2021 Elanco or its affiliates

Reverse of
2D code area
to be kept print free



320 mm