

Gleptoferron Labiana 200 mg/ml

Solution for Injection

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder and manufacturer responsible for batch release:

LABIANA LIFE SCIENCES S. A.

C/Venus, 26

08228 Terrassa (Barcelona). Spain.

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Gleptoferron Labiana 200 mg/ml Solution for Injection.

3. STATEMENT OF THE ACTIVE SUBSTANCE AND OTHER INGREDIENT

Dark, brown slightly viscous solution.

Active substance:

Iron(III)200.0 mg/ml

(as Gleptoferron)..... 532.6 mg)

Excipients:

Phenol...(As preservative).....5 mg/ml

4. INDICATION

For the prevention and treatment of iron deficiency anaemia.

5. CONTRAINDICATIONS

Do not use in case of:

- Hypersensitivity to gleptoferron complex or any of the excipients.
- In animals with hepatic and/or renal disease.
- Do not administer to piglets suspected to suffer from deficiency of vitamin E and/or selenium.
- Do not use in clinically diseased animals, especially not in case of diarrhoea.

6. ADVERSE REACTIONS

There are normally no undesirable side effects associated with the use of the product. Sometimes a slight staining of muscle tissue may occur at the administration site. Occasional deaths have occurred in piglets following the administration of parenteral iron preparations. These deaths have been associated with maternal dietary deficiency of vitamin E and/or selenium.

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

Very rarely piglet deaths have been reported which have been attributed to an increased susceptibility to infection due to temporary blocking of the reticuloendothelial system.

The frequency of adverse reactions is defined using the following convention:

- Very common (more than 1 in 10 animals displaying adverse reaction(s) during the course of one treatment).
- Common (more than 1 but less than 10 animals in 100 animals).
- Uncommon (more than 1 but less than 10 animals in 1,000 animals).
- Rare (more than 1 but less than 10 animals in 10,000 animals).

7. TARGET SPECIES

Neonatal pigs.

8. DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION

Use only automatic syringe equipment.

Swab the septum before use. The product is administered as a single 1 mL (200 mg iron) dose by deep intramuscular injection into the hind limb midway between the stifle joint and the base of the tail. Injections should be administered as follows:

For the prevention of iron deficiency anaemia: not later than the third day of life.

For the treatment of iron deficiency anaemia: at the onset of clinical anaemia normally within the first three weeks of life.

It is advisable to stretch the skin at the injection site to minimize leakage after withdrawal of the needle.

9. ADVICE ON CORRECT ADMINISTRATION

Do not use Gleptoferron Labiana 200 mg/ml solution for injection if you notice visible signs of deterioration.

10. WITHDRAWAL PERIOD

Meat and offal: zero days.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

40C/75%RH, 30C/65%RH, 25C/60%RH.

Shelf life after first opening the container: 28

days. Do not use this veterinary medicinal product after the expiry date which is stated on the container after "EXP". The expiry date refers to the last day of that month. Shelf life as packaged for sale.

12. SPECIAL WARNING(S)

Special warnings for each target species:

The sachet in the low-density polyethylene collapsible bottles with a nominal capacity of 100 ml should not be opened until the product is required for use.

Producto/Product	Gleptoferron Labiana	Colores	Control de calidad
Dosis/Strength	100 y 200 ml	 P Black	Fecha
Envase/Pack Size	Prospecto	 Troquel	Registros
Material/Material			Fecha
Dimensiones/Dimensions	126 x 210 mm		Márqueting
Laetus/Laetus			Fecha
Código/Item Code	71204/001		
Fecha edición/Date of Issue			
Tipografía/Fonts	Segoe UI / 8 pt		
Programa/Software	InDesign CC	OBSERVACIÓN Tonos de color no válidos para impresión	
Cliente	 LABIANA	Proveedor	

Special precautions for use in animals:

Normal aseptic injection techniques should be practised.

Avoid the introduction of contamination during use.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known hypersensitivity to iron dextran, or those with haemochromatosis should avoid contact with the product.

Care should be taken to avoid accidental self-injection, as well as contact with the eyes and mouth.

In case of accidental injection, seek medical advice immediately and show the package leaflet or label to the physician.

Wash hands after use.

The sachet in the low-density polyethylene collapsible bottles with a nominal capacity of 100 ml and 200 ml should not be opened until the product is required for use.

Interaction with other medicinal products and other forms of interaction:

There are no known interactions between the product and other medicaments. There are no known other forms of interaction. Do not mix with other products prior to administration.

Overdose (symptoms, emergency procedures, antidotes):

Overdosage with the product is unlikely to result in signs of intoxication.

Incompatibilities:

None known.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Medicines should not be disposed of via wastewater or household waste. Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

15. OTHER INFORMATION

• Low-density polyethylene collapsible bottles (LDPE) with a nominal capacity of 100 ml and 200 ml closed with a chlorobutyl rubber closure and an aluminium ring seal. Each bottle is sealed in a sachet. The sachet is a polyester/polyethylene laminate. Do not open the sachet until ready to use the veterinary medicinal product.

• High-density polyethylene collapsible bottles (HDPE) with a nominal capacity of 100 ml and 200 ml closed with a chlorobutyl rubber closure and an aluminium ring seal.

Carton box with 1, 10, 20 or 40 vials of 100 ml or 1, 10 or 20 vials of 200 ml.

Not all pack sizes may be marketed.

For any information about this veterinary medicinal product, please contact the local representative of the marketing authorisation holder.

For animal treatment only - To be supplied only on veterinary prescription.

Controlled Medicines/Ubat Terkawal
Phenol as preservative

16. Pharmacodynamic properties

Injectable iron-carbohydrate complexes are established haematinic agents in veterinary medicine. Following intramuscular injection, the complex is absorbed and metabolized to release the iron for utilization and/or storage in accordance with the nutritional status of the animal. In iron deficient states, the iron is utilized for the synthesis of haemoglobin and other iron-containing molecules. Excess iron is stored principally in the liver.

17. Pharmacokinetic particulars

Absorption of the product has been shown to be rapid. Over 95% of the administered iron (1mL/200 mg iron administered at three days of age) was absorbed by 24 hours after injection



Producto/Product	Gleptoferron Labiana	Colores	Control de calidad
Dosis/Strength	100 y 200 ml	 P Black	Fecha
Envase/Pack Size	Prospecto	 Troquel	Registros
Material/Material			Fecha
Dimensiones/Dimensions	126 x 210 mm		Márqueting
Laetus/Laetus			Fecha
Código/Item Code	71204/001		
Fecha edición/Date of Issue			
Tipografía/Fonts	Segoe UI / 8 pt	OBSERVACIÓN	
Programa/Software	InDesign CC	Tonos de color no válidos para impresión	
Cliente		Proveedor	
			