

Lincofarm TR 44% w/w Soluble Powder

Composition:

Contains per 100 g:

Lincomycin Hydrochloride.....44,000 mg

Butylparaben*30 mg

*Antimicrobial preservative

Product description:

White to almost white powder.

Pharmacodynamics:

Lincomycin is a lincosamide antibiotic derived from *Streptomyces lincolnensis* which inhibits protein synthesis. Lincomycin binds to the 50S sub-unit of the bacterial ribosome close to the peptidyl transfer centre and interferes with the peptide chain elongation process by causing premature peptidyl-tRNA dissociation from the ribosome.

Lincomycin is active against some gram-negative bacteria (*Clostridium perfringens*) and mycoplasmas (*Mycoplasma hyopneumoniae*).

While the lincosamides are generally considered to be bacteriostatic agents, the activity depends on the sensitivity of the organism and concentration of the antibiotic. Lincomycin may be either bactericidal or bacteriostatic.

Resistance to lincomycin is frequently conferred by plasmid-borne factors (erm genes) coding for methylases modifying the ribosomal binding site and frequently leading to cross-resistance to other antimicrobials of the macrolides, lincosamides and streptogramins group. However, the most prevalent mechanism in mycoplasmas is the alteration of the binding site through mutational events (chromosomal resistance). Lincomycin resistance mediated by efflux pumps, or by inactivating enzymes, has also been described. There is often complete cross-resistance between lincomycin and clindamycin.

Pharmacokinetics:

In pigs, lincomycin is rapidly absorbed following oral administration.

Crossing the intestinal barrier, lincomycin is widely distributed to all tissues, especially the lungs and joint cavities; the volume of distribution is about 1 litre.

Approximately 50% of lincomycin is metabolised in the liver. Lincomycin undergoes enterohepatic circulation.

Lincomycin is eliminated unchanged or in the form of various metabolites in bile and urine. High concentrations of the active form are observed in the intestine.

Chickens were administered lincomycin hydrochloride in the drinking water at a level of approximately 34 mg/litre (5.1-6.6 mg/kg body weight) for seven days.

Metabolites comprised more than 75% of total residues in the liver. Lincomycin and one unknown metabolite comprised >50% of the muscle residue at zero hours.

The excreta contained mostly unmetabolised lincomycin (60-85%) during treatment.

Indication:

Pigs

Treatment and metaphylaxis of enzootic pneumonia caused by *Mycoplasma hyopneumoniae*.

The presence of the disease in the group must be established before the product is used.

Chickens

Treatment and metaphylaxis of necrotic enteritis caused by *Clostridium perfringens*. The presence of the disease in the group must be established before the product is used.

Recommended dose:

Pigs:

Enzootic pneumonia: 10 mg lincomycin per kg of body weight (corresponding to 22.73 mg product per kg bodyweight) for 21 consecutive days.

Chickens:

Necrotic enteritis: 5 mg lincomycin per kg of body weight (corresponding to 11.36 mg product per kg bodyweight) for 7 consecutive days. The concentration to be used depends on the actual body weight and the water consumption of the animals and can be calculated according to the following formula:

$$\frac{\text{Dosage (mg product per kg body weight per day)}}{\text{Average daily water intake (litre/animal)}} \times \frac{\text{Mean body weight (kg) of animals to be treated}}{\text{litre}} = \frac{\text{mg product per litre drinking water}}$$

The use of suitably calibrated weighing equipment is recommended if part packs are used. The daily amount is to be added to the drinking water in such a way that all medication will be consumed within 24 hours. Medicated drinking water should be freshly prepared every 24 hours. No other source of drinking water should be available.

Controlled Medicine

Keep out of reach of children

Jauhi dari kanak-kanak

Instructions of use:

For use in drinking water.

Dosing guidance and recommended doses:

To ensure a correct dosage, body weight should be determined as accurately as possible to avoid underdosing.

The intake of medicated water depends on the physiological and clinical condition of the animals. In order to obtain the correct dosage, the concentration of the lincomycin has to be adjusted accordingly.

The uptake of water should be monitored frequently.

The medicated water should be the only source of drinking water for the animals for the entire duration of the treatment period.

After the end of the medication period, the water supply system should be cleaned appropriately to avoid intake of sub-therapeutic amounts of the active substance.

Contraindications:

Do not use in cases of known hypersensitivity to the active substance or to any of the excipients.

Do not administer, and do not allow access to water containing lincomycin, to rabbits, hamsters, guinea pigs, chinchillas, horses or ruminants as this could result in severe gastro-intestinal disturbance.

Do not use in cases of known resistance to lincosamides.

Do not use in cases of hepatic dysfunction.

Warning and Precautions:Special warnings for each target species

Medicated drinking water uptake can be affected by the severity of the disease. In case of insufficient uptake of water, pigs should be treated parenterally.

The susceptibility of *Mycoplasma hyopneumoniae* to antimicrobial agents is difficult to test in vitro owing to technical constraints. In addition, there is a lack of clinical breakpoints for both *M. hyopneumoniae* and *C. perfringens*. Where possible, therapy should be based on local (regional, farm level) epidemiological information concerning the response of enzootic pneumonia/necrotic enteritis to treatment with lincomycin.

Special precautions for use

Special precautions for use in animals

Use of the veterinary medicinal product preferably should be based on identification of the target pathogen and susceptibility testing of the bacteria isolated from the animal. See also the above special warnings for each target species.

Official, national and regional antimicrobial policies should be taken into account when the veterinary medicinal product is used.

Use of the veterinary medicinal product deviating from the instructions given in the summary of product characteristics may increase the prevalence of bacteria resistant to the lincomycin and may decrease the effectiveness of treatment with other lincosamides, macrolides and streptogramin B due to the potential for cross-resistance.

Repeated or prolonged use should be avoided by improving the farm management and hygiene practices.

Special precautions to be taken by the person administering the veterinary medicinal product to animals.

This product contains lincomycin and lactose monohydrate, either of which can cause allergic reactions in some people. People with known hypersensitivity to lincomycin or any other lincosamide, or to lactose monohydrate, should avoid contact with the veterinary medicinal product.

Care should be taken not to raise and inhale any dust.

Contact with skin and eyes should be avoided. Personal protective equipment consisting of approved dust masks, gloves and safety glasses should be worn when handling and mixing the product. If respiratory symptoms develop following exposure, seek medical advice and show this warning to the physician.

In case of accidental exposure to the skin, eyes or mucous membranes, wash the affected area thoroughly with plenty of water. If symptoms such as skin rash or persistent eye irritation appear after exposure, seek medical advice immediately and show the package leaflet or label to the physician.

Wash hands and any exposed skin with soap and water immediately after use.

Do not eat, drink or smoke while handling the product.

Interactions with other medicaments:

Antagonism may exist between lincomycin and macrolides such as erythromycin and other bactericidal antibiotics; concurrent use is therefore not recommended due to competitive binding at the 50S ribosomal subunit of the bacterial cell.

The bioavailability of lincomycin may decrease in the presence of gastric antacids or activated charcoal, pectin or kaolin. Lincomycin can potentiate neuromuscular effects of anaesthetics and muscle relaxants.

Use during pregnancy, lactation or lay

Laboratory studies in rats have not produced any evidence of teratogenic effects, although foetotoxicity has been reported. The safety of the veterinary medicinal product has not been established during pregnancy, lactation or lay in the target species. Use according to the benefit-risk assessment by the responsible veterinarian.

Side Effects:

On rare occasions, pigs given lincomycin- medicated water may develop diarrhoea/soft stools and/or mild swelling of the anus within the first 2 days after onset of treatment. On rare occasions some pigs may show reddening of the skin and mild irritable behaviour. These conditions are usually self-correcting within 5-8 days without discontinuing the lincomycin treatment. Allergic/hypersensitive reactions occur on rare occasions.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reactions)
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

Symptoms and treatment of overdose:

A dosage greater than 10 mg lincomycin per kg of body weight may cause diarrhoea and loose stools in pigs.

In case of accidental overdose, the treatment must be stopped and restarted at the recommended dose level.

There is no specific antidote, treatment is symptomatic.

Withdrawal period:

Pigs:

Meat and offal: 1 day.

Chickens:

Meat and offal: 5 days.

Not authorised for use in laying birds producing eggs for human consumption.

Storage condition:

Keep the product in its original packaging at room temperatures below 30 C, away from sunlight.

Shelf-life as packaged for sale: 3 years

Shelf-life after first opening the 200 g sachet: 24 hours.

Shelf-life after first opening the 25 kg bag: 30 days

Shelf-life after dilution: 24 hours.

Packaging:

- 200 g Sachet
- 25 kg Bag

Manufacturer:

FARMABASE SAÚDE ANIMAL LTDA.

Praça Emílio Marconato, No. 1000. Galpões A1 to A6
Chácaras Primavera, Núcleo Residencial Doutor João
Aldo Nassif,

CEP: 13916-074, Jaguariúna/SP – Brazil

Phone: +55 (19) 3837-9900

CNPJ: 073.174.377/0001-30

Brazilian industry

Marketing Authorization Holder:

F.E Venture Sdn. Bhd., Lot 3 & 5, Jalan PJS 11/8,
Bandar Sunway, 46150, Petaling Jaya,
Selangor, Malaysia.

MAL registration number:

Date of revision: October 13th 2021

