



Cattle	MIC _{90s}	MBC _{90s}
	µg/ml	
<i>Mannheimia haemolytica</i>	0.5	1
<i>Pasteurella multocida</i>	1	2
<i>Histophilus somni</i>	1	2
Pigs	MIC _{90s}	MBC _{90s}
	µg/ml	
<i>Actinobacillus pleuropneumoniae</i>	4	4
<i>Pasteurella multocida</i>	1	2
<i>Haemophilus parasuis</i>	0.5	0.5
<i>Bordetella bronchiseptica</i>	2	4
Sheep	MIC ₉₀	
	µg/ml	
<i>Fusobacterium necrophorum</i>	MIC ₉₀ : 64	
<i>Dichelobacter nodosus</i>	0.008 – 0.016	

SHELF LIFE AS PACKAGED FOR SALE
3 years

SHELF LIFE AFTER FIRST OPENING OF CONTAINER
Once open use within 28 days

STORAGE CONDITION
Store in a cool and dry place below 30°C.

OTHER PACKAGING AVAILABLE
Cardboard box containing 1 vial of 50, 100, or 250ml.
Not all pack sizes may be marketed.

ENVIRONMENTAL PROPERTIES / DISPOSAL
Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements. Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

DATE OF REVISION OF PACKAGE INSERT
03/ 2022

Zactran® is a registered trademark of Boehringer Ingelheim Animal Health France SCS, used under licence.

Three mechanisms are generally considered responsible for resistance to the macrolide class of compounds. This is often referred to as MLSB resistance as it affects macrolides, lincosamides and streptogramins. The mechanisms involve the alteration of the ribosomal target site, the utilization of active efflux mechanism and the production of inactivating enzymes.

PHARMACOKINETICS

Cattle :

Gamithromycin administered subcutaneously into the neck of cattle at a single dose of 6 mg/kg body weight, resulted in rapid absorption with peak plasma concentrations observed after 30 to 60 min with a long plasma half-life (> 2 days). The bioavailability of the compound was > 98 % with no gender differences. The volume of distribution at steady-state was 25 l/kg. Gamithromycin levels in lung reached a maximum in less than 24 hr, with lung-to-plasma ratio of > 264 indicating that gamithromycin was absorbed rapidly into the target tissue for BRD. In vitro plasma protein binding studies determined that the mean concentration of the free active substance was 74 %. Biliary excretion of the unchanged drug substance was the major route of elimination.

Pigs :

Gamithromycin administered intramuscularly in pigs at single dose of 6 mg/kg body weight, resulted in rapid absorption with peak plasma concentrations observed after 5 to 15 min, with a long plasma half-life (about 4 days). The bioavailability of gamithromycin was > 92 %. The compound is absorbed rapidly into the target tissue for SRD. Accumulation of gamithromycin in the lung has been demonstrated by high and sustained concentrations in the lung and bronchial fluid which far exceed those in blood plasma. The volume of distribution at steady-state was approximately 39 l/kg. In vitro plasma protein binding studies determined that the mean concentration of the free active drug was 77 %. Biliary excretion of the unchanged drug was the major route of elimination.

Sheep :

Gamithromycin administered subcutaneously into the neck of sheep at a single dose of 6 mg/kg body weight is rapidly absorbed, and maximum plasma concentrations were observed between 15 minutes and 6 hours after dosing (2.30 hours on average) with high absolute bioavailability of 89 %. Gamithromycin skin concentrations were much higher than the plasma concentrations resulting in skin/plasma concentration ratios of approximately 21, 58, and 138 at two, five and ten days post-dosing, respectively, demonstrating extensive distribution and accumulation in skin tissue.

OIC CENTRALISÉ
MODIFICATION

- Design (couleur, photo, pictogramme)
- Position et taille des zones de textes (incluant blue box)
- Police/corps de caractère
- Taille de l'OIC
- Autres....

PACK OPS LYON GRAPHIC ARTWORK APPROVAL

MYZACTRAN/leaflet/008/Apr2021	VERSION	PTG: 0137 Version : B 230119	COLORS
PRODUCT: NOT ZACTRAN VERSO	INITIALS: NSe - Composer RH		Noir/Black
COUNTRY: MY	VERSION - DATE:		
Dimensions à plat (mm): 160 x 360	A 14/04/20 11h00	E 05/07/2021	
Dimensions plié(e)s: 160 x 30	B 28/04/20 15h00	F 05/08/2021	
PRINTED BASIS CODE (if applicable):	C 22/05/20 15h00	G 03/03/2022	
NA	D 01/06/2021		
	CANCEL AND REPLACE:		
	MYZACTRAN/leaflet/007/Apr2021		

! Please use the official PANTONE® (P) matching system for an accurate color representation.

J110320

Marketing approval:	Regulatory approval:
COMPLETE NAME:	COMPLETE NAME:
DATE + SIGNATURE:	DATE + SIGNATURE:

NOT PRINTABLE NON IMPRIMABLE	TECHNICAL NOTE(S) - INFO(S) TECHNIQUE
	Artwork identification(s)
	Text areas / zones de textes
	Dieline / Découpe