

Veterinary Package Insert

TP-GENTA 300 KIU/G WATER SOLUBLE POWDER

PRODUCT DESCRIPTION

White to off-white colour water soluble powder, which contains 300,000IU of gentamycin sulphate per gram in sachet.

PHARMACODYNAMICS

Gentamicin is a bactericidal aminoglycoside antibacterial, whose antimicrobial action results from acting on the 30 S subunit of the ribosomes, preventing protein synthesis. Prevents the entire initiation phase, disrupting the ordering of messenger RNA and causing an incorrect reading of the genetic code by the transfer RNA. It also disrupts the permeability of the bacterial membrane.

PHARMACOKINETICS

Gentamicin is barely absorbed orally and consequently plasma concentrations are practically undetectable. It remains in the gastrointestinal tract and its distribution to the rest of the organs and tissues is red.

INDICATION

Pig:

In weaned animals.

Treatment of digestive infections caused by non-invasive *Escherichia coli* strains sensitive to gentamicin.

Treatment of swine dysentery caused by microorganisms sensitive to gentamicin.

TARGETED SPECIES

Pigs.

RECOMMENDED DOSAGE AND ADMINISTRATION

The product is recommended for oral administration in drinking water.

Administer 2,000 IU of gentamicin sulfate / kg live weight (equivalent to 66.7 mg of medicine per 10 kg of live weight) for 3 consecutive days.

The weight of the animals should be determined as accurately as possible to avoid insufficient dosing.

Water consumption depends on the clinical situation of the animal and the time of the year. To ensure a dosage correct, the concentration of gentamicin in the water will be adjusted taking into account daily consumption.

Depending on the recommended dose, the number and weight of the animals that should be treated, the exact daily dose of the medication by applying the following formula:

g of medicine / liter of drinking water	=	Dose of active substance (IU/kg live weight/day)
		x average weight of animals (kg)

		IU of active substance in g of medicine (IU/g)
		x Average consumption of
		water (liters/day)

*Note: 300,000 IU of gentamicin sulfate is equivalent to 473.9mg of gentamicin sulfate, when 1mg of gentamicin sulfate is equivalent to 633 IU.

CONTRAINDICATIONS

Do not use in case of hypersensitivity to aminoglycosides or to any of the excipients.

WARNINGS & PRECAUTIONS.

Special precautions for use in animals

Good clinical practice requires basing treatment on sensitivity tests of bacteria isolated from sick animals. If this is not possible, treatment should be based on local epidemiological information. (regional, at the exploitation level) on the sensitivity of different strains of bacterial species usually involved in the infectious process.

The use of the drug in conditions other than those recommended in the Technical Data Sheet may increase the prevalence of gentamicin resistant bacteria and decrease the effectiveness of aminoglycoside treatment such as consequence of the appearance of cross resistance.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

People with known hypersensitivity to gentamicin should avoid contact with the medication.

Handle the medication with caution. During handling, wear consistent personal protective equipment in gloves, work overalls, safety glasses and protective mask.

Avoid direct contact with the skin, mucosa and inhalation of the product. In case of accidental contact, wash immediately with plenty of water. If symptoms appear after exposure, such as a rash, see with a doctor and show him the leaflet or label.

Do not eat, drink or smoke while handling the product.

Wash hands after use.

INTERACTIONS WITH OTHER MEDICATIONS

Do not administer together with other aminoglycosides or bacteriostatic antibiotics.

Revised November 2021

PREGNANCY AND LACTATION

Its use is not recommended during pregnancy or lactation.

ADVERSE EFFECTS

Intestinal dysbiosis may occur.

OVERDOSE & TREATMENT

No information.

WITHDRAWAL PERIOD

Meat: 14 days

MAXIMUM RESIDUAL LIMIT (MRL)

Substance/ Drug Definition of residues in which MRL was set	Food	Maximum Residue Limits (MRLs) in food µg/kg
Gentamycin	Muscle & fat (porcine)	100
	Liver (porcine)	200
	Kidney (porcine)	1000

STORAGE CONDITION

Store below 30°C. Protect from direct sunlight.

SHELF LIFE

2 years.

After 1st opening, reconstitution or dilution, use within 24 hours.

PACKING

200g

500g

1000g

THYE PHARMA SDN. BHD.

(manufacturer & product registration holder)

No.2 Lorong Industri Ringan Permatang

Tinggi 8,

Kawasan Industri Ringan Permatang Tinggi,

14000 Bukit Mertajam,

Pulau Pinang.