

PF MONENSIN PREMIX 20% W/W GRANULE

Composition:

Monensin (as Monensin Sodium).....200mg

Exp. Qs.....1g

Description

Grey or taupe granule

Pharmacodynamic

Monensin is an ionophore antibiotic derived from *Streptomyces* that forms complexes with monovalent cations, including sodium and potassium. The complexes are transported in a nonpolar manner across the bacterial cell membrane. As such, it acts as a Na⁺/H⁺ antiporter. Monensin blocks intracellular protein transport, resulting in antibacterial and antimalarial effects. Monensin also increases the production of propionic acid and thus prevents bloat.

Pharmacokinetic

Ionophores are absorbed orally. They are rapidly and extensively metabolized by the liver and the numerous metabolites are excreted by bile and eliminated in the feces

Target Species

Broiler Chickens

Indication

As an aid in the prevention of coccidiosis caused by *Eimeria acervulina*, *E. mitis*, *E. necatrix*, *E. maxima*, *E. tenella*, and *E. brunetti* in broiler chickens.

Route of Administration

For oral administration

Recommended dose

Important - must be thoroughly mixed in feeds before use.

Broiler chickens:

Thoroughly mix 500 g (0.50 kg) of PF MONENSIN PREMIX 20% W/W GRANULE in one metric tonne (1000 kg) of feed to provide monensin at 100 g monensin activity per metric tonne of feed. It is recommended that MONENSIN PREMIX 20% W/W GRANULE be mixed with a small quantity of one of the feed ingredients before it is incorporated in the finished feed.

Feed the medicated feed continuously as the sole ration.

Contraindications

Do not use MONENSIN premix for treatment of outbreaks of coccidiosis.

Do not allow canines, horses or other equines, or guinea fowl access to formulations containing monensin. Ingestion of monensin by these species has been fatal.

Interaction with other medicinal products

Poultry consuming monensin should not be treated with products containing tiamulin. Severe growth depression may occur.

Pregnancy and lactation

Do not feed to replacement and laying chickens

Side effects

None under correct use of the product

Symptoms and treatment of overdose

In case of overdosage, possible suppression of growth and intoxication (expressed of loss of appetite, tremor, paresis of limbs and death) could occur.

Storage

Store in a cool dry place (<30°C). Strictly avoid light and heat exposure.

Warning and Precaution

WARNINGS:

1. Do not feed to replacement and laying chickens
2. When mixing and handling MONENSIN premix, use protective clothing, impervious gloves and a dust mask. Operators should wash thoroughly with soap and water after handling.
3. No pre-slaughter withdrawal is required when this drug product is used according to the label directions.

CAUTIONS:

1. Do not use MONENSIN premix for treatment of outbreaks of coccidiosis.
2. Do not exceed recommended levels as reduced average daily gains may result.
3. Do not allow canines, horses or other equines, or guinea fowl access to formulations containing monensin. Ingestion of monensin by these species has been fatal.
4. May be used in feeds containing the pellet binding agents bentonite (2%), Attapulgit (2%), Kaolin (2.5%), Lignin Sulfonate (4%), Carboxy Methylcellulose (0.1%) or Agri-Colloid.
5. Poultry consuming monensin should not be treated with products containing tiamulin. Severe growth depression may occur.
6. NOT TO BE USED AFTER EXPIRATION DATE.
- Jauhkan ubat dari kanak-kanak/Keep out of reach of children

Shelf life

24 months

Shelf life after opening

24 hours

Shelf life after reconstitution or dilution

24 hours

Withdrawal period

No preslaughter withdrawal time is required when this drug product is used according to the label directions

Packaging

25kg

Revised date

24/08/2021

Product Registration Holder:

PANFAST MARKETING (M) SDN.BHD.

PENANG SCIENCE PARK 1102, MUKIM 13, LORONG PERINDUSTRIAN BUKIT
MINYAK 18, KAWASAN PERINDUSTRIAN BUKIT MINYAK,
14100 SIMPANG AMPAT, PENANG, MALAYSIA
TEL: +604-5016868 FAX: +604-5016866

Manufactured by:

SHANDONG QILU KING-PHAR PHARMACEUTICAL
CO., LTD.

21 QINGLONG ROAD, PINGYIN, JINAN, 250400, CHINA
Tel: +86 531 83127771 Fax : +86 531 83127774