

URISOL

(Sodium Bicarbonate, Tartaric acid, Citric acid and Sodium Citrate Effervescent Granules Sachet)

COMPOSITION:

Each sachet contains:

Sodium Bicarbonate BP	1.76g
Tartaric Acid BP	890mg
Citric Acid Anhydrous BP	720mg
Sodium Citrate Anhydrous USP	630mg

PRESENTATION:

28 Sachets in a Carton

PRODUCT DESCRIPTION:

Granules: White to off white coloured, lemon flavoured granules.

Reconstituted Solution: The reconstituted solution is clear and lemon flavored.

CLINICAL PHARMACOLOGY:

PHARMACODYNAMICS

ATC code: G04BX Other urologicals; Used in the treatment of urological problems

Sodium Bicarbonate, Tartaric Acid, Citric Acid and Sodium Citrate Effervescent granules have gastric antacid as well as urinary alkalinising properties.

Sodium Bicarbonate increases the alkali reserve of the plasma and increases excretion of urine, which is rendered less acidic. Sodium Citrate is used to make the urine alkaline in the treatment of urinary tract infections. Citric acid increases the secretion of urine and renders it less acidic. It is also used in the preparation of effervescent granules to aid effervescence.

PHARMACOKINETICS AND DRUG METABOLISM

Sodium Citrate and Citric Acid: Sodium citrate is metabolized to bicarbonates which increases urinary pH by increasing the excretion of free bicarbonate ions without producing systemic alkalosis when administered in recommended doses.

Sodium bicarbonate is excreted through renal and also via lungs by forming CO₂.

Sodium citrate, citric acid and absorbed tartaric acid are excreted through the urine.

INDICATIONS AND USAGE:

Indicated for relieving of discomfort in mild urinary tract infection; symptomatic relief of dysuria; to enhance the action of certain antibiotics especially some sulphonamides; in gout as urinary alkalinisers to prevent crystallisation of urates and dissolution of uric acid stones.

DOSAGE AND ADMINISTRATION:***Adults:***

Dissolve the contents of 1 – 2 sachets in a glass of cold water. It may be taken 4 times daily for 5 days unless advised to do so by the physician.

Children:***Children under 12 years old***

Do not give unless advised to do so by physician.

Method of Administration

Dissolve the contents of 1 – 2 sachets in a glass of cold water. It may be taken 4 times daily for 5 days unless advised to do so by the physician.

SIDE EFFECTS:

The tartrate component of Sodium Bicarbonate, Tartaric Acid, Citric Acid and Sodium Citrate Effervescent Granules Sachet may be incompletely absorbed. Because of this, Sodium Bicarbonate, Tartaric Acid, Citric Acid and Sodium Citrate Effervescent Granules may exert a mild laxative effect. Prolonged and excessive use may cause systemic alkalosis and/or hypernatremia.

Sodium bicarbonate may cause flatulence. Mild diuresis may occur.

INTERACTIONS WITH OTHER MEDICINAL PRODUCTS AND OTHER FORM OF INTERACTION:

General: Alkalisiation of the urine due to the use of Sodium Bicarbonate, Tartaric Acid, Citric Acid and Sodium Citrate Effervescent Granules Sachet, theoretically, may result in a decreased therapeutic effect of the following medications: chlorpropamide, lithium, salicylates and tetracyclines.

Alternatively, alkalisiation of the urine due to the use of Sodium Bicarbonate, Tartaric Acid, Citric Acid and Sodium Citrate Effervescent Granules Sachet, theoretically, may result in an increased therapeutic effect of the following medications: amphetamines, ephedrine/pseudoephedrine.

Antacids:

Concomitant use of Sodium Bicarbonate, Tartaric Acid, Citric Acid and Sodium Citrate Effervescent Granules Sachet with an antacid by patients with compromised renal function may result in the absorption of dangerously high amounts of aluminium.

Co-administration of antacids with citrates, such as contained in Sodium Bicarbonate, Tartaric Acid, Citric Acid and Sodium Citrate Effervescent Granules Sachet, may result in systemic alkalosis, while concurrent administration of antacids with sodium citrate and sodium bicarbonate may lead to the development of calcium stones in patients with uric acid stones. It may also cause hypernatraemia.

Quinolones:

The sodium citrate in Sodium Bicarbonate, Tartaric Acid, Citric Acid and Sodium Citrate Effervescent Granules Sachet may reduce the solubility of quinolone antibiotics, such as ciprofloxacin, norfloxacin, or ofloxacin, in the urine. Patients should therefore be monitored for signs of crystalluria and nephrotoxicity.

Salicylates:

Co-administration of salicylates with the sodium citrate in Sodium Bicarbonate, Tartaric Acid, Citric Acid and Sodium Citrate Effervescent Granules Sachet may increase urinary excretion and reduce therapeutic effects of salicylates as a result of urinary alkalinisation.

Tetracyclines:

Due to an increase in intragastric pH, the absorption of tetracyclines may be decreased when it is co-administered with sodium bicarbonate. Sodium Bicarbonate, Tartaric Acid, Citric Acid and Sodium Citrate Effervescent Granules Sachet should therefore not be taken within 1 to 2 hours of tetracycline administration.

Ketoconazole:

Since the sodium bicarbonate in Sodium Bicarbonate, Tartaric Acid, Citric Acid and Sodium Citrate Effervescent Granules Sachet may increase gastrointestinal pH, the absorption of ketoconazole may be markedly reduced with concurrent administration. Patients should wait at least 2 hours before taking Sodium Bicarbonate, Tartaric Acid, Citric Acid and Sodium Citrate Effervescent Granules Sachet following ketoconazole administration.

Methenamine:

Sodium Bicarbonate, Tartaric Acid, Citric Acid and Sodium Citrate Effervescent Granules Sachet should not be administered with urinary tract antiseptics which require acidic urine, such as methenamine mandelate and methenamine hippurate.

Laxatives:

Concurrent administration of citrates with laxatives may have an additive effect.

SPECIAL WARNING AND PRECAUTION FOR USE:

Do not give this medicine to children under 12 years of age unless advised to do so by a doctor. Patients should be advised to consult your doctor or pharmacist before using this medicine if you are taking other medicines or if you have kidney problems, heart problems or high blood pressures.

Patients should be advised to consult a doctor if pain or irritation persists for more than 48 hours or if you notice blood in your urine.

This preparation contains 650mg of sodium per sachet and should be taken into account if on a low sodium diet. Also contains soya bean products and sugars.

The clinical condition of the patient should be evaluated and laboratory determinations (eg. serum electrolytes, acid-base balance) obtained periodically during Sodium Bicarbonate, Tartaric Acid, Citric Acid and Sodium Citrate Effervescent Granules Sachet therapy, particularly in patients with renal disease.

PREGNANCY AND LACTATION:

Pregnancy:

Studies regarding the effect of citrates on pregnancy have not been done.

Nursing Mothers:

Caution should be exercised when administered to a nursing mother.

OVERDOSAGE:

Treatment is symptomatic and supportive and consists mainly of correction of fluid and electrolyte balance. Consult a doctor in known cases of overdosage.

CONTRAINDICATION:

Renal failure or hypernatraemia; in conjunction with hexamine mandelate or hexamine hippurate therapy because an acid urine is needed.

Caution is advised in overt and occult cardiac failure.

Concomitant use of urinary alkalinisers and quinolone antibiotics should be avoided; crystalluria may be more likely to occur with alkaline urine.

STORAGE CONDITION:

Store below 30°C. Store in the original package in order to protect from moisture.

After dissolving the sachet as recommended for dosing, it is to be used whole at a time and do not store for later use.

KEEP ALL MEDICINES OUT OF REACH AND SIGHT OF CHILDREN

MANUFACTURED BY:



Ind-Swift Ind-Swift Limited

Off. NH-21, Village Jawaharpur,
Tehsil Derabassi,
District SAS Nagar (Mohali)
Punjab-140507, India.

PRODUCT REGISTRATION HOLDER

Healol Pharmaceuticals Sdn Bhd,

74-3, Jalan Wangsa Delima 6, KLSC Wangsa Maju, 53300, Kuala Lumpur, Malaysia

DATE OF REVISION

November 2021