

VETRICOX 5MG/ML SOLUTION

Description and Composition

VETRICOX 5MG/ML SOLUTION is a clear, light yellow to yellow brown coloured solution. Each ml contains 5mg Diclazuril as its active ingredient.

Pharmacodynamics

Diclazuril is an anticoccidial of the benzene acetonitrile group and has an anticoccidial activity against *Eimeria* species. Depending on the coccidia species, diclazuril has a coccidiocidal effect on the asexual or sexual stages of the development cycle of the parasite. Treatment with diclazuril causes interruption of the coccidial cycle and of excretion of oocysts for approximately 2 to 3 weeks after administration.

Pharmacokinetics

Biotransformation of diclazuril in chicken is low and the major compound found in plasma, tissues, urine and faeces is the parent substance. It was also concluded that although poorly absorbed, diclazuril is detectable in plasma soon after administration with a t_{max} of about 6 hours.

Indication

For the treatment of Coccidiosis caused by *E. tenella*, *E. maxima* and *E. acervulina* in broiler chickens.

Recommended Dose

For oral administration.

Dilute 0.25ml to 0.5ml (0.05ml to 0.1ml per kg body weight per day) of Vetricox 5mg/ml Solution per 1L of drinking water and administered for 2 consecutive days. This is equivalent to 0.25mg to 0.5mg Diclazuril per kg body weight per day.

Route of Administration

For oral administration.

Contraindication

None known.

Warning and Precautions

Special warnings for each target species:

None known.

Special precautions for use

Special precautions for use in animals:

- Do not use on animals that have been in shock or experience hypersensitivity reaction to this drug.
- Use as directed by your veterinarian.
- Read the instruction manual (precautions) thoroughly before use.
- Do not use arbitrarily because safety and efficacy have not been established other than with the specified breed.
- In case of non-compliance with the withdrawal period, it may remain in livestock products such as meat, so be sure to monitor the time accurately.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

- The operator should wear protective equipment such as gloves, a mask and protective clothing to avoid direct contact with skin or inhalation.
- In case of contact with skin or eyes while handling the drug, immediately rinse thoroughly and seek medical attention as soon as any abnormal symptoms are found.

Interaction with Other Medicaments

None known.

Pregnancy and Lactation

No residue data in eggs were provided; therefore, the use should be restricted to animals from which eggs are not produced for human consumption.

Side Effects

None known.

Symptoms and Treatment of Overdose

Chickens for fattening tolerated a 25-fold overdose of diclazuril for 37 days without any observable negative effects on performance, haematology, necropsy and histopathology.

Storage Condition

Store below 30°C.

Shelf life

3 years.

To be used within 24 hours after dilution in drinking water.

After first opening the container, any product remaining after 24 hours should be discarded.

Withdrawal Period

Chickens: 5 days

Packing

100ml, 200ml, 500ml, 1L.

Revised date: 20.12.2021

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com