



Package Insert

1. Name of the medicinal product

MediThroat Spray 3.0 mg/ml

2. Qualitative and quantitative composition

The active substance is Benzydamine HCl 0.3% w/v

3. Pharmaceutical form

Throat Spray that is a clear, colourless solution and smells of strawberry.

4. Clinical particulars

4.1 Therapeutic indications

Temporary relief of painful conditions of the mouth and throat including tonsillitis, sore throat, radiation mucositis, aphthous ulcers, pharyngitis, swelling, redness, inflammatory conditions, post orosurgical and periodontal procedures.

4.2 Posology and method of administration

Route of administration: Oromucosal.

Adults and children over 12 years: 2-4 sprays (1-2 mg) directly onto the sore/inflamed area and swallow gently. Repeat every 1½ to 3 hours as necessary.

Children (6-12 years): 2 sprays (1 mg) directly onto sore/inflamed area and swallow gently. Repeat every 1½ to 3 hours as necessary.

Children under 6 years: Not recommended.

Uninterrupted treatment should not exceed seven days, unless under medical supervision.

Instruction for Use:

- (1) Hold the bottle in an upright position.
- (2) Prime the spray the first time using it (max. 5 sprays). Point the spray-tube away from the face. Press down firmly until a fine spray appears from the end of the spray-tube.
- (3) Aim the spray-tube at the affected area and press down. One press releases one actuation.
- (4) After finishing the desired number of puffs, wipe the end of the spray-tube with a clean tissue.

This will prevent the spray-tube from getting blocked.

- (5) Do not squeeze anything into the spray-tube if it becomes blocked.



4.3 Contraindications

Patients with known hypersensitivity to benzydamine or to the other components listed in section 6.

4.4 Special warnings and precautions for use

If a sore throat is either caused or complicated by a bacterial infection, appropriate antibacterial therapy should be considered in addition to the use of MediThroat Spray.

4.5 Interaction with other medicinal products and other forms of interaction

There are no known drug interactions with benzydamine.

4.6 Use in Pregnancy

The safety of benzydamine hydrochloride has not been established in pregnant patients. Risk to benefit ratio should be established if MediThroat Spray is to be used in these patients.

4.7 Effects on ability to drive and use machines

Medithroat spray does not affect the ability to drive or to operate the machine.

4.8 Adverse Effects

MediThroat Throat Spray is generally well tolerated and side- effects are minor. The following adverse reactions have been reported after use of benzydamine hydrochloride in solution form:

Local Adverse Reactions:

The most commonly reported reaction is oral numbness. Occasional burning or stinging sensation may occur. Other local adverse effects were less common and included dryness or thirst , tingling , warm feeling in mouth and altered sense of taste.

Systemic Adverse Reactions:

These were uncommon and consisted mainly of nausea, vomiting, retching, gastrointestinal disorders , dizziness , headache and drowsiness.

Hypersensitivity reactions occur very rarely but may be associated with pruritis, rash, urticaria, photodermatitis and occasionally laryngospasm or bronchospasm.

4.9 Overdose

There are no known cases of overdosage with MediThroat Spray. There is no specific antidote for benzydamine. If an excessive quantity is ingested, the treatment should be symptomatic.



5. Pharmacological properties

Pharmacodynamics:

Benzydamine is an anti-inflammatory analgesic agent structurally unrelated to the steroid group. Benzydamine differs chemically from other non-steroidal anti-inflammatory agents in that it is a base rather than an acid. In common with the aspirin-like drugs, benzydamine possesses an antipyretic activity.

Pharmacokinetics:

Benzydamine is well absorbed following oral administration. Following topical administration, benzydamine is well absorbed into the inflamed local mucosa where it exerts anti-inflammatory and local anaesthetic actions. Plasma benzydamine levels following use are low and proportionate to the amount actually ingested.

Benzydamine and its metabolites are excreted largely in the urine. Metabolism is largely by oxidative pathways, although dealkylation can be shown.

Benzydamine has been detected in blood and urine following gargling of solution form. Most of the absorbed dose is eliminated in the first 24 hours. Repeated administration for 7 days does not result in accumulation of benzydamine in plasma.

6. Pharmaceutical particulars

6.1 List of excipients

Povidone (K-30)、Methyl Paraben 0.30%w/v (preservative), Sodium Benzoate 0.02 %w/v(preservative)、Saccharin Sodium, Sodium Bicarbonate、Strawberry Flavor、Polysorbate 20、Ethanol 8.936%w/v、Purified Water.

6.2 Incompatibilities

Not applicable.

6.3 Special precautions for storage

Store in a cool, dry place. Protect from direct sunlight, store below 30°C.

Always keep the bottle upright during use and when stored.

Use within 6 months after opening.

6.4 Nature and contents of container

Pack size: Each bottle contains 15 ml and 34 ml of solution and comes in a carton.

6.5 Special precautions for disposal and other handling

No special requirements.



YUAN CHOU CHEMICAL & PHARMACEUTICAL CO., LTD
No. 272 , Nangang 3rd Rd., Nantou City , Nantou County 54067, Taiwan R.O.C

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. Name and Address of Manufacturer

Yuan Chou Chemical and Pharmaceutical Co. Ltd.

No. 272, Nangang 3rd Rd, Nantou City, Nantou County 54067, Taiwan.

8. Name and Address of Product Registration Holder

World Medicare Supplies Sdn. Bhd.

Pt 5740 (1st floor) Jalan TS 2/1E,

Taman Semarak 2, 71800 Nilai,

Negeri Sembilan.

9. Date of Revision

June 2024

Font size times new roman 13