

CENMICIN 100/50 mg/ml solution for injection

STATEMENT OF THE ACTIVE SUBSTANCES AND OTHER INGREDIENTS

Each ml contains:

Active substances:

Spectinomycin (sulphate tetrahydrate) 100 mg

Lincomycin (hydrochloride) 50 mg

Excipients:

Benzyl alcohol (E1510) (as preservative) 9 mg

Excipients, n.q.

Clear and colourless solution

INDICATIONS

Treatment of infections caused by strains sensitive to the association lincomycin / spectinomycin

Cattle:

Pneumonia caused by *Pasteurella multocida* and *Mycoplasma* spp.

Pigs:

Swine dysentery caused by *Brachyspira hyodysenteriae*.

Enzootic pneumonia caused by *Mycoplasma hyopneumoniae*.

Sheep:

Pneumonia caused by *Pasteurella multocida* and *Mycoplasma* spp.

Footrot caused by *Dichelobacter nodosus* and *Fusobacterium necrophorum*.

CONTRAINDICATIONS

Do not use in animals with known hypersensitivity to lincomycin, spectinomycin or other antibacterial lincosamides or aminocyclitols and/or any of its excipients. Do not use in horses, as lincomycin causes haemorrhagic colitis and diarrhoea with very serious results. Do not use in animals suffering from pre-existing infections of *Monilia* spp.

SIDE EFFECTS

In very rare case, after intramuscular administration, pain and local irritation may occur.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reactions)
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

If you notice any side effects, even those not already listed in this package leaflet or you think that the medicine has not worked, please inform your veterinary surgeon.

TARGET SPECIES

Cattle, sheep and pigs.

DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION

Instructions for Use:

Route of Administration: Intramuscular

Cattle and pigs: 10 mg of lincomycin + 20 mg of spectinomycin per kg bw/day (equivalent to 2 ml of medicine/10 kg liveweight/day), for 3-5 consecutive days.

Sheep:

Footrot: 5 mg of lincomycin + 10 mg of spectinomycin per kg/bw (equivalent to 1 ml of medicine/10 kg liveweight) as a single dose.

Pneumonia: 10 mg of lincomycin + 20 mg of spectinomycin per kg/bw/day (equivalent to 2 ml of medicine/10 kg liveweight/day), for 3-5 consecutive days.

The weight of the animals should be determined as accurately as possible to avoid under-dosing.

ADVICE ON CORRECT ADMINISTRATION

No applicable.

WITHDRAWAL PERIOD

Meat: 21 days.

Milk: Do not administer to lactating animals producing milk for human consumption.

SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Store below 25°C

Do not use this veterinary medicinal product after the expiry date which is stated on the label or carton after EXP.

Shelf-life after first opening the immediate packaging: 28 days at below 25°C.

SPECIAL WARNINGS

Special warnings for each target species: None.

Special precautions for use in animals:

Good clinical practice requires basing the treatment on susceptibility tests on bacteria isolated from the sick animals. If this is not possible, treatment should be based on local epidemiological information (regional, at farm level) on the sensitivity of the different strains of bacterial species commonly involved in the infectious process.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known hypersensitivity to any of the active substances should avoid contact with the medication.

Do not smoke, eat or drink while handling the medication.

Wash hands after use.

Avoid contact with skin and eyes. If this occurs, wash the affected area with plenty of water.

In the case of accidental self-injection, seek medical advice immediately and show the doctor the information leaflet or label.

If symptoms appear following exposure, such as a skin rash, seek medical advice and show the doctor the information leaflet or label. Swelling of the face, lips or eyes, or difficulty in breathing, are more serious signs and require immediate medical attention.

Use during pregnancy, lactation or lay:

Do not administer to pregnant or lactating females.

Interaction with other medicinal products and other forms of interaction:

Do not administer with macrolides or aminoglycosides.

Overdose (symptoms, emergency procedures, antidotes):

In general, on receiving three times the therapeutic dose, the only symptom is local irritation.

SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Medicines should not be disposed of via wastewater or household waste.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local regulation.

OTHER INFORMATION

Pharmacodynamic properties

SPECTINOMYCIN: Spectinomycin is a bacteriostatic antibacterial; some authors include it within the group of aminoglycosides, while others consider it independent of this group, since it has a different chemical structure from these, although its mechanism of action and

bacterial spectrum are quite similar. It works by inhibiting bacterial protein synthesis by attaching itself to the 30S subunit of the ribosome.

It disrupts the ordering of the messenger RNA and causes an incorrect reading of the genetic code by the transfer RNA.

LINCOMYCIN: Antibacterial from the lincosamide group. It presents a mechanism of action and a bacterial spectrum very similar to that of macrolides. It works by inhibiting the synthesis of bacterial proteins by binding to the 50S subunit of the ribosome, preventing the coupling of transfer RNA molecules.

It is primarily bacteriostatic, but at high concentrations it can be bactericidal.

ASSOCIATION: There is a synergistic effect in the association of lincomycin and spectinomycin in the ratio of 1:2, translating into greater efficacy against different pathological processes than that which they have separately.

Action spectrum:

The association of lincomycin and spectinomycin is active against Gram-negative bacteria such as *Pasteurella multocida* and *Actinobacillus pleuropneumoniae* and Gram-positive such as *Staphylococcus aureus* and β -hemolytic *Streptococcus*, as well as mycoplasmas such as *Mycoplasma hyopneumoniae*, *Mycoplasma bovis*, and *Mycoplasma suis*.

Pharmacokinetic particulars

SPECTINOMYCIN:

After intramuscular (IM) administration it is quickly absorbed and reaches maximum blood concentrations within 20 minutes, with detectable concentrations being maintained for 12 hours.

It penetrates little into the tissues and its distribution is fundamentally extracellular. It is excreted by glomerular filtration and approximately 80% can be recovered unchanged in urine during the first 24-48 hours.

LINCOMYCIN:

After IM administration it is well absorbed. It has a plasma peak between 2-4 hours after administration, with therapeutic concentrations persisting for approximately 6-8 hours.

It is widely distributed throughout the body, crossing the placental barrier. The concentration they reach in tissues is several times higher than the serum one when crossing cell walls. It is metabolized in the liver, locating high concentration in the bile; approximately 30% of the dose is excreted in the urine during the first 4 hours and about 14% with feces. It is also excreted in milk.

ASSOCIATION:

The pharmacokinetic parameters of the association are practically unchanged, especially spectinomycin, coinciding with those that each have separately.

INCOMPATIBILITIES:

In the absence of compatibility studies, this medicinal product must not be mixed with other drugs.

Formats:

Box containing 1 vial of 100 ml

Box containing 1 vial of 250 ml

Box containing 1 vial of 500 ml

Box containing 10 vials of 100 ml

Box containing 10 vials of 250 ml

Box containing 12 vials of 500 ml

Not all pack sizes may be marketed.

**For Veterinary use only. To be supplied only on veterinary prescription
To be administered under veterinary control or supervision**

DATE OF REVISION OF LEAFLET

April 2022

MARKETING AUTHORIZATION HOLDER

YENHER AGRO-PRODUCTS Sdn. Bhd.

1628, Jalal IKS Simpang Ampat 1, Taman IKS Simpang Ampat, 14100 Simpang Ampat, Seberang Perai Selatan, Penang, Malaysia

MANUFACTURER

CENAVISA S.L.

Camí Pedra Estela s/nº, Reus,

43205 Tarragona España

Tel 34-977-75 72 73 www.cenavisa.com

REGISTRATION NUMBER IN MALAYSIA

OC cenavisa
LABORATORIOS

Tel 34-977 75 72 73

www.cenavisa.com

E-mail: cenavisa@cenavisa.com