

AMPROFEED 250G/KG POWDER

Description and Composition

AMPROFEED 250G/KG POWDER is an off-white to yellow powder. Each kg contains 250g Amprolium (as hydrochloride) as its active ingredient.

Pharmacodynamics

By mimicking its structure, amprolium competitively inhibits thiamine utilization by the parasite. Prolonged high dosages can cause thiamine deficiency in the host; excessive thiamine in the diet can reduce or reverse the anticoccidial activity of the drug.

Amprolium is thought to act primarily upon the first generation schizont in the cells of the intestinal wall, preventing differentiation of the merozoites. It may suppress the sexual stages and sporulation of the oocysts.

Pharmacokinetics

None known.

Indication

Turkeys: As an aid in the prevention of coccidiosis.

Broiler chickens: As an aid in the prevention of deaths from coccidiosis.

Replacement chickens: As an aid in the prevention of death from coccidiosis in chickens intended as conventional floor managed layers. As an aid in the prevention of deaths from coccidiosis in chickens intended as caged layers.

Calves: As an aid in the prevention and treatment of coccidiosis caused by *Eimeria bovis* and *E. zuernii*.

Recommended Dose

Turkeys:

500g of product per tonne of feed as sole ration from one day of age up to and including 16 weeks of age.

Broiler chickens:

500g or 1kg of product per tonne of feed as sole ration from one day of age until birds are slaughtered.

Replacement chickens:

1) As an aid in the prevention of death from coccidiosis in chickens intended as conventional floor managed layers.

Heavy exposure to coccidiosis: 320g to 500g of product per tonne of feed as sole ration from one day of age until 14 weeks as follows:

Age of chickens	Amount of product per tonne of feed
0-5 weeks	500g
5-9 weeks	320g to 500g
9-14 weeks	320g

Light to moderate exposure to coccidiosis: 240g to 320g of product per tonne of feed as sole ration from one day of age until 14 weeks as follows:

Age of chickens	Amount of product per tonne of feed
0-5 weeks	320g
5-9 weeks	240g to 320g
9-14 weeks	240g

2) As an aid in the prevention of death from coccidiosis in chickens intended as caged layers

500g or 1kg of product per tonne of feed as sole ration from one day of age until birds are placed in cages.

Calves:

Prevention: 250mg of Amprolium per 50kg of body weight (5mg/kg) per day for 21 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

Treatment: 500mg of Amprolium per 50kg of body weight (10mg/kg) per day for 5 days. The proper amount of Amprolium medicated supplement is mixed with the amount of ration usually consumed in one day.

Route of Administration

For oral administration via feed.

Contraindications

Do not administer to debilitated animals.

Warning and Precautions

Special precautions for use:

Special precautions for use in animals:

Chickens and turkeys:

1. Do not use amprolium medicated feed for treatment of outbreaks of coccidiosis.
2. If unexpected deaths occur, obtain an accurate diagnosis and follow veterinarian's or poultry pathologist's recommendations.

Calves:

1. Do not use in feeds for calves intended for future breeding.
2. For a satisfactory diagnosis, a microscopic examination of the feces should be done by a veterinarian or diagnostic laboratory before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

Interaction with Other Medicaments

No drug interactions are reported in animals.

Pregnancy and Lactation

Do not feed to laying birds in production.

Side Effects

Central nervous system (CNS) signs are caused by thiamine deficiency, which may be reversed by adding thiamine to the diet.

Symptoms and Treatment of Overdose

Toxicity is observed only at high doses. Central nervous system (CNS) signs are caused by thiamine deficiency, which may be reversed by adding thiamine to the diet.

Storage Condition

Store below 30°C.

Shelf life

3 years.

To be used within 24 hours after mixing into feed.

Withdrawal Period

Chickens and turkeys: No withdrawal period is required when treated according to the label.

Do not feed to laying birds in production.

Calves: 7 days

Packing

10kg and 25kg.

Revised date: 18.4.22

	Manufacturer & Product Registration Holder Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com
---	---