

Septol Antiseptic Germicide Liquid (Chloroxylenol 5.0% w/v)

Name and Strength of Active Substance(s)

Chloroxylenol 5.0% w/v
Denatured Alcohol 96% 20.83% v/v

Product Description

A light to dark, amber colored solution with pine fragrance.

Pharmacodynamics

Chloroxylenol is a substituted phenol which has been widely used for many years as an ingredient of antiseptic/disinfectant products intended for external use. It is known to be bactericidal in low concentration to a wide range of Gram positive and Gram-negative bacteria.

Pharmacokinetics

Chloroxylenol is well-absorbed when applied to the skin. It is extensively metabolised in the body, probably by the liver, and rapidly excreted, mainly in the urine, as sulphate and glucuronide conjugates. Chloroxylenol has a low systemic toxicity, even at dosage levels many times higher than those likely to be absorbed during normal usage of SEPTOL solution.

Indications

Chloroxylenol 5.0 % w/v (SEPTOL) used for first aid and medical uses such as cuts, bites, abrasions, insect stings and used for personal hygiene conditions such as douching (when medically advised), dandruff, spots and pimples.

Instruction for use

First Aid and Medical:

For cuts, bites, abrasions and insect stings: wash the area with one tablespoonful of SEPTOL diluted in a 250ml of water and cover with dry gauze or lint.

Personal Hygiene:

For douching (when medically advised): Two teaspoonfuls of SEPTOL diluted in 1L of warm water.

For dandruff: One tablespoonful of SEPTOL diluted in 500ml of warm water. Saturate hair and scalp for 10 minutes, then shampoo.

For spots and pimples (not for eczematous condition): Bathe the affected area daily with one tablespoonful of SEPTOL diluted in 250ml of warm water.

SEPTOL Solution can be used as a household disinfectant on surfaces or in laundry.

Household Cleaning:

Laundry: 4 capful of SEPTOL to 2L of water

Floors and other hard surfaces: 5 capful of SEPTOL to 2.4L of water

Lavatories, sinks (non-acrylic), drains etc.: Use undiluted SEPTOL Solution.

Route of Administration

For external use only.

Contraindications

Hypersensitivity to chloroxylenol or to any of the excipients listed in the leaflet.

Warning and Precautions

For external use only. If swallowed, wash out mouth and drink plenty of water or milk. If contact is made with eyes, wash thoroughly with cold water. In both cases consult your doctor. Keep out of the reach of children.

Interactions with Other Medicaments

Not applicable.

Pregnancy & Lactation

No effects during pregnancy are anticipated, since systemic exposure from topical chloroxylenol use is negligible. Chloroxylenol can be used during pregnancy.

It is unknown whether chloroxylenol or its metabolites are excreted in human milk. A risk to the newborns/infants cannot be excluded. Application of the product to the breast is not recommended during breast feeding.

Adverse Effects / Undesirable Effects

Adverse events which have been associated with chloroxylenol are given below, tabulated by system organ class and frequency. Frequencies are defined as: Very common ($\geq 1/10$); Common ($\geq 1/100$ and $1/10$); Uncommon ($\geq 1/1000$ and $1/100$); Rare ($\geq 1/10,000$ and $1/1000$); Very rare ($< 1/10,000$); Not known (cannot be estimated from the available data). Within each frequency grouping, adverse events are presented in order of decreasing seriousness.

System Organ Class	Frequency	Adverse Events
Immune System Disorders	Not known	Hypersensitivity
Skin and Subcutaneous Tissue Disorders	Not known	Skin sensitisation, dermatitis contact, skin discolouration, application site burn

Symptoms and Treatment of Overdose

Topical application of undiluted chloroxylenol can cause skin burning. Symptoms reported include corrosion of the oral mucosa, larynx, and the gastrointestinal tract, bradycardia, hypotension, and renal failure. Large amounts may cause CNS depression. Pulmonary aspiration of chloroxylenol-based disinfectants may result in pneumonia, acute respiratory distress syndrome, and cardiorespiratory arrest. There have been reports of death by excessive consumption.

Oral ingestion may result in pharyngeal erosion, laryngeal oedema, stomatitis, bradycardia, hypotension, renal failure and CNS depression. Pulmonary aspiration following ingestion may result in pneumonia, acute respiratory distress syndrome and cardiorespiratory arrest. There have been reports of death by excessive consumption.

In the case of ingestion or excess exposure, seek medical advice immediately. Careful observation of airway patency for 24-48 hours should be made post-ingestion.

Effects on ability to drive and use machines

Not applicable.

Storage Conditions

Keep container tightly closed. Protect from heat and light. Store below 30°C.

Dosage Forms and Packaging Available

30ml, 50ml, 250ml, 500ml and 750ml bottle

1L bottle

5L bottle (For Export Only)

Registration Number:

MAL

Manufactured by & Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.
Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia.

Date of Revision:

1 July 2022