

1. Product Name:

VEINOSMIN Film Coated Tablets

2. Qualitative and Quantitative Composition:

Each film-coated tablet contains:

Diosmin 450mg and Hesperidin 50mg

Excipients:

Microcrystalline cellulose powder, maize starch, colloidal anhydrous silica, povidone K 30, sodium lauryl sulphate, citric acid, purified water, magnesium stearate, purified talc, croscarmellose sodium, titanium dioxide, yellow iron oxide, red iron oxide

3. Dosage Form:

Film-coated tablet

4. Product Description:

Peach colored oval shaped film coated tablet having break line on one side.

5. Pharmacodynamics:

In pharmacology:

VEINOSMIN exerts an action on the vascular return system:

- at the venous level, it reduces venous distensibility and reduces venous stasis;
- at the microcirculatory level, it normalises capillary permeability and reinforces capillary resistance.

In clinical pharmacology:

Controlled, double-blind studies using methods that allow demonstrating and quantifying the activity on venous haemodynamics have confirmed the pharmacological properties of this medicinal product in humans.

- dose/effect relationship:

Statistically-significant dose-effect relationships have been demonstrated for the following venous plethysmography parameters: capacitance, distensibility and emptying time. The best dose/effect ratio is obtained with 2 tablets.

- venotonic activity:

It increases venous tone: venous occlusion plethysmography with a mercury strain gauge revealed a reduction in venous emptying time.

- microcirculatory activity:

Controlled, double-blind studies have demonstrated a statistically-significant difference between this medicinal product and placebo. In patients with signs of capillary fragility, it increases capillary resistance as measured by angiostrometry.

6. Pharmacokinetics:

In humans, following oral administration of the medicinal product with carbon 14-labelled diosmin:

- Excretion is essentially faecal and urinary excretion is on average 14% of the administered quantity,
- The elimination half-life is 11 hours,
- The product is highly metabolised, this metabolism is revealed by the presence of different phenol acids in the urine.

7. Indication:

- Treatment of symptoms related to venolymphatic insufficiency (heavy legs, pain, early morning restless legs).
- Treatment of functional symptoms related to acute hemorrhoidal attack.

8. Recommended Dose:

- Venolymphatic insufficiency: 2 tablets daily in two divided doses, midday and evening at meal times.
- Acute hemorrhoidal attack : 6 tablets daily (in 3 divided doses) for the first 4 days, then 4 tablets per day (in 2 divided doses) for the next 3 days.

Paediatric population

The safety and efficacy of VEINOSMIN 500mg in children and adolescents under 18 years of age have not been established.

9. Route of Administration:

Oral

10. Contraindications:

Hypersensitivity to the active substance or to any of the excipients.

11. Warnings and Precautions:

Hemorrhoidal attack:

The administration of this product does not preclude treatment for other anal conditions. The treatment must be short-term. If symptoms do not subside promptly, a proctological examination should be performed and the treatment should be reviewed.

12. Interactions with Other Medicaments:

None

13. Pregnancy and Lactation:

Pregnancy:

There are no or limited amount of data from the use of micronised purified flavonoid fraction in pregnant women.

Animal studies do not indicate reproductive toxicity.

As a precautionary measure, it is preferable to avoid the use of VEINOSMIN during pregnancy.

Breast-feeding:

It is unknown whether the active substance/metabolites are excreted in human milk.

A risk to the newborns/infants cannot be excluded.

A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from VEINOSMIN therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

Fertility:

Reproductive toxicity studies showed no effect on fertility in male and female rats.

14. Side Effects:

Summary of the safety profile

Side effects reported with VEINOSMIN in clinical trials are of mild intensity. They consist mainly in gastro intestinal events (diarrhoea, dyspepsia, nausea, vomiting).

Tabulated list of adverse reactions

The following adverse effects or events have been reported and are ranked using the following frequency: very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$), not known (cannot be estimated from the available data).

System Organ Class	Frequency	Preferred Term
Nervous system disorders	Rare	Dizziness
		Headache
		Malaise
Gastrointestinal disorders	Common	Diarrhoea
		Dyspepsia
		Nausea
		Vomiting
	Uncommon	Colitis
	Not known*	Abdominal pain
Skin and subcutaneous tissue disorders	Rare	Pruritus
		Rash
		Urticaria
	Not known*	Isolated face, lip, eyelid oedema. Exceptionally, Quincke's oedema

* Post-marketing experience

15. Symptoms and Treatment of Overdose:

Symptoms

There is limited experience with VEINOSMIN overdose. The most frequently reported adverse events in overdose cases were gastrointestinal events (such as diarrhoea, nausea, abdominal pain) and skin events (such as pruritus, rash).

Management

Management of overdose should consist in treatment of clinical symptoms.

16. Effects on Ability to Drive and Use Machines:

None

17. Storage Condition:

Store below 30°C.

18. Shelf Life:

36 months

Manufactured by:

Saga Lifesciences Limited

Survey No. 198/2 & 198/3, Chachrawadi Vasna,
Ta Sannand District,
Ahmedabad – 382210,
Gujarat, India

Product Registration Holder:

Perniagaan Normahas Sdn Bhd
1-5-10, 5th Block C, Diamond
Square(Jalan Semarak API),
Jalan 1/50, Off Jalan
Gombak 53000 Kuala Lumpur

Date of revision: 04 August 2025