

HP GLUCOSAMINE CAPSULE 500MG

VICOSxx-02

DESCRIPTION

Reddish orange opaque and white opaque with 'CM500' printed on one end and 'hovid' on the other end of the capsule filled with white to light yellowish granules.

COMPOSITION

Each capsule contains Glucosamine Sulphate Potassium Chloride 663.2 mg (equivalent to 500mg of Glucosamine Sulphate).

PHARMACODYNAMICS

Glucosamine possesses antiarthrosic-arthrotrophic activities. It stimulates the anabolic metabolism of osteo-cartilaginous tissues via stimulation of the biosynthesis of the mucopolysaccharides (which are the essential components of the cartilage ground substances) and of the bone mesenchymal tissues. Glucosamine also acts to improve the viscosity of synovial fluid by increasing synovial fluid production, thereby providing lubricant activity.

PHARMACOKINETICS**Absorption**

Glucosamine sulphate is well absorbed (with a proportion close to 90%) following oral administration.

Protein binding

It is known that glucosamine and its metabolites bind to α - and β -globulins.

Distribution

Mainly distributed to bone tissues and articular cartilages.

INDICATIONS

Adjunctive therapy in osteoarthritis.

CONTRAINDICATIONS

Contraindicated in patients sensitive to glucosamine sulphate and shellfish.

WARNINGS AND PRECAUTIONS

HP Glucosamine is not meant to replace NSAIDs in cases of intense pain, as this is a causal therapy. It is advisable to take an anti-inflammatory drug in the initial treatment of osteoarthritis together with glucosamine sulphate.

Therapeutic effect can only be seen after approximately 1 week from the beginning.

HP Glucosamine is to be taken under medical supervision in pregnant and lactating mothers, and in geriatric patients.

SIDE EFFECTS**Cardiovascular**

Peripheral oedema, tachycardia were reported in a few patients following larger clinical trials investigating oral administration in osteoarthritis. Causal relationship has not been established.

Central nervous system

Drowsiness, headache, insomnia have been observed rarely during therapy (less than 1%).

Gastrointestinal

Nausea, vomiting, diarrhoea, dyspepsia or epigastric pain, constipation, heartburn and anorexia have been described rarely during oral therapy with glucosamine.

Skin

Skin reactions such as erythema and pruritus have been reported with therapeutic administration of glucosamine.

DRUG INTERACTIONS

None known.

SYMPTOMS OF OVERDOSE AND TREATMENT

None known.

DOSAGE AND ADMINISTRATION

For mild to moderate osteoarthritis symptoms: Oral, 1 capsule 2 times daily for 6 weeks, taken 15-30 minutes before meal.

For severe osteoarthritis symptoms: Oral, 1 capsule 3 times daily for 8 weeks, taken 15-30 minutes before meal.

Maintenance therapy should be followed for 3-4 months by administration of 1 capsule twice daily.

Note: The information given here is limited. For further information consult your doctor or pharmacist.

Storage: Store below 30°C. Protect from moisture.

Presentation/Packing:

Glass/Plastic bottle of 60's.

Product Registration Holder:

HOVID Pharmacy Sdn. Bhd.
121, Jalan Tunku Abdul Rahman,
30010 Ipoh, Perak, Malaysia.

Manufactured by:

HOVID Bhd.
Lot 56442, 7½ Miles, Jalan Ipoh/Chemor,
31200 Chemor, Perak, Malaysia

Date of Revision: September 2022