

QKP MADURAMICIN PREMIX 1% W/W POWDER

Composition:

Maduramicin Ammonium.....10mg

Exp. Qs.....1g

Description

White Colour Powder

Pharmacodynamic

Maduramicin is considerably more potent as a coccidiostat than the other polyether ionophores that are used as coccidiostatic feed additives. These compounds are branchchained, polyoxygenated carboxylic acids that act as mobile carriers of cations by rendering cations lipid-soluble, thereby enabling them to pass across membranes. This process disrupts cationic cross-membrane gradients and is responsible for their anticoccidial activity.

As other polyether ionophores, maduramicin ammonium possesses moderate activity against many gram-positive bacteria, but with no activity against gram-negative organisms.

Pharmacokinetic

Maduramicin ammonium is absorbed (to an unknown extent) and excreted rapidly in chicken. Under continuous administration a distribution/metabolism steady state is observed after three days in the excreta but after six days in plasma.

Indication

As an aid in the prevention of coccidiosis in broiler chickens caused by *Eimeria tenella*, *E. (Eimeria) acervulina*, *E. (Eimeria) maxima*, *E. (Eimeria) necatrix*, *E. (Eimeria) mitis*, and *E. (Eimeria) brunetti*.

Route of Administration

Oral

Recommended dose

5 mg/kg (milligrams per kilogram) (0.0005%) of maduramicin ammonium in the complete feed. Feed this medicated feed continuously as the sole ration from day one to five days prior to slaughter.

Contraindication

Dangerous for equine species.

This feedingstuff contains an ionophore: simultaneous use with certain medicinal substances (e.g. tiamulin) can be contraindicated.

Interaction with other medicinal products

it would be advisable to avoid concurrent administration of maduramicin ammonium with tiamulin and certain other medicinal substances

Pregnancy and lactation

Do not feed to replacement, laying or breeding chickens

Side effects

Although maduramicin ammonium, in common with other ionophores, can inhibit the growth of Gram-positive micro-organisms, it is unlikely that this would result in any measurable adverse effect on the overall gut flora.

Symptoms and treatment of overdose

Do not exceed the recommended level of 5 ppm (mg/kg) as reduced weight gains may result with overdosage.

Storage

Store in a cool dry place (<30°C). Strictly avoid light and heat exposure.

Warning and Precaution

Keep out of reach of children / Jauhi ubat dari kanak-kanak

1. Consult a veterinarian or a poultry pathologist if losses exceed 0.5% in a two-day period.
2. Do not use this feed for treatment of outbreaks of coccidiosis.
3. Do not use in feeds containing pellet binding agents with the exception of bentonite, Pel-Aid or lignosulfonate.
4. Do not feed to replacement or breeding chickens.

Shelf life

24 months

Shelf life after opening

24 hours

Shelf life after dilution or constitution

24 hours

Withdrawal period

Treated chickens must not be slaughtered for use in food for at least 5 days after treatment with this medicated feed.

Packaging

25kg

Revised date

12/1/2022

Product Registration Holder:

PANFAST MARKETING (M) SDN.BHD.

PENANG SCIENCE PARK 1102, MUKIM 13, LORONG PERINDUSTRIAN BUKIT MINYAK 18, KAWASAN PERINDUSTRIAN BUKIT MINYAK, 14100 SIMPANG AMPAT, PENANG, MALAYSIA
TEL: +604-5016868 FAX: +604-5016866

Manufactured by:

SHANDONG QILU KING-PHAR PHARMACEUTICAL CO., LTD.

21 QINGLONG ROAD, PINGYIN, JINAN, 250400, CHINA
Tel: +86 531 83127771 Fax : +86 531 83127774