

AETOS KETOTECH ^{30mg} PLASTER

COMPOSITION

Each plaster contains:

Ketoprofen 30 mg

APPEARANCE

Plaster spraying colorless, translucent adhesives to a flesh-colored nonwoven fabric and covered with a release paper.

PHARMACODYNAMICS

Pharmacotherapeutic group: Anti-inflammatory preparations, non-steroids for topical use. ATC code: M02AA10.

Mechanism of action

Ketoprofen is an inhibitor of both the cyclo-oxygenase and lipoxygenase pathways. In addition, ketoprofen is a powerful inhibitor of bradykinin (a chemical mediator of pain and inflammation).

Pharmacodynamic effects

Inhibition of prostaglandin synthesis provides for potent anti-inflammatory, analgesic and antipyretic effects. Lipoxygenase inhibitors appear to attenuate cell-mediated inflammation and thus retard the progression of tissue destruction in inflamed joints. Ketoprofen stabilises the lysosomal membranes against osmotic damage and prevents the release of lysosomal enzymes that mediate tissue destruction in inflammatory reactions.

PHARMACOKINETICS

By cutaneous route, absorption is very low. The percutaneous application of 50-150 mg of ketoprofen produces plasma levels of the active ingredient of 0.08- 0.15 µg/mL approx. 5-8 hours after application. The systemic passage of the plaster compared to that of the oral formulations of ketoprofen is around 5 per cent, which enables a local effect to be obtained without systemic incidence.

INDICATIONS

For the treatment of the signs and symptoms of the following:

- Arthritis deformans (rheumatoid arthritis) • Periarthritis humero-scapularis (frozen, painful or stiff shoulder)
- Peritendinitis • Tendinitis • Muscular pain • Pain and swelling resulted from trauma (contusion, distortion etc.)

RECOMMENDED DOSAGE

Adult: Apply one plaster to the affected area after removal of release paper twice daily.

ROUTE OF ADMINISTRATION

Topical

CONTRAINDICATIONS

1. Patients with hypersensitivity to this medicine or any of its components.
2. History of any photosensitivity reaction.
3. Patients who have or had Aspirin-induced asthma or asthma attack caused by other non- steroidal anti-inflammatory drugs (NSAID).
4. Patients with hypersensitivity to the following drugs: tiaprofenic acid, fenofibrate, bezafibrate, ciprofibrate, oxybenzone. They may cause cross-hypersensitivity with ketoprofen.
5. Sun exposure, even in the case of hazy sun, including UV light from solarium, during the treatment and 2 weeks after its discontinuation.
6. Third trimester of pregnancy
7. Pathological skin changes such as dermatosis, eczema or acne, infected skin lesions, or open wounds.

WARNING AND PRECAUTION

Risk of GI Ulceration, Bleeding and Perforation with NSAID

Serious GI toxicity such as bleeding, ulceration and perforation can occur at any time, with or without warning symptoms, in patients treated with NSAID therapy. Although minor upper GI problems (e.g. dyspepsia) are common, usually developing early in therapy, prescribers should remain alert for ulceration and bleeding in patients treated with NSAIDs even in the absence of previous GI tract symptoms.

Studies to date have not identified any subset of patients not at risk of developing peptic ulceration and bleeding. Patients with prior history of serious GI events and other risk factors associated with peptic ulcer disease (e.g. alcoholism, smoking, and corticosteroid therapy) are at increased risk. Elderly or debilitated patients seem to tolerate ulceration or bleeding less than other individuals and account for most spontaneous reports for fatal GI events.

1. Used with caution in patients with reduced heart, liver or renal function: isolated cases of systemic adverse reactions affecting renal function have been reported.
2. During application of this plaster or 2 weeks after its application, avoid exposure to UV light on application area as it has been reported that photosensitivity to UV may occur during or after application.
3. The topical use of large amounts of product may give rise to systemic effects such as hypersensitivity and asthma.
4. The treatment should be interrupted if rash appears.
5. The recommended length of treatment should not be exceeded due to the risk of developing contact dermatitis and photosensitivity reactions increases over time.
6. Hands should be washed thoroughly after each application of the product.

- 7. Treatment should be discontinued immediately upon development of any skin reaction including cutaneous reactions after co-application of octocrylene containing products.
- 8. Do not use with occlusive dressings.
- 9. Patients with asthma combined with chronic rhinitis, chronic sinusitis, and/or nasal polyposis have a higher risk of allergy to aspirin and/or NSAIDs than the rest of the population.
- 10. The use of topical products, especially if it is prolonged, may give rise to phenomena of sensitisation or local irritation.
- 11. Keep out of the sight and reach of children.
- 12. Since skin infection can be masked, if you use this patch for inflammation with infection, use this product in combination with appropriate antibiotics or antifungal agents.

INTERACTION WITH OTHER MEDICAMENTS

Interactions are unlikely as serum concentrations following topical administration are low. It is, however, advisable to monitor patients under treatment with coumarinic substances.

STATEMENT ON USAGE DURING PREGNANCY AND LACTATION

Pregnancy

During the first and second trimester:

As the safety of ketoprofen in pregnant women has not been evaluated, the use of ketoprofen during the first and second trimester of pregnancy should be avoided.

During the third trimester of pregnancy:

All prostaglandin synthetase inhibitors including ketoprofen may induce cardiopulmonary and renal toxicity in the foetus. At the end of the pregnancy, prolonged bleeding time in both the mother and child may occur. Therefore, ketoprofen is contraindicated during the last trimester of pregnancy.

Breast-feeding

No data are available on excretion of ketoprofen in human milk. Ketoprofen is not recommended in nursing mothers.

ADVERSE REACTIONS

The following CIOMS frequency rating is used: Very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1000$ to $< 1/100$); rare ($\geq 1/10\,000$ to $< 1/1000$); very rare ($< 1/10\,000$), not known (cannot be estimated from the available data).

Infections and infestations	Skin and subcutaneous tissue disorders
Not known: Secondary impetigo	Uncommon: Local skin reactions such as rash, erythema, eczema, pruritus and burning sensation, application site burn.
Blood and lymphatic system disorders	Rare: Photosensitisation, urticaria, bullous/contact/exfoliative/ vesicular dermatitis, phlyctenular eczema, blister, photosensitivity reaction, allergic reaction, skin exfoliation, skin oedema.
Not known: Eosinophilia	
Immune system disorders	Renal and urinary disorders
Not known: anaphylactic shock, angioedema, hypersensitivity reactions	Very rare: Cases of aggravation of previous renal insufficiency, acute renal failure
Eye disorders	
Not known: Eyelid oedema	
Vascular disorders	General disorders and administration site conditions
Not known: Vasculitis	Not known: Pyrexia
Gastrointestinal disorders	Injury, poisoning and procedural complications
Not known: Peptic ulcer, gastrointestinal bleeding, diarrhoea, lip oedema	Not known: Wound complication

OVERDOSE AND TREATMENT

Overdosage is unlikely to be caused by topical administration. However, the topical use of large amounts of product may give rise to systemic effects such as hypersensitivity and asthma. If they occur, treatment should be supportive and symptomatic.

STORAGE

Store below 30°C in a light-resistant tightly closed container

SHELF LIFE

3 years

DOSAGE FORM AND PACKAGING AVAILABLE

7 plasters in a polyethylene-aluminium pouch, 50 pouches in an outer carton.

Manufactured by:

Sinil Pharmaceutical Co., Ltd.

28, Boksanggol-gil, Anseong-myeon, Chungju-si, Chungcheongbuk-do, Korea.

Revision date: December 2022

Title : Aetos Ketotech Plaster (Leaflet)
Size : 123mm(w) x 260mm(h)
Colour : CMYK
Font : Arial Bold (size 8) , Myriad Regular (size 7) , Myriad Pro Semibold Italic (size 7)

123 mm

AETOS KETOTECH 30mg PLASTER

COMPOSITION

Each plaster contains:
Ketoprofen 30 mg

APPEARANCE

Plaster spraying colorless, translucent adhesives to a flesh-colored nonwoven fabric and covered with a release paper.

PHARMACODYNAMICS

Pharmacotherapeutic group: Anti-inflammatory preparations, non-steroids for topical use. ATC code: M02AA10.

Mechanism of action

Ketoprofen is an inhibitor of both the cyclo-oxygenase and lipoxygenase pathways. In addition, ketoprofen is a powerful inhibitor of bradykinin (a chemical mediator of pain and inflammation).

Pharmacodynamic effects

Inhibition of prostaglandin synthesis provides for potent anti-inflammatory, analgesic and antipyretic effects. Lipoxygenase inhibitors appear to attenuate cell-mediated inflammation and thus retard the progression of tissue destruction in inflamed joints. Ketoprofen stabilises the lysosomal membranes against osmotic damage and prevents the release of lysosomal enzymes that mediate tissue destruction in inflammatory reactions.

PHARMACOKINETICS

By cutaneous route, absorption is very low. The percutaneous application of 50-150 mg of ketoprofen produces plasma levels of the active ingredient of 0.08- 0.15 µg/mL approx. 5-8 hours after application. The systemic passage of the plaster compared to that of the oral formulations of ketoprofen is around 5 per cent, which enables a local effect to be obtained without systemic incidence.

INDICATIONS

For the treatment of the signs and symptoms of the following:
• Arthritis deformans (rheumatoid arthritis) • Periarthritis humelo-scapularis (frozen, painful or stiff shoulder)
• Peritendinitis • Tendinitis • Muscular pain • Pain and swelling resulted from trauma (contusion, distortion etc.)

RECOMMENDED DOSAGE

Adult: Apply one plaster to the affected area after removal of release paper twice daily.

ROUTE OF ADMINISTRATION

Topical

CONTRAINDICATIONS

1. Patients with hypersensitivity to this medicine or any of its components.
2. History of any photosensitivity reaction.
3. Patients who have or had Aspirin-induced asthma or asthma attack caused by other non- steroidal anti-inflammatory drugs (NSAID).
4. Patients with hypersensitivity to the following drugs: tiaprofenic acid, fenofibrate, bezafibrate, ciprofibrate, oxybenzone. They may cause cross-hypersensitivity with ketoprofen.
5. Sun exposure, even in the case of hazy sun, including UV light from solarium, during the treatment and 2 weeks after its discontinuation.
6. Third trimester of pregnancy
7. Pathological skin changes such as dermatosis, eczema or acne, infected skin lesions, or open wounds.

WARNING AND PRECAUTION

Risk of GI Ulceration, Bleeding and Perforation with NSAID

Serious GI toxicity such as bleeding, ulceration and perforation can occur at any time, with or without warning symptoms, in patients treated with NSAID therapy. Although minor upper GI problems (e.g. dyspepsia) are common, usually developing early in therapy, prescribers should remain alert for ulceration and bleeding in patients treated with NSAIDs even in the absence of previous GI tract sympto ms.

Studies to date have not identified any subset of patients not at risk of developing peptic ulceration and bleeding. Patients with prior history of serious GI events and other risk factors associated with peptic ulcer disease (e.g. alcoholism, smoking, and corticosteroid therapy) are at increased risk. Elderly or debilitated patients seem to tolerate ulceration or bleeding less than other individuals and account for most spontaneous reports for fatal GI events.

1. Used with caution in patients with reduced heart, liver or renal function: isolated cases of systemic adverse reactions affecting renal function have been reported.
2. During application of this plaster or 2 weeks after its application, avoid exposure to UV light on application area as it has been reported that photosensitivity to UV may occur during or after application.
3. The topical use of large amounts of product may give rise to systemic effects such as hypersensitivity and asthma.
4. The treatment should be interrupted if rash appears.
5. The recommended length of treatment should not be exceeded due to the risk of developing contact dermatitis and photosensitivity reactions increases over time.
6. Hands should be washed thoroughly after each application of the product.

260 mm

[말레이시아]하루펜 설명서
123 X 260 양면인쇄

123 mm

7. Treatment should be discontinued immediately upon development of any skin reaction including cutaneous reactions after co-application of octocrylene containing products.
8. Do not use with occlusive dressings.
9. Patients with asthma combined with chronic rhinitis, chronic sinusitis, and/or nasal polyposis have a higher risk of allergy to aspirin and/or NSAIDs than the rest of the population.
10. The use of topical products, especially if it is prolonged, may give rise to phenomena of sensitisation or local irritation.
11. Keep out of the sight and reach of children.
12. Since skin infection can be masked, if you use this patch for inflammation with infection, use this product in combination with appropriate antibiotics or antifungal agents.

INTERACTION WITH OTHER MEDICAMENTS

Interactions are unlikely as serum concentrations following topical administration are low. It is, however, advisable to monitor patients under treatment with coumarinic substances.

STATEMENT ON USAGE DURING PREGNANCY AND LACTATION

Pregnancy

During the first and second trimester:

As the safety of ketoprofen in pregnant women has not been evaluated, the use of ketoprofen during the first and second trimester of pregnancy should be avoided.

During the third trimester of pregnancy:

All prostaglandin synthetase inhibitors including ketoprofen may induce cardiopulmonary and renal toxicity in the foetus. At the end of the pregnancy, prolonged bleeding time in both the mother and child may occur. Therefore, ketoprofen is contraindicated during the last trimester of pregnancy.

Breast-feeding

No data are available on excretion of ketoprofen in human milk. Ketoprofen is not recommended in nursing mothers.

ADVERSE REACTIONS

The following CIOMS frequency rating is used: Very common (≥1/10); common (≥1/100 to <1/10); uncommon (≥ 1/1000 to <1/100); rare (≥ 1/10 000 to <1/1000); very rare (<1/10 000), not known (cannot be estimated from the available data).

Infections and infestations Not known: Secondary impetigo	Skin and subcutaneous tissue disorders Uncommon: Local skin reactions such as rash, erythema, eczema, pruritus and burning sensation, application site burn.
Blood and lymphatic system disorders Not known: Eosinophilia	Rare: Photosensitisation, urticaria, bullous/contact/exfolia- tive/ vesicular dermatitis, phlyctenular eczema, blister, photosensitivity reaction, allergic reaction, skin exfoliation, skin oedema.
Immune system disorders Not known: anaphylactic shock, angioedema, hypersensitivity reactions	Renal and urinary disorders Very rare: Cases of aggravation of previous renal insufficiency, acute renal failure
Eye disorders Not known: Eyelid oedema	General disorders and administration site conditions Not known: Pyrexia
Vascular disorders Not known: Vasculitis	
Gastrointestinal disorders Not known: Peptic ulcer, gastrointestinal bleeding, diarrhoea, lip oedema	Injury, poisoning and procedural complications Not known: Wound complication

OVERDOSE AND TREATMENT

Overdosage is unlikely to be caused by topical administration. However, the topical use of large amounts of product may give rise to systemic effects such as hypersensitivity and asthma. If they occur, treatment should be supportive and symptomatic.

STORAGE

Store below 30°C in a light-resistant tightly closed container

SHELF LIFE

3 years

DOSAGE FORM AND PACKAGING
AVAILABLE

7 plasters in a polyethylene-aluminium pouch, 50 pouches in an outer carton.

Manufactured by:

Sinil Pharmaceutical Co., Ltd.

28, Boksanggol-gil, Angseong-myeon, Chungju-si,
Chungcheongbuk-do, Korea.

Revision date: December 2022

260 mm