

TERIPIN-S Solution for Injection 1mg/8.5mL

(Terlipressin Acetate 1mg/8.5mL)

【Appearance】

Colorless and transparent injection in a colorless and transparent ampoule

【Composition】

Each ampoule (8.5 mL) contains
- Active ingredient: Terlipressin Acetate ----- 1.0 mg (0.85 mg as Terlipressin)
- Inactive Ingredients: Acetic acid anhydride, sodium acetate trihydrate, sodium chloride and water for injection

【Indications】

- Bleeding oesophageal varices.
- Treatment of type 1 hepatorenal syndrome, characterized by spontaneous acute renal insufficiency, in patient suffering from severe cirrhosis with ascites.

【Dosage and Administration】

Bleeding oesophageal varices

Adults: Initially an i.v. injection of 1.7 mg terlipressin is given every 4 hours. The treatment should be maintained until bleeding has been controlled for 24 hours, but up to a maximum of 48 hours. After the initial dose, the dose can be adjusted to 0.85mg terlipressin i.v. every 4 hours in patients with body weight < 50 kg or if adverse effects occur.

Type 1 hepatorenal syndrome

2.55 to 3.4 mg of terlipressin every 24 hours as 3 or 4 administrations.
In the absence of any reduction of the serum creatinine after 3 days of treatment, cessation of Teripin-S Injection treatment is advised.
In the other cases, Teripin-S Injection treatment is to be pursued until the obtaining either of a serum creatinine less than 130 µmol/litre or of a drop of at least 30 % in the serum creatinine with respect to the value measured at the time of diagnosis of hepatorenal syndrome.
The standard average duration of treatment is 10 days.

【Method of Administration】

I.V. Injection

【Caution】

1. Warning
Cut ampoule injections carefully to minimize glass fragments, as fragments of glass can be mixed in while cutting the container and cause adverse effects. Take special caution when using with children and the elderly and infirm.

【Contraindications】

Contraindicated in pregnancy.
Hypersensitivity to terlipressin acetate or any of the excipients

【Special warnings and precautions for use】

Blood pressure, heart rate and fluid balance should be monitored during treatment.
To avoid local necrosis at the injection site, the injection must be given i.v.
Caution should be exercised in treating patients with hypertension or recognised heart disease.
In patients with septic shock with a low cardiac output, Teripin-S should not be used.
Children and the elderly: Particular caution should be exercised in the treatment of children and elderly patients, as experience is limited in these groups.
There is no data available regarding dosage recommendation in these special patient categories.

【Adverse events】

Table: Frequency of undesirable effects

SYSTEM ORGAN CLASS	Frequency		
	COMMON	UNCOMMON	RARE
Metabolism and nutrition disorders		Hyponatraemia if fluid not monitored	
Nervous system Disorders	Headache		
Cardiac disorders	Bradycardia	Atrial fibrillation Ventricular extrasystoles Tachycardia Chest pain Myocardial infarction Fluid overload with pulmonary oedema Torsade de pointes Cardiac failure	
Vascular disorders	Peripheral vasoconstriction Peripheral ischaemia Facial pallor Hypertension	Intestinal ischaemia Peripheral cyanosis Hot flushes	
Respiratory thoracic and mediastinal disorders		Respiratory distress Respiratory failure	Dyspnoea
Gastrointestinal disorders	Transient abdominal cramps Transient diarrhoea	Transient nausea Transient vomiting	
Skin and subcutaneous tissue disorders		Skin necrosis	
Pregnancy, puerperium and perinatal conditions		Uterine hypertonus Uterine ischaemia	
General disorders and administration site disorders		Injection site necrosis	

Reporting of suspected adverse reactions

【Interaction】

The hypotensive effect of non-selective beta-blockers on the portal vein is increased with terlipressin. Concomitant treatment with medicinal products with a known bradycardic effect (e.g. propofol, sufentanil) may lower the heart rate and cardiac output. These effects are due to reflexogenic inhibition of cardiac activity via the vagus nerve due to the elevated blood pressure.

【Administration to Pregnant Women】

Pregnancy

Treatment with Teripin-S during pregnancy is contraindicated.
Teripin-S has been shown to cause uterine contractions and increased intrauterine pressure in early pregnancy and may decrease uterine blood flow. Glypressin may have harmful effects on pregnancy and foetus.
Spontaneous abortion and malformation have been shown in rabbits after treatment with Teripin-S.
Breast-feeding

It is not known whether terlipressin is excreted in human breast milk. The excretion of terlipressin in milk has not been studied in animals. A risk to the suckling child cannot be excluded. A decision on whether to continue/discontinue breast-feeding or to continue/discontinue therapy with terlipressin should be made taking into account the benefit of breast-feeding to the child and the benefit of terlipressin therapy to the woman.

【Effects on ability to drive and use machines】

No studies on the effects on the ability to drive and use machines have been performed.

【Treatment for Overdose】

The recommended dose (2mg/4 hours) should not be exceeded as the risk of severe circulatory adverse effects is dose-dependent.

【Pharmacodynamic properties】

Pharmacotherapeutic group: Posterior pituitary lobe hormones (vasopressin and analogues) (H 01 BA 04)
Teripin-S may be regarded as a circulating depot of lysine vasopressin. Following intravenous injection, three glycyI moieties are enzymatically cleaved from the N-terminus to release lysine vasopressin.
The slowly released vasopressin reduces blood flow in the splanchnic circulation in a prolonged manner, thereby helping to control bleeding from ruptured oesophageal varices.

【Pharmacokinetics properties】

Teripin-S is administered by bolus intravenous injection. It shows a biphasic plasma level curve which indicates that a two compartment model can be applied.
The half-life of distribution (T_{1/2α}) is about 8 -10 minutes.
The half-life of elimination (T_{1/2β}) is about 50 -70 minutes.
Lysine vasopressin reaches maximum plasma levels about 1 - 2 hours following intravenous administration and has a duration of activity of 4 - 6 hours.

【Storage Condition】

Store in a refrigerator (2 - 8 °C).
Keep the ampoules in the outer carton in order to protect from light.

Keep out of reach of children.

Jauhi daripada kanak-kanak.

【Packaging Unit】

8.5 mL/ampoule × 5 ampoules / Box

【Shelf Life】

2 years

Manufacturer

Hanlim Pharm. Co., Ltd.
2-27, Yeongmun-ro, Cheoin-gu, Yongin-si, Gyeonggi-do, Korea

Product Registration Holder

Eucogen Pharma Sdn. Bhd.
6A, Jalan Sungai Burung U 32/U, Bukit Rimau, Seksyen 32, 40460 Shah Alam, Selangor, Malaysia

Date of Revision : 21 June 2022

'22. 06. 21
P36-00

MANUFACTURED BY
 **HANLIM PHARM.CO.,LTD.**
2-27, Yeongmun-ro, Cheoin-gu, Yongin-si, Gyeonggi-do, Korea
<http://www.hanlim.com>

HanLim Instructions

File Name(제품명)	테리핀 에스주 (등록용)	Paper Material	
Type(Material)	설명서	Coating	무코팅
Date(Y/M/D)	2022, 06, 21	Coaptation(접착)	
Version No.	P32-00	OTC/RX	RX
Country	말레이시아	라벨폴림	
Font Size	본문 : 6 Point	Presswork(인쇄소)	영일인쇄
Color	 검정	담당	■ 무역 : 이영석
Korea Drug Code		제조번호 인쇄가능 여부	
Bar Code		수정사항	내용
Size(규격) (접지)	120 × 265mm	비고	