

160x160

220929

200mg/ ml iron dextran injectable solution

FOR VETERINARY USE ONLY

WOOGENE FERRUM 200 inj.**COMPOSITION**

Each 1ml contains

Active ingredients

Iron dextran 200mg/ml

Excipients

Phenol 5mg/ml

PRODUCT DESCRIPTION

Woogene Ferrum 200 inj. is dark brown solution use for prevention and treatment of iron deficiency anemia in piglet. The product is packed in an amber bottle sealed with rubber stopper and aluminum cap.

PHARMACODYNAMIC

Iron is necessary for myoglobin and hemoglobin in the transport and utilization of oxygen. While neither stimulating erythropoiesis nor correcting hemoglobin abnormalities not caused by iron deficiency, iron administration does correct both physical signs and decreased hemoglobin levels secondary to iron deficiency. Ionized iron is a component in the enzymes cytochrome oxidase, succinic dehydrogenase, and xanthine oxidase.

PHARMACOKINETIC

After IM injection, iron dextran is slowly absorbed primarily via the lymphatic system. About 60% of the drug is absorbed within 3 days of injection and up to 90% of the dose is absorbed after 1 - 3 weeks. The remaining drug may be absorbed slowly over several months.

After absorption, the reticuloendothelial cells of the liver, spleen, and bone marrow gradually clear the drug from plasma. The iron is cleaved from the dextran component and the dextran is then metabolized or excreted. The iron is immediately bound to protein elements to form hemosiderin, ferritin or transferrin. Iron crosses the placenta, but in what form is unknown. Only traces of iron are excreted in milk.

Iron is not readily eliminated from the body. Iron liberated by the destruction of hemoglobin is reused by the body and only small amounts are lost by the body via hair and nail growth, normal skin desquamation, and GI tract sloughing. Accumulation can result with repeated dosing as only trace amounts of iron are eliminated in the feces, bile, or urine.

ENVIRONMENTAL PROPERTIES

Avoid release to the environment.

INDICATION

Prevention and treatment of iron deficiency anemia in piglet

RECOMMENDED DOSE

Inject 1ml per head via IM

DIRECTION OF USE

Intramuscular administration

SHELF LIFE & STORAGE CONDITION

Shelf Life as packaged for Sale: 2 Years

Shelf Life after first opening of container and reconstitution: 24 hours

WARNING/PRECAUTION

Contraindication

-Do not use this product to animals that have shock or hypersensitivity to this product before.

-Do not use this product to pregnant or lactating animals.

-Inject slowly to avoid hypersensitivity in piglets that were born from the mother pig with vitamin E and/or selenium deficiency.

Special warning for each target species

-Use only for designated animal.

-Use according to label indications, dosage and route of administrations.

-Use under veterinary prescription and after consulting with veterinarian.

Special precautions for storage

-Keep out of reach of children and animal.

-Keep unused medicine in a dry place and avoid direct sunlight

-Not intended for repacking, avoid transferring to other vials may affect quality.

-Do not remove the rubber stopper and use disposable needle to prevent contamination.

-Follow the storage conditions on the label to keep the validity and safety of the product.

WITHDRAWAL PERIOD

0 day

WARNING AND PRECAUTION

N/A

INTERACTION WITH OTHER MEDICINES

N/A

PREGNANT AND LACTATING ANIMALS

Do not use in pregnant or lactating animals.

ADVERSE EFFECTS

N/A

OVERDOSE

N/A

INCOMPATIBILITIES

Oxidizing agents

CONTROLLED MEDICINE**DISPOSAL OF CONTAINER**

Dispose of used package by wrapping in paper and placing in garbage.

DOSAGE FORM AND PACKING AVAILABLE

50ml, 100ml

DATE OF REVISION OF PACKAGE INSERT

18th February 2022

Distributor :



F.E Venture Sdn. Bhd.
Lot 3 & 5, Jalan PJS 11/8, Bandar Sunway,
46150 Petaling Jaya Selangor D.E., Malaysia.



Frontier In Bio & Green Technology

WOOGENE B&G

230, Jeongmunsongsan-ro, Yanggam-myeon,
Hwaseong-si, Gyeonggi-do, Republic of KOREA
Tel. 82-2-795-2361 Fax. 82-2-797-9863