
UniScab Lotion 1% w/v

Composition and Strength of Active Ingredients:

Gamma benzene hexachloride.....1% w/v

Preservatives:

Potassium sorbate..... 0.1% w/v

Cetylpyridinium chloride.....0.1% w/v

Product Description

White coloured lotion with floral smell.

Pharmacodynamics

Gamma benzene hexachloride is a central nervous system (CNS) stimulant when absorbed systematically. Following absorption through the chitinous exoskeleton of arthropods, Gamma benzene hexachloride is presumed to stimulate the nervous system, resulting in convulsions and death.

Pharmacokinetics

Absorption

The absorption of Gamma benzene hexachloride through the skin is highly variable. Although the incidence of adverse effects appears to be higher in infants and young children than in adults, there have been no studies to demonstrate that children absorb more Gamma benzene hexachloride than adults.

Lotion- In one study, a mean peak blood concentration of 3 nanograms per mL occurred in infants and children 6 hours after total body application of Gamma benzene hexachloride for scabies.

Distribution

Gamma benzene hexachloride is highly lipid- soluble and is stored in adipose tissue and other lipophilic tissue. Gamma benzene hexachloride is also stored in the white matter of the brain and may persist in low levels for up to 2 weeks following topical application.

Metabolism

A study in 9 healthy adults demonstrated a decreased half- life and increased clearance of antipyrine when preceded by a topical Gamma benzene hexachloride 1% application of 12 hour duration. This suggested that Gamma benzene hexachloride induces microsomal drug metabolism.

Elimination Half-life

17.9 to 21.4 hours

Indication

UniScab Lotion is indicated for the treatment of scabies infestation caused by *Sarcoptes Scabiei*

Use should be restricted to patients who have failed treatment with or cannot tolerate other medications that pose less risk.

Recommended Dosage

For Scabies (*Sarcoptes Scabiei*)

- i) It is recommended 30ml of UniScab lotion applied in a thin layer to all areas of the body from the neck down and thoroughly washed off after 8 hours. UniScab lotion should not be applied immediately after a bath or shower and should not be used by persons with extensive dermatitis or in pregnant or lactating women.
- ii) The lotion should be applied thinly and rubbed in thoroughly from the neck down to the soles of the feet, with special attention under nail and to the hands, feet and interdigital areas. Avoid applying UniScab lotion to open cuts or excoriations.
- iii) The mite does not infest areas of sebaceous gland activity in adult or children, and the face and scalp need not be treated. However, scabies may affect the head in infants,
- iv) The lotion should be left on for 8 to 12 hours and removed by thorough washing.
- v) One application is usually curative although many patients exhibit persistent pruritus after treatment. It is not known how soon after UniScab Lotion administration that a second dose can safely be readministered. Retreatment is not recommended.

Route of Administration

Topical

Contraindications

Gamma benzene hexachloride is contraindicated for premature neonates because their skin may be more permeable than full term infants and their liver enzymes may not be sufficiently developed. It is also contraindicated for patients with known seizure disorders and for individuals with a known sensitivity to the products or any of its components. The other contraindications are Norwegian (crusted) scabies and other skin conditions (e.g., atopic dermatitis, psoriasis) that may increase systemic absorption.

Warning and Precautions

WARNINGS:

Gamma benzene hexachloride should only be used in patients who cannot tolerate or have failed first-line treatment with safer medications for the treatment of scabies.

Neurologic Toxicity

Seizures and deaths have been reported following Gamma benzene hexachloride use with repeat or prolonged application, but also in rare cases following a single application used according to directions. Gamma benzene hexachloride should be used with caution for infants, children, the elderly, and individuals with others skin condition (e.g, atopic dermatitis, Psoriasis) and in those who weigh <110 Ibs (50kg) as they may be at risk of serious neurotoxicity.

Contraindication

Gamma benzene hexachloride is contraindicated in premature infants and individuals with known uncontrolled seizure disorders.

- NOT first-line agent
- Use only as directed
- CNS toxicity may be greater in the young, use with extreme caution in infants, children and pregnant and nursing mothers
- Avoid contact with eyes or mucous membranes
- Avoid inflamed or excoriated skin
- Avoid simultaneous use with other creams, ointments or oils
- Condition which may increase the risk of seizure (eg. HIV, history of head trauma or prior seizure, CNS tumor, excessive alcohol consumption)
- Elderly
- Concomitant medications which lower the seizure threshold
- Individuals weighing less than 50kg (110 lbs).
- Just put a very thin layer for the UniScab lotion
- Wash the UniScab lotion off your skin after 8 to 12 hours.
- You may still itch after you use UniScab Lotion. Even after all the scabies bugs are dead, they still make your skin itch for a few weeks. This does not mean you need more UniScab lotion
- Avoid retreatment with UniScab lotion.

Interactions with Other Medicaments

Drug interaction:

Oils may enhance absorption of gamma benzene hexachloride, therefore, patient or caregivers applying UniScab lotion should be warned about simultaneous use of creams, ointments or oils.

The following drug interactions and/or related problems have been selected on the basis of their potential clinical significance (possible mechanism in parentheses where appropriate)- not necessarily inclusive (major clinical significance);

Skin, scalp or hair preparation, others such as cream, lotions, ointments or oils (simultaneous application may increase the percutaneous absorption of Gamma benzene hexachloride).

Pregnancy & Lactation

Adequate and well- controlled studies in humans have not been done because Gamma benzene hexachloride is absorbed through the skin and has the potential of causing CNS toxicity, some clinicians do not recommend the use of Gamma benzene hexachloride during pregnancy. If Gamma benzene hexachloride is used, however, the recommended dosage should not be treated more than twice during pregnancy.

For breast feeding, problems in human have not been documented. However, Gamma benzene hexachloride is systematically absorbed and is distributed into breast milk. Although the concentrations found in human blood following topical application of Gamma benzene hexachloride make it unlikely

that breast milk will contain amounts of Gamma benzene hexachloride sufficient to cause toxicity, an alternate method of feeding the infants should be used for 2 days.

Side Effects

Gamma benzene hexachloride is absorbed through the skin and has the potential for CNS toxicity, especially in infants, children and possibly the elderly.

The following side effects have been selected on the basis of their potential clinical significance (possible signs and symptoms in parentheses where appropriate)- not necessarily inclusive.

Those indicating need for medical attention.

Incidence rare:

Skin irritation not present before therapy-if Gamma benzene hexachloride is applied incorrectly or repeatedly, the incidence of skin irritation is increased; skin rash.

Symptoms of CNS toxicity:

Convulsions; dizziness, clumsiness or unsteadiness; fast heartbeat, muscle cramps; nervousness; restlessness or irritability; vomiting.

Those indicating need for medical attention, if they occur and continue or are bothersome after medication is discontinued. Itching of skin-acquired sensitivity to mites and their products may continue for one to several weeks

Other adverse effects in single patients that have been associated with Gamma benzene hexachloride are disseminated intravascular coagulation and subsequent death after oral ingestion and aplastic anaemia after prolonged topical exposure (twice daily application for three weeks).

Symptoms and Treatment of Overdose

Symptoms of overdose

Mild to moderate toxicity: Vomiting, headache, nausea, dizziness, diarrhea and tremor.

Severe Toxicity: Seizures, CNS depression, agitation, ataxia, hypotension, respiratory depression, metabolic acidosis and coma. Children appear to be more susceptible to the toxic effects of Gamma benzene hexachloride and more likely to manifest severe effects.

For the treatment of toxicity

- If accidental ingestion occurs, it may be life threatening and prompt gastric lavage is recommended.
- Since oils favor absorption, saline cathartics should be administered rather than oily cathartics for intestinal evacuation.
- If CNS manifestation occur, they may be treated by the administration of pentobarbital, phenobarbital or diazepam.

Effects on Ability to Drive and Use Machine

No effects were reported.

Storage Condition

Store below 30°C. Keep in a cool and dry place. Protect from light. Keep the bottle tightly closed.

Dosage Forms and Packaging Available

100ml bottle

1L bottle (For Export Only)

Registration Number:

XXXX

Manufactured by & Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia.

Date of Revision:

23 February 2023