



Introflor 300 mg/ml

Solution for Injection

NAME PRODUCT

Introflor 300 mg/ml Solution for Injection

DESCRIPTION APPEARANCE

A light-yellow solution.

QUALITATIVE AND QUANTITATIVE COMPOSITION PER ML

Florfenicol 300 mg.

TARGET SPECIES

Cattle and swine.

PHARMACODYNAMICS PROPERTIES AND

PHARMACOKINETIC DATA

Pharmacotherapeutic group: Antibacterial for systemic use: Amphenicols.

Pharmacodynamic properties

Florfenicol is a synthetic broad spectrum antibiotic effective against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol acts by inhibiting protein synthesis at the ribosomal level and is bacteriostatic. Laboratory tests have shown that florfenicol is active against the most commonly isolated bacterial pathogens involved in bovine and porcine respiratory disease which include *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni* and *Arcanobacterium pyogenes* for cattle and *Actinobacillus pleuropneumoniae* and *Pasteurella multocida* for swine. Acquired resistance to florfenicol is mediated by efflux pumps linked to a *floR* gene. Such resistance has not yet been identified in target pathogens other than *Pasteurella multocida*. Resistance to florfenicol and other antimicrobials has been demonstrated in food-derived *Salmonella typhimurium*. Co-resistance to florfenicol and other antibiotics (such as ceftiofur) has been observed in microorganisms of the *Enterobacteriaceae* family. Cross-resistance between phenicols, lincosamides, oxazolidinones, pleuromutins and streptogramin A antibiotics is known.

Pharmacokinetic particulars

Cattle:

Intramuscular administration at the recommended dose of 20 mg/kg maintains efficacious blood levels in cattle for 48 hours. Maximum mean serum concentration (C_{max}) of 3.37 µg/ml occurs at 3.3 hours (t_{max}) after dosing. The mean serum concentration 24 hours after dosing was 0.77 µg/ml.

Subcutaneous administration at the recommended dose of 40 mg/kg maintains efficacious blood levels in cattle (i.e. above the MIC_{50} of the main respiratory pathogens) for 63 hours. The mean elimination half-life is 18.3 hours. After intravenous administration of florfenicol to swine, the mean plasma clearance is 5.2 ml/min/kg and the mean volume of distribution is 948 ml/kg. The mean elimination half-life is 2.2 hours.

Swine:

After the first intramuscular administration of florfenicol in swine, maximum serum concentrations between 3.8 and 13.6 µg/ml are reached after 1.4 hours, the mean half-life for elimination is 3.6 hours. After the second intramuscular administration in swine, maximum serum concentrations between 3.7 and 3.8 µg/ml are reached after 1.8 hours. 12 to 24 hours after the i.m. administration, serum concentrations fall below 1 µg/ml, the MIC_{50} of the target porcine pathogens. Florfenicol concentrations achieved in lung tissue reflect plasma concentrations in an approximately 1:1 ratio. After intramuscular administration to swine, florfenicol is rapidly excreted, mainly in urine. Florfenicol is mostly metabolised.

INDICATIONS

Cattle:

Diseases caused by florfenicol-susceptible bacteria. Metaphylactic and therapeutic treatment of respiratory tract infections in cattle due to *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni*. The presence of the disease in the herd should be established before metaphylactic treatment.

Swine:

Treatment of acute outbreaks of swine respiratory disease caused by strains of *Actinobacillus pleuropneumoniae* and *Pasteurella multocida* susceptible to florfenicol.

CONTRAINDICATIONS

Not for use in cattle producing milk for human consumption.

Do not use in adult bulls, rams and boars intended for breeding purposes.

Do not use in piglets weighing less than 2 kg.

Do not use in the case of known hypersensitivity to the active substance or to any of the excipients.

WARNING AND PRECAUTIONS

Special warnings for each target species

Not authorised for use in animals producing milk for human consumption including pregnant animals intended to produce milk for human consumption.

Special precautions for use

Special precautions for use in animals

A separate needle or an inoculation gun should be used to withdraw the solution for injection in order to avoid puncturing the rubber stopper too frequently. The product should be used in conjunction with susceptibility testing and take into account official and local antimicrobial policies. Use of the veterinary medicinal product that deviates from the specifications in this product information can increase the prevalence of resistance to florfenicol and impair the effectiveness of therapy with other active substance (e.g. ceftiofur) as a result of cross-resistance.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Administration should be done carefully to avoid accidental self-injection. In case of accidental self-injection, seek medical advice immediately and show the package leaflet or the label to the doctor. Do not use if you are known to be sensitive to propylene glycol and polyethylene glycol.

INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION

None known.

USE DURING PREGNANCY, LACTATION

Cattle:

Studies in laboratory animals have not revealed any evidence of teratogenic or foetotoxic effects. However, the effect of florfenicol on bovine and ovine reproductive performance and pregnancy has not been assessed. Use only according to the benefit/risk assessment by the responsible veterinarian.

Swine:

The safety of the product in sows during pregnancy and lactation has not been proven. The use of the product during pregnancy and lactation in sows can therefore not be recommended.



Manufactured by:
Interchemie werken De Adelaar Eesti AS, Vanapere tee 14,
Püünsi, Viimsi, Harju county 74013, Estonia

Holland



DII16-MY



Introflor 300 mg/ml

Solution for Injection

ADVERSE REACTIONS (FREQUENCY AND SERIOUSNESS)

Cattle:

A decrease in food consumption and transient softening of the faeces may occur during the treatment period. The treated animals recover quickly and completely upon termination of treatment.

Intramuscular and subcutaneous administration may cause inflammatory lesions at the injection site which may persist for 14 days.

Swine:

Commonly observed adverse effects are transient diarrhoea and/or peri-anal and rectal erythema/oedema which may affect 50% of the animals. These effects can be observed for one week.

Under field conditions approximately 30% of treated swine presented with pyrexia (40 °C) associated with either moderate depression or moderate dyspnea a week or more after administration of the second dose.

Transient swelling lasting up to 5 days may be observed at the site of injection. Inflammatory lesions at the injection site may be seen up to 28 days.

OVERDOSE (SYMPTOMS, EMERGENCY PROCEDURES, ANTIDOTES), IF NECESSARY

Swine:

In swine after administration of 3 times the recommended dose or more a reduction in feeding, hydration and weight gain has been observed.

After administration of 5 times the recommended dose or more vomiting has also been noted.

WITHDRAWAL PERIODS

Meat and offal:

Cattle : IM use (20 mg/kg body weight, twice) : 30 days.

SC use (40 mg/kg body weight, once): 44 days.

Swine : 18 days.

Milk:

Not authorised for use in animals producing milk for human consumption including pregnant animals intended to produce milk for human consumption.

AMOUNT(S) TO BE ADMINISTERED AND ADMINISTRATION ROUTE

To ensure a correct dosage body weight should be determined as accurately as possible to avoid underdosing.

For treatment

Cattle:

Intramuscular use: 20 mg/kg body weight (1 ml/15 kg), twice at a 48-h interval.

Subcutaneous use: 40 mg/kg body weight (2 ml/15 kg), administered once only.

The dose should not exceed 10 ml per injection site.

The injection should only be given in the neck.

A large-bore injection needle (16-G) should be used.

Swine:

Intramuscular use: 15 mg/kg body weight (1 ml/20 kg), twice at a 48-h interval.

The dose should not exceed 3 ml per injection site.

The injection should only be given in the neck.

A large-bore injection needle (16-G) should be used.

For metaphylaxis

Cattle:

Subcutaneous use: 40 mg/kg body weight (2 ml/15 kg), administered once only.

The dose should not exceed 10 ml per injection site.

The injection should only be given in the neck.

A large-bore injection needle (16-G) should be used.

It is recommended to treat animals in the early stages of disease and to evaluate the response to treatment within 48 hours after the second injection. If clinical signs of respiratory disease persist 48 hours after the last injection, treatment should be changed using another formulation or another antibiotic and continued until clinical signs have resolved. The stopper of the injection bottle should be disinfected before each withdrawal. Use a dry, sterile cannula and syringe. Do not pierce the stopper more than 25 times.

STORAGE INSTRUCTIONS

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.

Store dark below 25 °C.

Shelf life after first opening the immediate packaging: 2 weeks.

Storage after first opening: Store dark and refrigerated (2 - 8 °C).

SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCTS OR WASTE MATERIALS

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

Keep out of reach of children / Jauhkan dari Kanak-Kanak

MARKETING AUTHORIZATION HOLDER

Danberg (M) Sdn Bhd (58493-A), Lot 1,
Jalan Bursa 23/4, Seksyen 23, 40300 Shahm Alam, Selangor,
Malaysia

Controlled Medicine / Ubat Terkawal

Reg. No.:



Manufactured by:

Interchemie werken De Adelaar Eesti AS, Vanapere tee 14,
Püünsi, Viimsi, Harju county 74013, Estonia

Holland