420x240mm   Font: Times New Roman - 7pt   45gsm with fold

VENUS® Enjoy Innovations

To be sold by retail on the prescription of an Oncologist/Cancer Hospital/ Institution only

PEMETREX-VENUS

PEMETREXED POWDER FOR CONCENTRATE FOR SOLUTION FOR INFUSION

A1. Product Name:
Pemetrex-venus (Pemetrexed Powder for concentrate for solution for infusion 500 mg)

A2. Name and Strength of Active Ingredient (s)
Active ingredient:
Pemetrexed Disodium
Strength: 500 mg

A3. Dosage Form:
Powder for concentrate for solution for infusion

A4. Product Description:
A White to light yellow or green yellow lyophilized powder.
After dilution/Recon.: A clear colorless solution.
List of compatible diluents: Sodium chloride 9 mg/ml (0.9 %) solution

A5. Pharmacodynamics:
ATC code: L01BA04
Pemetrexed (pemetrexed) is a multi-targeted anti-cancer antifolate agent that exerts its action by disrupting crucial folate-dependent metabolic processes essential for cell replication.
In vitro studies have shown that pemetrexed behaves as a multitargeted antifolate by inhibiting thymidylate synthase (TS), dihydrofolate reductase (DHFR) and glycinamide ribonucleotide formyltransferase (GARFT), which are key folate-dependent enzymes for the de novo biosynthesis of thymidine and purine nucleotides. Pemetrexed is transported into cells by both the reduced folate carrier and membrane folate binding protein transport systems. Once in the cell, pemetrexed is rapidly and efficiently converted to polyglutamate forms by the enzyme folylpolyglutamate synthetase. The polyglutamate forms are retained in cells and are even more potent inhibitors of TS and GARFT. Polyglutamation is a time- and concentration-dependent process that occurs in tumour cells and, to a lesser extent, in normal tissues. Polyglutamated metabolites have an increased intracellular half-life resulting in prolonged drug action in malignant cells.

A6. Pharmacokinetics:
The pharmacokinetic properties of pemetrexed following single-agent administration have been evaluated in 426 cancer patients with a variety of solid tumours at doses ranging from 0.2 to 838 mg/m^2 infused over a 10-minute period. Pemetrexed has a steady-state volume of distribution of 9 l/m^2. In vitro studies indicate that pemetrexed is approximately 81 % bound to plasma proteins. Binding was not notably affected by varying degrees of renal impairment. Pemetrexed undergoes limited hepatic metabolism. Pemetrexed is primarily eliminated in the urine, with 70 % to 90 % of the administered dose being recovered unchanged in urine within the first 24 hours following administration. In Vitro studies indicate that pemetrexed is actively secreted by OAT3 (organic anion transporter. Pemetrexed total systemic clearance is 91.8 ml/min and the elimination half-life from plasma is 3.5 hours in patients with normal renal function (creatinine clearance of 90 ml/min). Between patient variability in clearance is moderate at 19.3 %. Pemetrexed total systemic exposure (AUC) and maximum plasma concentration increase proportionally with dose. The pharmacokinetics of pemetrexed are consistent over multiple treatment cycles.
The pharmacokinetic properties of pemetrexed are not influenced by concurrently administered cisplatin. Oral folic acid and intramuscular vitamin B_{12} supplementation or N-acetyl-cysteine (NAC) injection do not affect the pharmacokinetics of pemetrexed.

A7. Indication:
Malignant pleural mesothelioma: Pemetrexed in combination with cisplatin is indicated for the treatment of chemotherapy naïve patients with unresectable malignant pleural mesothelioma.
Non-small cell lung cancer: Pemetrexed in combination with cisplatin is indicated for the first line treatment of patients with locally advanced or metastatic non-small cell lung cancer other than predominantly squamous cell histology. Pemetrexed is indicated as monotherapy for the maintenance treatment of locally advanced or metastatic non-small cell lung cancer other than predominantly squamous cell histology in patients whose disease has not progressed immediately following platinum-based chemotherapy.
Pemetrexed is indicated as monotherapy for the second line treatment of patients with locally advanced or metastatic non-small cell lung cancer other than predominantly squamous cell histology.
Pemetrexed in combination with pembrolizumab and platinum chemotherapy, is indicated for the first line treatment of patients with metastatic non-squamous NSCLC, with no EGFR or ALK genomic tumor aberrations.

A8. Recommended Dose:
Pemetrexed must only be administered under the supervision of a physician qualified in the use of anti-cancer chemotherapy.
Pemetrexed in combination with cisplatin: The recommended dose of Pemetrexed is 500 mg/m^2 of body surface area (BSA) administered as an intravenous infusion over 10 minutes on the first day of each 21-day cycle. The recommended dose of cisplatin is 75 mg/m^2 BSA infused over two hours approximately 30 minutes after completion of the pemetrexed infusion on the first day of each 21-day cycle. Patients must receive adequate anti-emetic treatment and appropriate hydration prior to and/or after receiving cisplatin (see also cisplatin product insert for specific dosing advice).
Pemetrexed as single agent: In patients treated for non-small cell lung cancer after prior chemotherapy, the recommended dose of Pemetrexed is 500 mg/m^2 BSA administered as an intravenous infusion over 10 minutes on the first day of each 21-day cycle.

Pemetrexed in combination with pembrolizumab: The recommended dose of Pemetrexed when administered with pembrolizumab and platinum chemotherapy for the initial treatment of metastatic non-squamous NSCLC in patients with a creatinine clearance (calculated by Cockcroft-Gault equation) of 45 mL/min or greater is 500 mg/m^2 as an intravenous infusion over 10 minutes administered after pembrolizumab and prior to carboplatin or cisplatin on Day 1 of each 21-day cycle for 4 cycles. Following completion of platinum-based therapy, treatment with Pemetrexed with or without pembrolizumab is administered until disease progression or unacceptable toxicity (see also pembrolizumab product insert for specific dosing advice).
Pre-medication regimen: To reduce the incidence and severity of skin reactions, a corticosteroid should be given the day prior to, on the day of, and the day after pemetrexed administration. The corticosteroid should be equivalent to 4 mg of dexamethasone administered orally twice a day.
To reduce toxicity, patients treated with pemetrexed must also receive vitamin supplementation. Patients must take oral folic acid or a multivitamin containing folic acid (350 to 1000 micrograms) on a daily basis. At least five doses of folic acid must be taken during the seven days preceding the first dose of pemetrexed, and dosing must continue during the full course of therapy and for 21 days after the last dose of pemetrexed. Patients must also receive an intramuscular injection of vitamin B12 (1000 micrograms) in the week preceding the first dose of pemetrexed and once every three cycles thereafter. Subsequent vitamin B12 injections may be given on the same day as pemetrexed.
Monitoring: Patients receiving pemetrexed should be monitored before each dose with a complete blood count, including a differential white cell count (WCC) and platelet count. Prior to each chemotherapy administration blood chemistry tests should be collected to evaluate renal and hepatic function. Before the start of any cycle of chemotherapy, patients are required to have the following: absolute neutrophil count (ANC) should be ≥ 1500 cells/mm^3 and platelets should be ≥ 100,000 cells/mm^3.
Creatinine clearance should be ≥ 45 ml/min.
The total bilirubin should be ≤ 1.5 times upper limit of normal. Alkaline phosphatase (AP), aspartate aminotransferase (AST or SGOT) and alanine aminotransferase (ALT or SGPT) should be ≤ 3 times upper limit of normal. Alkaline phosphatase, AST and ALT ≤ 5 times upper limit of normal is acceptable if liver has tumour involvement.
Dose adjustments: Dose adjustments at the start of a subsequent cycle should be based on nadir haematologic counts or maximum non-haematologic toxicity from the preceding cycle of therapy. Treatment may be delayed to allow sufficient time for recovery. Upon recovery patients should be retreated using the guidelines in Tables 1, 2 and 3, which are applicable for Pemetrexed used as a single agent or in combination with cisplatin.

Table 1 - Dose modification table for Pemetrexed (as single agent or in combination) and cisplatin-Haematologic toxicities

Nadir ANC < 500/mm^3 and nadir platelets ≥ 50,000/mm^3	75% of previous dose (both Pemetrexed and cisplatin)
Nadir platelets < 50,000/mm^3 regardless of nadir ANC	75% of previous dose (both Pemetrexed and cisplatin)
Nadir platelets < 50,000/mm^3 with bleedinga, regardless of nadir ANC	50% of previous dose (both Pemetrexed and cisplatin)

a. These criteria meet the National Cancer Institute Common Toxicity Criteria (CTC v2.0; NCI 1998) definition of ≥ CTC Grade 2 bleeding.
If patients develop non-haematologic toxicities ≥ Grade 3 (excluding neurotoxicity), Pemetrexed should be withheld until resolution to less than or equal to the patient's pre-therapy value. Treatment should be resumed according to the guidelines in Table 2.

Table 2. Dose Modification Table for Pemetrexed (as Single Agent or in Combination) and Cisplatin - Non-Haematologic Toxicities a, b

	Dose of Pemetrexed (mg/m^2)	Dose for Cisplatin (mg/m^2)
Any Grade 3 or 4 toxicities except mucositis	75% of previous dose	75% of previous dose
Any diarrhoea requiring hospitalisation (irrespective of grade) or Grade 3 or 4 diarrhoea	75% of previous dose	75% of previous dose
Grade 3 or 4 mucositis	50% of previous dose	100% of previous dose

a National Cancer Institute Common Toxicity Criteria (CTC v2.0; NCI 1998)
b Excluding neurotoxicity
In the event of neurotoxicity, the recommended dose adjustment for Pemetrexed and cisplatin is documented in Table 3. Patients should discontinue therapy if Grade 3 or 4 neurotoxicity is observed.

Table 3. Dose Modification Table for Pemetrexed (as Single Agent or in Combination) and Cisplatin - Neurotoxicity

CTC a Grade	Dose of Pemetrexed (mg/m^2)	Dose for Cisplatin (mg/m^2)
0-1	100% of previous dose	100% of previous dose
2	100% of previous dose	100% of previous dose

a National Cancer Institute Common Toxicity Criteria (CTC v2.0; NCI 1998)
Treatment with Pemetrexed should be discontinued if a patient experiences any haematologic or non-haematologic Grade 3 or 4 toxicity after 2 dose reductions or immediately if Grade 3 or 4 neurotoxicity is observed.

Special populations
Elderly: In clinical studies, there has been no indication that patients 65 years of age or older are at increased risk of adverse reaction compared to patients younger than 65 years old. No dose reductions other than those recommended for all patients are necessary.
Paediatric population: There is no relevant use of Pemetrexed in the paediatric population in malignant pleural mesothelioma and non-small cell lung cancer.
Patients with renal impairment (Standard Cockcroft and Gault formula or Glomerular Filtration Rate measured Tc99m-DPTA serum clearance method)
Pemetrexed is primarily eliminated unchanged by renal excretion. In clinical studies, patients with creatinine clearance of ≥ 45 ml/min required no dose adjustments other than those recommended for all patients. There are insufficient data on the use of pemetrexed in patients with creatinine clearance below 45 ml/min; therefore the use of pemetrexed is not recommended.

Patients with hepatic impairment: No relationships between AST (SGOT), ALT (SGPT), or total bilirubin and pemetrexed pharmacokinetics were identified. However patients with hepatic impairment such as bilirubin > 1.5 times the upper limit of normal and/or aminotransferase > 3.0 times the upper limit of normal (hepatic metastases absent) or > 5.0 times the upper limit of normal (hepatic metastases present) have not been specifically studied.

A9. Route of Administration:
For precautions to be taken before handling or administering Pemetrexed and for instructions on reconstitution and dilution of Pemetrexed before administration.

A10. Contraindication:
Hypersensitivity to the active substance or to any of the excipients.
Breast-feeding.
Concomitant yellow fever vaccine.

A 11. Warning and Precautions:
General: Pemetrexed can suppress bone marrow function as manifested by neutropenia, thrombocytopenia and anaemia (or pancytopenia). Myelosuppression is usually the dose-limiting toxicity. Patients should be monitored for myelosuppression during therapy and pemetrexed should not be given to patients until absolute neutrophil count (ANC) returns to ≥1500 cells/mm^3 and platelet count returns to ≥100,000 cells/mm^3. Dose reductions for subsequent cycles are based on nadir ANC, platelet count and maximum non-haematologic toxicity seen from the previous cycle.
Less toxicity and reduction in Grade 3/4 haematologic and non-haematologic toxicities such as neutropenia, febrile neutropenia and infection with Grade 3/4 neutropenia were reported when pre-treatment with folic acid and vitamin B_{12} was administered. Therefore, all patients treated with pemetrexed must be instructed to take folic acid and vitamin B_{12} as a prophylactic measure or N-acetyl-cysteine (NAC) to reduce treatment-related toxicity.
Skin reactions have been reported in patients not pre-treated with a corticosteroid. Pre-treatment with dexamethasone (or equivalent) can reduce the incidence and severity of skin reactions.
An insufficient number of patients has been studied with creatinine clearance of below 45 ml/min. Therefore, the use of pemetrexed in patients with creatinine clearance of < 45 ml/min is not recommended.
Patients with mild to moderate renal insufficiency (creatinine clearance from 45 to 79 ml/min) should avoid taking non-steroidal anti-inflammatory drugs (NSAIDs) such as ibuprofen, and acetylsalicylic acid (> 1.3 g daily) for 2 days before, on the day of, and 2 days following pemetrexed administration.
In patients with mild to moderate renal insufficiency eligible for pemetrexed therapy NSAIDs with long elimination half-lives should be interrupted for at least 5 days prior to, on the day of and at least 2 days following pemetrexed administration.
Serious renal events, including acute renal failure, have been reported with pemetrexed alone or in association with other chemotherapeutic agents. Many of the patients in whom these occurred had underlying risk factors for the development of renal events including dehydration or pre-existing hypertension or diabetes. Nephrogenic diabetes insipidus and renal tubular necrosis were also reported in post marketing setting with pemetrexed alone or with other chemotherapeutic agents. Most of these events resolved after pemetrexed withdrawal. Patients should be regularly monitored for acute tubular necrosis, decreased renal function and signs and symptoms of nephrogenic diabetes insipidus (e.g. hypernatraemia).
The effect of third space fluid, such as pleural effusion or ascites, on pemetrexed is not fully defined. A phase 2 study of pemetrexed in 31 solid tumour patients with stable third space fluid demonstrated no difference in pemetrexed dose normalized plasma concentrations or clearance compared to patients without third space fluid collections. Thus, drainage of third space fluid collection prior to pemetrexed treatment should be considered, but may not be necessary.
Due to the gastrointestinal toxicity of pemetrexed given in combination with cisplatin, severe dehydration has been observed. Therefore, patients should receive adequate antiemetic treatment and appropriate hydration prior to and/or after receiving treatment.
Serious cardiovascular events, including myocardial infarction and cerebrovascular events have been uncommonly reported during clinical studies with pemetrexed, usually when given in combination with another cytotoxic agent. Most of the patients in whom these events have been observed had pre-existing cardiovascular risk factors.
Immunodepressed status is common in cancer patients. As a result, concomitant use of live attenuated vaccines is not recommended.
Pemetrexed can have genetically damaging effects. Sexually mature males are advised not to father a child during the treatment and up to 6 months thereafter. Contraceptive measures or abstinence are recommended. Owing to the possibility of pemetrexed treatment causing irreversible infertility, men are advised to seek counselling on sperm storage before starting treatment.
Women of childbearing potential must use effective contraception during treatment with pemetrexed.
Cases of radiation pneumonitis have been reported in patients treated with radiation either prior, during or subsequent to their pemetrexed therapy. Particular attention should be paid to these patients and caution exercised with use of other radiosensitising agents. Cases of radiation recall have been reported in patients who received radiotherapy weeks or years previously.
Excipients: Pemetrex-Venus 500 mg powder for concentrate for solution for infusion. This medicinal product contains approximately 54 mg of sodium per vial. To be taken into consideration by patients on a controlled sodium diet.

A12. Interaction With Other Medicaments:
Pemetrexed is mainly eliminated unchanged renally by tubular secretion and to a lesser extent by glomerular filtration. Concomitant administration of nephrotoxic drugs (e.g. aminoglycoside, loop diuretics, platinum compounds, cyclosporin) could potentially result in delayed clearance of pemetrexed. This combination should be used with caution. If necessary, creatinine clearance should be closely monitored.
Concomitant administration of substances that are also tubularly secreted (e.g. probenecid, penicillin) could potentially result in delayed clearance of pemetrexed. Caution should be made when these drugs are combined with pemetrexed. If necessary, creatinine clearance should be closely monitored.
In patients with normal renal function (creatinine clearance ≥ 80 ml/min), high doses of non-steroidal anti-inflammatory drugs (NSAIDs, such as ibuprofen > 1600 mg/day) and acetylsalicylic acid at higher dose (≥ 1.3 g daily) may decrease pemetrexed elimination and, consequently, increase the occurrence of pemetrexed adverse events. Therefore, caution should be made when administering higher doses of NSAIDs or acetylsalicylic acid, concurrently with pemetrexed to patients with normal function (creatinine clearance ≥ 80 ml/min).

In patients with mild to moderate renal insufficiency (creatinine clearance from 45 to 79 ml/min), the concomitant administration of pemetrexed with NSAIDs (e.g. ibuprofen) or acetylsalicylic acid at higher dose should be avoided for 2 days before, on the day of, and 2 days following pemetrexed administration.
In the absence of data regarding potential interaction with NSAIDs having longer half-lives such as piroxicam or rofecoxib, the concomitant administration with pemetrexed in patients with mild to moderate renal insufficiency should be interrupted for at least 5 days prior to, on the day of, and at least 2 days following pemetrexed administration. If concomitant administration of NSAIDs is necessary, patients should be monitored closely for toxicity, especially myelosuppression and gastrointestinal toxicity.
Pemetrexed undergoes limited hepatic metabolism. Results from in vitro studies with human liver microsomes indicated that pemetrexed would not be predicted to cause clinically significant inhibition of the metabolic clearance of drugs metabolised by CYP3A, CYP2D6, CYP2C9 and CYP1A2.
Interactions common to all cytotoxics: Due to the increased thrombotic risk in patients with cancer, the use of anticoagulation treatment is frequent. The high intra-individual variability of the coagulation status during diseases and the possibility of interaction between oral anticoagulants and anticancer chemotherapy require increased frequency of INR (International Normalised Ratio) monitoring, if it is decided to treat the patient with oral anticoagulants.
Concomitant use contraindicated: Yellow fever vaccine: risk of fatal generalised vaccinale disease.
Concomitant use not recommended: Live attenuated vaccines (except yellow fever, for which concomitant use is contraindicated): risk of systemic, possibly fatal, disease. The risk is increased in subjects who are already immunosuppressed by their underlying disease. Use an inactivated vaccine where it exists (poliomyelitis).

A13. Pregnancy and Lactation
Contraception in males and females: Women of childbearing potential must use effective contraception during treatment with pemetrexed. Pemetrexed can have genetically damaging effects. Sexually mature males are advised not to father a child during the treatment and up to 6 months thereafter. Contraceptive measures or abstinence are recommended.
Pregnancy: There are no data from the use of pemetrexed in pregnant women but pemetrexed, like other anti-metabolites, is suspected to cause serious birth defects when administered during pregnancy. Animal studies have shown reproductive toxicity. Pemetrexed should not be used during pregnancy unless clearly necessary, after a careful consideration of the needs of the mother and the risk for the foetus.
Breast-feeding: It is not known whether pemetrexed is excreted in human milk and adverse reactions on the suckling child cannot be excluded. Breast-feeding must be discontinued during pemetrexed therapy.
Fertility: Owing to the possibility of pemetrexed treatment causing irreversible infertility, men are advised to seek counselling on sperm storage before starting treatment.

A14. Side Effects:
Summary of the safety profile: The most commonly reported undesirable effects related to pemetrexed, whether used as monotherapy or in combination, are bone marrow suppression manifested as anaemia, neutropenia, leukopenia, thrombocytopenia and gastrointestinal toxicities, manifested as anorexia, nausea, vomiting, diarrhoea, constipation, pharyngitis, mucositis, and stomatitis. Other undesirable effects include renal toxicities, increased aminotransferases, alopecia, fatigue, dehydration, rash, infection/sepsis and neuropathy. Rarely seen events include Stevens-Johnson syndrome and Toxic epidermal necrolysis.
Tabulated list of adverse reactions:
The table below provides the frequency and severity of undesirable effects that have been reported in > 5 % of 168 patients with mesothelioma who were randomised to receive cisplatin and pemetrexed and 163 patients with mesothelioma randomised to receive single agent cisplatin. In both treatment arms, these chemonaive patients were fully supplemented with folic acid and vitamin B_{12}.
Frequency estimate: Very common (≥1/10), Common (≥1/100 to <1/10), Uncommon (≥1/1000 to <1/100), Rare (≥1/10,000 to <1/1000), Very rare (<1/10,000) and not known (cannot be estimated from available data).
Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

System Organ Class	Frequency	Event*	Pemetrexed/ Cisplatin (N=168)		Cisplatin (N=163)	
			All grades toxicity (%)	Grade 3-4 toxicity (%)	All grades toxicity (%)	Grade 3-4 toxicity (%)
Blood and lymphatic system disorders	Very common	Neutrophils/ Granulocytes decreased	56.0	23.2	13.5	3.1
		Leukocytes decreased	53.0	14.9	16.6	0.6
		Haemoglobin decreased	26.2	4.2	10.4	0.0
		Platelets decreased	23.2	5.4	8.6	0.0
Metabolism and nutrition disorders	Common	Dehydration	6.5	4.2	0.6	0.6
Nervous system disorders	Very common	Neuropathy-Sensory	10.1	0.0	9.8	0.6
	Common	Taste disturbance	7.7	0.0***	6.1	0.0***
Eye disorders	Common	Conjunctivitis	5.4	0.0	0.6	0.0
Gastrointestinal disorders	Very common	Diarrhoea	16.7	3.6	8.0	0.0
		Vomiting	56.5	10.7	49.7	4.3
		Stomatitis/ Pharyngitis	23.2	3.0	6.1	0.0
		Nausea	82.1	11.9	76.7	5.5
		Anorexia	20.2	1.2	14.1	0.6
		Constipation	11.9	0.6	7.4	0.6
	Common	Dyspepsia	5.4	0.6	0.6	0.0
Skin and subcutaneous tissue disorders	Very common	Rash	16.1	0.6	4.9	0.0
		Alopecia	11.3	0.0***	5.5	0.0***
Renal and urinary disorders	Very common	Creatinine elevation	10.7	0.6	9.8	1.2
		Creatinine clearance decreased***	16.1	0.6	17.8	1.8
General disorders and administration site conditions	Very common	Fatigue	47.6	10.1	42.3	9.2

*Refer to National Cancer Institute CTC version 2 for each grade of toxicity except the term "creatinine clearance decreased"
** which is derived from the term "renal/genitourinary other".
*** According to National Cancer Institute CTC (v2.0; NCI 1998), taste disturbance and alopecia should only be reported as Grade 1 or 2.
For the purpose of this table a cut off of 5 % was used for inclusion of all events where the reporter considered a possible relationship to pemetrexed and cisplatin.
Clinically relevant CTC toxicities that were reported in ≥ 1 % and ≤ 5 % of the patients that were randomly assigned to receive cisplatin and pemetrexed include: renal failure, infection, pyrexia, febrile neutropenia, increased AST, ALT, and GGT, urticaria and chest pain.
Clinically relevant CTC toxicities that were reported in < 1 % of the patients that were randomly assigned to receive cisplatin and pemetrexed include arrhythmia and motor neuropathy.
The table below provides the frequency and severity of undesirable effects that have been reported in > 5 % of 265 patients randomly assigned to receive single agent pemetrexed with folic acid and vitamin B12 supplementation and 276 patients randomly assigned to receive single agent docetaxel. All patients were diagnosed with locally advanced or metastatic non-small cell lung cancer and received prior chemotherapy.

System Organ Class	Frequency	Event*	Pemetrexed (N=265)		Docetaxel (N=276)	
			All grades toxicity (%)	Grade 3-4 toxicity (%)	All grades toxicity (%)	Grade 3-4 toxicity (%)
Blood and lymphatic system disorders	Very common	Neutrophils/ Granulocytes decreased	12.1	4.2	34.1	27.2
		Leukocytes decreased	12.1	4.2	34.1	27.2
		Haemoglobin decreased	19.2	4.2	22.1	4.3
	Common	Platelets decreased	8.3	1.9	1.1	0.4
Gastrointestinal disorders	Very common	Diarrhoea	12.8	0.4	24.3	2.5
		Vomiting	16.2	1.5	12.0	1.1
		Stomatitis/ Pharyngitia	14.7	1.1	17.4	1.1
		Nausea	30.9	2.6	16.7	1.8
		Anorexia	21.9	1.9	23.9	2.5
	Common	Constipation	5.7	0.0	4.0	0.0
Hepatobiliary disorders	Common	SGPT (ALT) elevation	7.9	1.9	1.4	0.0
		SGOT (AST) elevation	6.8	1.1	0.7	0.0
Skin and subcutaneous tissue disorders	Very common	Rash/ Desquamation	14.0	0.0	1.8	0.0
	Common	Pruritus	6.8	0.4	1.8	2.5
		Alopecia	6.4	0.4***	37.7	2.2***
General disorders and administration site conditions	Very common	Fatigue	34.6	5.3	35.9	5.4
	Common	Fever	8.3	0.0	7.6	0.0

*Refer to National Cancer Institute CTC version 2 for each grade of toxicity.

**According to National Cancer Institute CTC (v2.0; NCI 1998), alopecia should only be reported as Grade 1 or 2.
For the purpose of this table a cut off of 5 % was used for inclusion of all events where the reporter considered a possible relationship to pemetrexed.
Clinically relevant CTC toxicities that were reported in ≥ 1 % and ≤ 5 % of the patients that were randomly assigned to pemetrexed include: infection without neutropenia, febrile neutropenia, allergic reaction/hypersensitivity, increased creatinine, motor neuropathy, sensory neuropathy, erythema multiforme, and abdominal pain.
Clinically relevant CTC toxicities that were reported in < 1 % of the patients that were randomly assigned to pemetrexed include supraventricular arrhythmias.
Clinically relevant Grade 3 and Grade 4 laboratory toxicities were similar between integrated Phase 2 results from three single agent pemetrexed studies (n=164) and the Phase 3 single agent pemetrexed study described above, with the exception of neutropenia (12.8 % versus 5.3 %, respectively) and alanine aminotransferase elevation (15.2 % versus 1.9 %, respectively). These differences were likely due to differences in the patient population, since the Phase 2 studies included both chemonaive and heavily pre-treated breast cancer patients with pre-existing liver metastases and/or abnormal baseline liver function tests.
The table below provides the frequency and severity of undesirable effects considered possibly related to study drug that have been reported in >5% of 839 patients with NSCLC who were randomized to receive cisplatin and pemetrexed and 830 patients with NSCLC who were randomized to receive cisplatin and gemcitabine. All patients received study therapy as initial treatment for locally advanced or metastatic NSCLC and patients in both treatment groups were fully supplemented with folic acid and vitamin B12.

System Organ Class	Frequency	Event*	Pemetrexed/ Cisplatin (N=839)		Gemcitabine/ Cisplatin (N=830)	
			All grades toxicity (%)	Grade 3-4 toxicity (%)	All grades toxicity (%)	Grade 3-4 toxicity (%)
Blood and lymphatic system disorders	Very common	Haemoglobin decreased	33.0*	5.6*	45.7*	9.9*
		Neutrophils/ Granulocytes decreased	29.0*	15.1*	38.4*	26.7*
		Leukocytes decreased	17.8	4.8*	20.6	7.6*
		Platelets decreased	10.1*	4.1*	26.6*	12.7*
Nervous system disorders	Common	Neuropathy-Sensory	8.5*	0.0*	12.4*	0.6*
		Taste disturbance	8.1	0.0***	8.9	0.0***
Gastrointestinal disorders	Very common	Nausea	56.1	7.2	53.4	3.9*
		Vomiting	39.7	6.1	35.5	6.1
		Anorexia	26.6	2.4*	24.2	0.7*
		Constipation	21.0	0.8	19.5	0.4
		Stomatitis/ Pharyngitis	13.5	0.8	12.4	0.1
		Diarrhoea without colostomy	12.4	1.3	12.8	1.6
	Common	Dyspepsia/ Heartburn	5.2	0.1	5.9	0.0
Skin and subcutaneous tissue disorders	Very common	Alopecia	11.9*	0***	21.4*	0.5***
	Common	Rash/ Desquamation	6.6	0.1	8.0	0.5
Renal and urinary disorders	Very common	Creatinine elevation	10.1*	0.8	6.9*	0.5
General disorders and administration site conditions	Very common	Fatigue	42.7	6.7	44.9	4.9

*P-values <0.05 comparing pemetrexed/cisplatin to gemcitabine/cisplatin, using Fisher Exact test.
**Refer to National Cancer Institute CTC (v2.0; NCI 1998) for each Grade of Toxicity.
***According to National Cancer Institute CTC (v2.0; NCI 1998), taste disturbance and alopecia should only be reported as Grade 1 or 2.
For the purpose of this table, a cut-off of 5% was used for inclusion of all events where the reporter considered a possible relationship to pemetrexed and cisplatin.
Clinically relevant toxicity that was reported in ≥ 1% and ≤ 5% of the patients that were randomly assigned to receive cisplatin and pemetrexed include: AST increase, ALT increase, infection, febrile neutropenia, renal failure, pyrexia, dehydration, conjunctivitis, and creatinine clearance decrease.
Clinically relevant toxicity that was reported in < 1% of the patients that were randomly assigned to receive cisplatin and pemetrexed include: GGT increase, chest pain, arrhythmia, and motor neuropathy.
Clinically relevant toxicities with respect to gender were similar to the overall population in patients receiving pemetrexed plus cisplatin.
The table below provides the frequency and severity of undesirable effects considered possibly related to study drug that have been reported in > 5% of 800 patients randomly assigned to receive single agent pemetrexed and 402 patients randomly assigned to receive placebo in the single-agent pemetrexed maintenance (JMEN: N= 663) and continuation pemetrexed maintenance (PARAMOUNT: N=539) studies. All patients were diagnosed with Stage IIIB or IV NSCLC and had received prior platinum-based chemotherapy. Patients in both study arms were fully supplemented with folic acid and vitamin B12.

System Organ Class	Frequency	Event*	Pemetrexed*** (N=800)		Placebo*** (N=402)	
			All grades toxicity (%)	Grade 3-4 toxicity (%)	All grades toxicity (%)	Grade 3-4 toxicity (%)
Blood and lymphatic system disorders	Very common	Hemoglobin decreased	18.0	4.5	5.2	0.5
	Common	Leukocytes decreased	5.8	1.9	0.7	0.2
		Neutrophils decreased	8.4	4.4	0.2	0.0
Nervous system disorders	Common	Neuropathy-Sensory	7.4	0.6	5.0	0.2
Gastrointestinal disorders	Very common	Nausea	17.3	0.8	4.0	0.2
		Anorexia	12.8	1.1	3.2	0.0
	Common	Vomiting	8.4	0.3	1.5	0.0
		Mucositis/ Stomatitis	6.8	0.8	1.7	0.0
Hepatobiliary disorders	Common	ALT (SGPT) elevation	6.5	0.1	2.2	0.0
		AST (SGOT) elevation	5.9	0.0	1.7	0.0
Skin and subcutaneous tissue disorders	Common	Rash/ Desquamation	8.1	0.1	3.7	0.0
General disorders and administration site conditions	Very common	Fatigue	24.1	5.3	10.9	0.7
	Common	Pain	7.6	0.9	4.5	0.0
		Edema	5.6	0.0	1.5	0.0
Renal Disorders	Common	Renal disorders****	7.6	0.9	1.7	0.0

Abbreviations: ALT = alanine aminotransferase; AST = aspartate aminotransferase; CTCAE = Common Terminology Criteria for Adverse Event; NCI = National Cancer Institute; SGOT = serum glutamic oxaloacectic aminotransferase; SGPT = serum glutamic pyruvic aminotransferase.
* Definition of frequency terms: Very common - ≥ 10%; Common - > 5% and < 10%. For the purpose of this table, a cutoff of 5% was used for inclusion of all events where the reporter considered a possible relationship to pemetrexed.
** Refer to NCI CTCAE Criteria (Version 3.0; NCI 2003) for each grade of toxicity. The reporting rates shown are according to CTCAE version 3.0.
*** Integrated adverse reactions table combines the results of the JMEN pemetrexed maintenance (N=663) and PARAMOUNT continuation pemetrexed maintenance (N=539) studies.
**** Combined term includes increased serum/blood creatinine, decreased glomerular filtration rate, renal failure and renal/genitourinary- other.
Clinically relevant CTC toxicity of any grade that was reported in ≥ 1% and ≤ 5% of the patients that were randomly assigned to pemetrexed include: febrile neutropenia, infection, decreased platelets, diarrhoea, constipation, alopecia, pruritis/itching, fever (in the absence of neutropenia), ocular surface disease (including conjunctivitis), increased lacrimation, dizziness and motor neuropathy.
Clinically relevant CTC toxicity that was reported in < 1% of the patients that were randomly assigned to pemetrexed include: allergic reaction/hypersensitivity, erythema multiforme, supraventricular arrhythmia and pulmonary embolism.
Safety was assessed for patients who were randomised to receive pemetrexed (N=800). The incidence of adverse reactions was evaluated for patients who received ≤ 6 cycles of pemetrexed maintenance (N=519), and compared to patients who received > 6 cycles of pemetrexed (N=281). Increases in adverse reactions (all grades) were observed with longer exposure. A significant increase in the incidence of possibly study-drug-related Grade 3/4 neutropenia was observed with longer exposure to pemetrexed (≤6 cycles: 3.3%, > 6cycles: 6.4%: p=0.046). No statistically significant differences in any other individual Grade 3/4/5 adverse reactions were seen with longer exposure.
Serious cardiovascular and cerebrovascular events, including myocardial infarction, angina pectoris, cerebrovascular accident and transient ischaemic attack have been uncommonly reported during clinical studies with pemetrexed, usually when given in combination with another cytotoxic agent. Most of the patients in whom these events have been observed had pre-existing cardiovascular risk factors.
Rare cases of hepatitis, potentially serious, have been reported during clinical studies with pemetrexed. Pancytopenia has been uncommonly reported during clinical trials with pemetrexed.
In clinical trials, cases of colitis (including intestinal and rectal bleeding, sometimes fatal, intestinal perforation, intestinal necrosis and typhlitis) have been reported uncommonly in patients treated with pemetrexed. In clinical trials, cases of interstitial pneumonitis with respiratory insufficiency, sometimes fatal, have been reported uncommonly in patients treated with pemetrexed.
Uncommon cases of oedema have been reported in patients treated with pemetrexed. Oesophagitis/ radiation oesophagitis has been uncommonly reported during clinical trials with pemetrexed.
Sepsis, sometimes fatal, has been commonly reported during clinical trials with pemetrexed.
During post marketing surveillance, the following adverse reactions have been reported in patients treated with pemetrexed:

Hyperpigmentation has been commonly reported.
Uncommon cases of acute renal failure have been reported with pemetrexed alone or in association with other chemotherapeutic agents. Nephrogenic diabetes insipidus and renal tubular necrosis have been reported in post marketing setting with an unknown frequency.
Uncommon cases of radiation pneumonitis have been reported in patients treated with radiation either prior, during or subsequent to their pemetrexed therapy.
Rare cases of radiation recall have been reported in patients who have received radiotherapy previously.
Uncommon cases of peripheral ischaemia leading sometimes to extremity necrosis have been reported.
Rare cases of bullous conditions have been reported including Stevens-Johnson syndrome and Toxic epidermal necrolysis which in some cases were fatal. Rarely, immune-mediated haemolytic anaemia has been reported in patients treated with pemetrexed. Rare cases of anaphylactic shock have been reported.
Erythematous oedema mainly of the lower limbs has been reported with an unknown frequency. Infectious and non-infectious disorders of the dermis, the hypodermis and/or the subcutaneous tissue have been reported with an unknown frequency (e.g. acute bacterial dermo-hypodermitis, pseudocellulitis, dermatitis).

A15. Symptoms and Treatment of overdose
Reported symptoms of overdose include neutropenia, anaemia, thrombocytopenia, mucositis, sensory polyneuropathy and rash. Anticipated complications of overdose include bone marrow suppression as manifested by neutropenia, thrombocytopenia and anaemia. In addition, infection with or without fever, diarrhoea, and/or mucositis may be seen. In the event of suspected overdose, patients should be monitored with blood counts and should receive supportive therapy as necessary. The use of calcium folinate / folinic acid in the management of pemetrexed overdose should be considered.

A16. Effect on Ability to Drive and Use Machine
No studies on the effects on the ability to drive and use machines have been performed. However, it has been reported that pemetrexed may cause fatigue.

A17. Preclinical Safety Data
Not Applicable

A18. Instructions For Use
1. Use aseptic technique during the reconstitution and further dilution of pemetrexed for intravenous infusion administration.
2. Calculate the dose and the number of Pemetrexed vials needed. Each vial contains an excess of pemetrexed to facilitate delivery of label amount.
3. Pemetrexed 500 mg
Reconstitute 500-mg vials with 20 ml of sodium chloride 9 mg/ml (0.9 %) solution for injection without preservative, resulting in a solution containing 25 mg/ml pemetrexed.
Gently swirl each vial until the powder is completely dissolved. The resulting solution is clear and ranges in colour from colourless to yellow or green-yellow without adversely affecting product quality. The pH of the reconstituted solution is between 6.6 and 7.8. Further dilution is required.
4. The appropriate volume of reconstituted pemetrexed solution must be further diluted to 100 ml with sodium chloride 9 mg/ml (0.9 %) solution for injection without preservative, and administered as an intravenous infusion over 10 minutes.
5. Pemetrexed infusion solutions prepared as directed above are compatible with polyvinyl chloride and polyolefin lined administration sets and infusion bags.
6. Parenteral medicinal products must be inspected visually for particulate matter and discolouration prior to administration. If particulate matter is observed, do not administer.
7. Pemetrexed solutions are for single use only. Any unused medicinal product or waste material must be disposed of in accordance with local requirements.
Preparation and administration precautions: As with other potentially toxic anticancer agents, care should be exercised in the handling and preparation of pemetrexed infusion solutions. The use of gloves is recommended. If a pemetrexed solution contacts the skin, wash the skin immediately and thoroughly with soap and water. If pemetrexed solutions contact the mucous membranes, flush thoroughly with water. Pemetrexed is not a vesicant. There is not a specific antidote for extravasation of pemetrexed. There have been few reported cases of pemetrexed extravasation, which were not assessed as serious by the investigator. Extravasation should be managed by local standard practice as with other non-vesicants.

A19. Storage Condition
Store below 30°C.
Unopened vial: This medicinal product does not require any special storage conditions.
Reconstituted and infusion solutions: When prepared as directed, reconstituted and infusion solutions of Pemetrex-venus contain no antimicrobial preservatives. Chemical and physical in-use stability of reconstituted and infusion solutions of Pemetrex-venus were demonstrated for 24 hours at refrigerated temperature. From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would not be longer than 24 hours at 2°C to 8°C.

A20. Shelf Life
Unopened Vial : 36 Months

A21. Therapeutical Code (If Any)
Pharmacotherapeutic group: Folic acid analogues
ATC code: L01BA04

Manufactured by: **VENUS REMEDIES LIMITED**
Hill Top Industrial Estate, Jharmajri, EPIP
Phase-I (Extn.), Bhatoli Kalan, Baddi,
Distt. Solan, Himachal Pradesh-173205, India

Product Registration Holder:
Healol Pharmaceuticals Sdn. Bhd.
74-3, Jalan Wangsa Delima 6, KLSC
Wangsa Maju, 53300, Kuala Lumpur,
Wilayah Persekutuan

Date of Revision : 08-01-2024

B81/PM500/4956-01