
UNIMED'S OXI-CLIN

(Hydrogen Peroxide 6% w/v solution)

Name and Strength of Active Substances

Hydrogen peroxide 6% w/v

Product Description

A clear, colorless liquid.

Pharmacodynamics

Hydrogen peroxide is used as a disinfectant and deodorant. It releases oxygen when applied to tissues, the effect lasts only as long as the oxygen is being released and is of short duration. The antimicrobial effect of the liberated oxygen is reduced in the presence of organic matter. It is used to cleanse wounds and ulcers in concentrations of up to 6%. Adhering and blood-soaked dressings may be released by the application of a solution of hydrogen peroxide. A 1.5% solution has been used as a mouthwash in the treatment of acute stomatitis and as a deodorant gargle.

Pharmacokinetics

No further information available.

Indications

UNIMED'S OXI-CLIN is an antiseptic for minor cuts or oral rinse.

Recommended Dose

Antiseptic

Dilute one part of UNIMED'S OXI-CLIN with one part of water. Soak cotton wool in the solution and dress the wound. Apply 2-3 times daily or when necessary.

Oral rinse

Dilute one part of UNIMED'S OXI-CLIN with five parts waters. Gargle for one minute and expel. Use 2-3 times daily.

Route of Administration

For topical or oromucosal use only.

Contraindications

Hypersensitivity to any of the ingredients.

Not for use in closed body cavities or on surgical wounds due to the risk of oxygen released into the circulation causing gas embolism.

Not for use as a disinfection agent for surgical instruments (particularly endoscopes) and as an enema.

Warning and Precautions

For external use only.

Keep all medicines away from children.

Not for use in closed body cavities or on surgical wounds due to risk of oxygen released into the circulation causing gas embolism.

Avoid normal skin.

Bleaches fabric.

Interactions with Other Medicaments

None known

Pregnancy & Lactation

Should be avoided if possible during pregnancy and lactation. No evidence is available as to the safety of use of this product in these conditions.

Adverse Effects / Undesirable Effects

Cases of gas embolism, sometimes resulting in cardiac arrest, have been reported when hydrogen peroxide has been instilled in closed body cavities or deep surgical wounds.

Strong solutions of hydrogen peroxide produce irritating burns on the skin and mucous membranes with a white eschar. The pain disappears after about 1 hour. Continued use of the product as a mouthwash may cause reversible hypertrophy of the papillae of the tongue.

Symptoms and Treatment of Overdose

Accidental ingestion may cause sore throat, gastric disturbances and vomiting. Sudden evolution of oxygen may cause injury by acute distension of the stomach and internal bleeding. Water may be given to drink. Ingestion of large volumes can lead to gas embolism following evolution of oxygen in the stomach.

Effects on ability to drive and use machines

None known.

Storage Conditions

Store below 30°C. Protect from light.

Dosage Forms and Packaging Available

60ml, 450ml, 500ml & 1L in white HDPE opaque bottle with red HDPE screw cap.

Registration Number:

MALXXXXX

Manufactured by & Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.
Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia.

Date of Revision:

02 February 2023