

GLEPTOFORT 200MG/ML INJECTION

Description and Composition

GLEPTOFORT 200MG/ML INJECTION is a dark brown solution. Each ml contains 200mg Iron (as gleptoferron) as its active ingredient and 5mg Phenol as its preservative.

Pharmacodynamics

Iron is an essential micronutrient. It takes a major role in the oxygen transport of haemoglobin and myoglobin, as well as it has a key role in enzymes, such as cytochromes, catalases, and peroxidases. Iron has a high recovery rate from metabolism and food ingested. Thus, deficiency occurs only very rarely in adult animals.

Pharmacokinetics

After intramuscular injection, the iron complex is absorbed into the lymphatic tissue within 3 days. Here, the complex is split to release Fe^{3+} which is stored as ferritin in the main storage organs (e.g. liver, spleen and the reticuloendothelial system). In the blood, free Fe^{3+} binds to transferrin (transport form) and is mainly used for the synthesis of haemoglobin.

Indication

For prevention and treatment of iron deficiency anaemia in piglets.

Recommended Dose

For strictly intramuscular injection.

Piglets: 1ml of the product which is equivalent to 200mg Iron (III) per animal. Inject once between the 1st and the 3rd day of life.

Route of Administration

To be given by intramuscular route in pigs.

Contraindication

Do not administer to piglets suspected to suffer from deficiency of vitamin E and/or selenium. Do not use in cases of hypersensitivity to the active substance or to any of the excipients. Do not use in clinically diseased animals, especially not in case of diarrhoea.

Warning and Precautions

Special warnings for each target species:

None known.

Special precautions for use

Special precautions for use in animals:

Not applicable

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known hypersensitivity to iron dextran, or those with haemochromatosis should avoid contact with the product.

Care should be taken to avoid accidental self-injection, as well as contact with the eyes and mouth.

In case of accidental injection, seek medical advice immediately and show the package leaflet or label to the physician.

Wash hands after use.

Interaction with Other Medicaments

The absorption of concomitantly administered oral iron may be reduced.

See also section Incompatibilities.

Pregnancy and Lactation

Not applicable.

Side Effects

Uncommonly discolouration of the tissue and/or slight, soft swelling may be observed at the site of injection. This should disappear within a few days. Also hypersensitivity reactions can occur.

Rarely deaths have occurred in piglets following the administration of parenteral iron dextran preparations. These deaths have been associated with genetic factors or deficiency of vitamin E and/or selenium.

Very rarely piglets deaths have been reported which have been attributed to an increased susceptibility to infection due to temporary blocking of the reticuloendothelial system.

The frequency of adverse reactions is defined using the following convention:

- very common (more than 1 in 10 animals treated displaying adverse reaction(s) during the course of one treatment)
- common (more than 1 but less than 10 animals in 100 animals treated)
- uncommon (more than 1 but less than 10 animals in 1,000 animals treated)
- rare (more than 1 but less than 10 animals in 10,000 animals treated)
- very rare (less than 1 animal in 10,000 animals treated, including isolated reports).

Symptoms and Treatment of Overdose

Transferrin-iron saturation levels leading to increased susceptibility for (systemic) bacterial disease, pain, inflammation reactions as well as abscess formation at the injection site may occur.

Persistent discolouration of muscle tissue at the injection site may occur.

Iatrogenic poisoning with following symptoms: pale mucous membranes, hemorrhagic gastroenteritis, vomiting, tachycardia, hypotension, dyspnoea, edema of the limbs, lameness, shock, death, liver damage. Supportive measures such as chelating agents can be used.

Incompatibilities

In the absence of compatibility studies, this veterinary medicinal product must not be mixed with other veterinary medicinal products.

Storage Condition

Store below 30°C. Protect from light.

Shelf life

3 years.

Following withdrawal of first dose, use the product within 24 hours. Any unused materials should be discarded.

Withdrawal Period

Pigs: Meat and offal: Zero days.

Packing

20ml, 50ml, 100ml, 250ml

Revised date: 1.11.2022

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com