

MARBOXIN 2% w/v INJECTION

Description and Composition

MARBOXIN 2% w/v INJECTION is a light buff to amber solution. Each ml contains 20mg of Marbofloxacin as its active ingredient and 10mg Benzyl Alcohol as its preservative.

Pharmacodynamics

Marbofloxacin is a synthetic, bactericidal antimicrobial, belonging to the fluoroquinolone group which acts by inhibition of DNA gyrase. It is effective against a wide range of Gram positive bacteria (in particular *Staphylococci*) and Gram negative bacteria (*Escherichia coli*, *Salmonella typhimurium*, *Campylobacter jejunii*, *Citrobacter*, *Enterobacter*, *Proteus* spp, *Klebsiella* spp, *Actinobacillus pleuropneumoniae*, *Bordetella bronchiseptica*, *Pasteurella haemolytica*, *Pasteurella multocida*, *Haemophilus* spp, *Moraxella* spp, *Pseudomonas aeruginosa*) as well as Mycoplasma (*Mycoplasma bovis*, *Mycoplasma dispar*, *Mycoplasma hyopneumoniae*).

Resistance to *Streptococcus* spp. may occur.

Pharmacokinetics

After subcutaneous administration in cattle and pigs at the recommended dose of 2mg/kg, marbofloxacin is readily absorbed and its bioavailability is close to 100%. It is weakly bound to plasma proteins (less than 10% in pigs and 30% in cattle), extensively distributed and in most tissues (liver, kidney, skin, lung, bladder, uterus digestive tract) it achieves higher concentrations than in plasma.

In cattle, marbofloxacin is eliminated slowly in pre-ruminating calves ($t_{1/2\beta}$ = 5-9 h) predominantly in the active form in urine (3/4) and faeces (1/4).

In pigs, marbofloxacin is eliminated slowly ($t_{1/2\beta}$ = 8-10 h) predominantly in the active form in urine (2/3) and faeces (1/3).

Indication

Indicated in the treatment of respiratory infections caused by susceptible strains of organisms in cattle and pigs.

Recommended Dose

The recommended dosage is 2mg per kg per day (1ml Marboxin 2% w/v per 10kg) in a single daily injection by subcutaneous or intravenous routes in cattle and by intramuscular route in pigs.

Treatment duration is 3 days by subcutaneous route and 3 to 5 days by intravenous route in cattle and 3 to 5 days by intramuscular route in pigs. The volume of injection should be limited to 10ml at each site of injection for pigs.

In order to reduce the risk of particulate contamination of the product, it is recommended that a draw-off needle be used to reduce the number of times the septum is punctured.

Route of Administration

To be given by subcutaneous or intravenous injection in cattle.

To be given by intramuscular injection in pigs.

Contraindications

None known.

Warning and Precautions

Special warnings for each target species:

None.

Special precautions for use

Special precautions for use in animals:

Official and local antimicrobial policies should be taken into account when the product is used. Fluoroquinolones should be reserved for the treatment of clinical conditions which have responded poorly,

or are expected to respond poorly, to other classes of antimicrobials. Whenever possible, fluoroquinolones should only be used based on susceptibility testing. Use of the product deviating from the instructions may increase the prevalence of bacteria resistant to the fluoroquinolones and may decrease the effectiveness of treatment with other quinolones due to the potential for cross resistance.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known hypersensitivity to fluoroquinolones should avoid using this product.

Wash hands after use.

Interaction with Other Medicaments

None known.

Pregnancy and Lactation

Not applicable.

Side Effects

No severe side-effects are to be expected at doses up to 5 times the recommended dose in cattle and pigs. In particular no lesions of the articular joints are encountered.

Subcutaneous injection is well tolerated. Transitory inflammatory reactions are sometimes observed at the injection site, but without clinical impact.

Symptoms and Treatment of Overdose

Overdosage may cause acute signs in the form of neurological disorders which should be treated symptomatically.

Storage Condition

Store below 30°C. Protect from light.

Shelf life

3 years.

Following withdrawal of first dose, use the product within 24 hours. Any unused material should be discarded.

Withdrawal Periods

Meat: Pre-ruminating calves (up to 100kg body weight): 6 days

Pigs: 4 days

Packing

20ml, 50ml, 100ml, 250ml.

Revised date: 9.9.22

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com