

AVIACOX 2.5% w/v SOLUTION

Description and Composition

AVIACOX 2.5% w/v SOLUTION is a clear colorless to yellow brown solution. Each ml contains 25mg Toltrazuril as its active ingredient.

Pharmacodynamics

Toltrazuril is an anticoccidial of the triazinetrione group, active against *Eimeria* spp. Toltrazuril induces changes in the fine structure of the developmental stages of coccidia. These are caused primarily by swelling of the endoplasmic reticulum and of the Golgi apparatus, abnormal changes to the perinuclear space and disturbances in cell division. Toltrazuril causes a decrease in the activity of respiratory chain enzymes in the parasites.

Pharmacokinetics

After oral administration, toltrazuril undergoes at least 50% absorption in poultry. The highest concentrations are to be found in the liver and kidneys of the poultry. The active substance is broken down rapidly. The main metabolite is toltrazuril sulfone.

Indication

For the treatment of coccidiosis in chickens and turkeys, caused by infections with various species of *Eimeria*:

Chickens: *E. acervulina*, *E. brunetti*, *E. maxima*, *E. mitis*, *E. necatrix*, *E. tenella*.

Turkeys: *E. adenoides* and *E. meleagrimitis*.

Recommended Dose

For oral administration via the drinking water.

Chickens and turkeys: The dose is 7mg toltrazuril per kg body weight per day (equivalent to 0.28ml of Aviacox 2.5% w/v Solution per kg body weight per day). Treatment is carried out on 2 consecutive days.

In case an automatic dose dispenser is used the medicinal product should be administered for one period of 8 hours per day for 2 consecutive days. Medicated drinking water should be refreshed every 24 hours.

In the case of continuous treatment over 24 hours, the volume of the product to be mixed into the drinking water for the animals to be treated is calculated according to the following formula:

Volume of the product required per litre drinking water:

$$\frac{0.28\text{ml Aviacox 2.5\% w/v Solution per kg bw per day}}{\text{Average drinking water intake in litres over 24 hours per animal}} \times \frac{\text{Average bw (kg) of the animals to be treated}}{1} = \frac{\text{ } \times \text{ml Aviacox 2.5\% w/v Solution per litre drinking water}}{1}$$

Total volume of the product required per day (24 h):

The calculated volume (x ml [product] per litre) must be multiplied by the total consumption of drinking water (1) per day (24 h).

In the case of treatment for a period of 8 hours per day, the volume of the product to be mixed into the drinking water for the animals to be treated is calculated according to the following formula:

Volume of the product required per litre drinking water:

$$\frac{0.28\text{ml Aviacox 2.5\% w/v Solution per kg bw per day}}{\text{Average drinking water intake in litres over 8 hours per animal}} \times \frac{\text{Average bw (kg) of the animals to be treated}}{1} = \frac{\text{ } \times \text{ml Aviacox 2.5\% w/v Solution per litre drinking water}}{1}$$

The volume of the product required for a treatment period of 8 hours:

The calculated volume (y ml [product] per litre) must be multiplied by the total consumption of drinking water (l) per 8-hour period.

The appropriate volume of solution must be added daily to the drinking water while stirring.

If volumes of between 1 and 4ml product are added per litre drinking water, solubility is guaranteed over the period of treatment.

In order to ensure that all the animals drink water evenly, sufficient space must be made available at the waterer. Free-range animals must be kept indoors during treatment.

After the end of the treatment, the watering system must be cleaned in an appropriate manner in order to prevent any exposure to residual subtherapeutic doses, particularly if liable to promote the development of resistance.

Predilution and the administration through a dosing pump (proportioner) are not recommended. Use preferably a bulk tank.

Route of Administration

For oral administration via the drinking water.

Contraindications

Do not use in animals with hypersensitivity to the active substance or to any of the excipients.

Warning and Precautions

Special warnings for each target species:

Good hygiene can reduce the risk of coccidiosis. It is therefore recommended any deficiencies in husbandry should be addressed in addition to treatment.

Poultry houses should be kept clean and dry.

It is recommended to treat all animals in a pen. For best results, treatment should be initiated before the clinical signs of disease have spread throughout the whole group.

Special precautions for use

Special precautions for use in animals:

As with any anticoccidials, frequent and prolonged use of an antiprotozoal of the same class may result in the development of resistances. It is important to keep to the recommended dose in order to minimise the risk of resistance.

If resistance is present it should be considered to use another antiprotozoal from another class/mechanism of action.

This veterinary medicinal product should not be used together with feed additives/or other veterinary medicinal products that might interfere with the efficacy of the product, like "coccidiostats" and "histomonostats".

For best results, treatment should be initiated before the clinical signs of disease have spread throughout the whole group.

The veterinary medicinal product is a strongly alkaline solution and should not be administered undiluted.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known hypersensitivity to toltrazuril should avoid contact with the veterinary medicinal product. The veterinary medicinal product is an alkaline solution.

Wear synthetic rubber gloves when handling the product.

Contact with skin and mucous membranes should be avoided.

In case of direct contact with the eyes or skin, wash immediately and thoroughly with water.

Do not ingest. In case of accidental ingestion, seek medical advice immediately and show the package leaflet or the label to the physician.

Do not eat, drink or smoke during use.

Wash hands after use.

Interactions with Other Medicaments

Combination of the product with antibiotics may result in reduced water intake in turkeys. The concomitant administration of other substances to the drinking water should be avoided.

Pregnancy and Lactation

Not applicable, see Withdrawal periods.

Side Effects

None known.

Symptoms and Treatment of Overdose

A reduction in drinking water intake may be the first sign of an overdose. This is observed only after an overdose with more than 10 times the recommended dose.

Storage Condition

Store below 30°C. Protect from light.

Shelf life

3 years.

To be used within 24 hours after dilution.

After first opening the container, any product remaining after 24 hours should be discarded.

Withdrawal Periods

Chickens and turkeys:

Meat and offal: 16 days

Eggs: Not authorised for use in laying birds producing eggs for human consumption.

Do not use within 4 weeks of onset of laying.

Packing

100ml, 200ml, 500ml, 1L.

Revised date: 23.2.2023

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com