

PIGICOX 5% SUSPENSION

Description and Composition

PIGICOX 5% SUSPENSION is an opaque white or yellowish suspension. Each ml contains 50mg Toltrazuril as its active ingredient and 2mg Sodium Benzoate as its preservative.

Pharmacodynamics

Toltrazuril is a triazinon derivative. It acts against coccidia of the genus *Cystoisospora* and *Eimeria*. It is active against all intracellular development stages of coccidia of the merogony (asexual multiplication) and gamogony (sexual phase). All stages are destroyed, thus the mode of action is coccidiocidal.

Pharmacokinetics

Pigs: After oral administration toltrazuril is slowly absorbed with a bioavailability of $\geq 70\%$. The main metabolite is characterised as toltrazuril sulfone. The elimination of toltrazuril is slow with a half-life elimination time around 3 days. The major route of excretion is via the faeces.

Indication

For the prevention of clinical signs of coccidiosis in neonatal piglets (3-5 days old) on farms with a confirmed history of coccidiosis caused by *Cystoisospora suis*.

Recommended Dose

For oral administration. Shake well before use.

Piglets: 20mg Toltrazuril per kg body weight per animal corresponding to 0.4ml of product per kg body weight on day 3 to 5 of life.

1 pump delivers 1ml of Pigicox 5% Suspension.

Route of Administration

For oral administration.

Contraindications

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

Warning and Precautions

Special warnings for each target species:

It is recommended to treat all animals in a pen.

Hygienic measures may reduce the risk of coccidiosis. It is therefore, recommended to improve concomitantly the hygienic conditions in the concerned facility, particularly dryness and cleanliness.

To obtain maximum benefit, animals should be treated before the expected onset of clinical signs, i.e. in the prepatent period.

To alter the course of an established clinical coccidial infection, in individual animals already showing signs of diarrhoea, additional supportive therapy may be required.

Treatment during an outbreak will be of limited value for the individual animal because of damage to the small intestine having already occurred.

As with any antiparasiticide frequent and repeated use of antiprotozoals from the same class may lead to the development of resistance.

If resistance is present it should be considered to use another antiprotozoal from another class and with a different mechanism of action.

Special precautions for use

Special precautions for use in animals:

Not applicable.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

People with known hypersensitivity to the active substance or any of the excipients should avoid contact with the veterinary medicinal product.

Avoid skin and eye contact with the product.

In case of accidental exposure to the skin or eyes, wash immediately with water.

Do not eat, drink or smoke while using the product.

Interactions with Other Medicaments

None known. In pigs there is no interaction in combination with iron supplementation.

Pregnancy and Lactation

Not applicable.

Side Effects

None known.

Symptoms and Treatment of Overdose

No signs of intolerance have been observed in healthy piglets with a threefold overdose.

Storage Condition

Store below 30°C.

Shelf life

3 years. After first opening the container, any product remaining after 24 hours should be discarded.

Withdrawal Period

Meat and offal: 77 days

Packing

200ml

TO BE PRESCRIBED AND TREATED WITH BY REGISTERED VETERINARY SURGEONS ONLY

Revised date:13.12.2022

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com